

**This Page Is Inserted by IFW Operations
and is not a part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- **BLACK BORDERS**
- **TEXT CUT OFF AT TOP, BOTTOM OR SIDES**
- **FADED TEXT**
- **ILLEGIBLE TEXT**
- **SKEWED/SLANTED IMAGES**
- **COLORED PHOTOS**
- **BLACK OR VERY BLACK AND WHITE DARK PHOTOS**
- **GRAY SCALE DOCUMENTS**

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

THIS PAGE BLANK (USPTO)

**(19) World Intellectual Property Organization
International Bureau**



(43) International Publication Date
3 January 2002 (03.01.2002)

(10) International Publication Number
WO 02/00860 A2

PCT

- (51) International Patent Classification⁷: C12N 9/00
MANNING, Gerard [IE/US]; 844 Fremont Street #4, Menlo Park, CA 94025 (US). CAENEPEEL, Sean [US/US]; 447 Orange Street #22, Oakland, CA 94610 (US). CHARYDCZAK, Glen [US/US]; 936 Sir Francis Drake Boulevard, Kentfield, CA 94904 (US).
- (21) International Application Number: PCT/US01/20171
- (22) International Filing Date: 26 June 2001 (26.06.2001).
- (25) Filing Language: English
(74) Agents: BURROUS, Beth, A. et al.; Foley & Lardner, Washington Harbour, Suite 500, 3000 K Street, N.W., Washington, DC 20007-5109 (US).
- (26) Publication Language: English
(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (30) Priority Data:
60/214,047 26 June 2000 (26.06.2000) US
- (71) Applicant (*for all designated States except US*): SUGEN, INC. [US/US]; 230 East Grand Avenue, South San Francisco, CA 94080-4811 (US).
- (72) Inventors; and
(75) Inventors/Applicants (*for US only*): PLOWMAN, Gregory [US/US]; 35 Winding Way, San Carlos, CA 94070 (US). WHYTE, David [US/US]; 2623 Barclay Way, Belmont, CA 94002 (US). SUDARSANAM, Sucha [US/US]; 20 Corte Patencio, Greenbrae, CA 94904 (US).
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

[Continued on next page]

- (54) Title:** NOVEL PROTEASES.

CCAGTCCAGGGGGAACGCAAGTCGCCCCGCTTCGGTGTACTTCCCTCAGCCTGTGGCCAGCGGACTTCAAG
GATAACTGGAGGATTGCGCGCTCCAGACAGGAAGTGGCCCTGGCAGGTGAGCCTGCAGACCGAACAGACAC
ATCTCGGAGAGCTCCCTTATTCGCGACAGACTGGGTATAAAGAGGACACCAACTCAGATTGTGGTGAGCCG
TGGTGTTCGGAGGATTGGAATGACCCGCCATTGGCCTGGGAGGTGAGCTCCGGATGGAAATGAGCAGT
TGTGTGGAGGGCCCTCACTGACCCAGCTGGGTGGTCTGCAGCTCGGCCACTGCAGCCAGGCCAACCAAGAGTA
CTCAGTGGTGCTTGGCACCTCCAAGCTGCAGCCCATGAATTCAGCAGGGCCCTCTGGGTCCCTGTGAGGGAC
ATCAATTATGCACCAAGTACTGGGGCGGGGCTTCATCACTGGGTGACGTTGCCCTTGTCCACTTCAACACCT
TGACCTTCAGTAGGACTCGTGACGCCATCTGCCCTCCGAGGCCCAATTCAACTGGAAGTTGGGACGAGTG
TTGGGTGACTGGCTGGAGCCAGGTTAAGCAGCGCTTTTCAGGCTCCACAGCCAATCTCATGCTGACCCGAGGC
TGACGAGGCTTGAGGTGTTTATCATGGACAACAACGAGGTGTACCGGCAATTACAAGAAGTCCCTCTCTCCCCCA
GTTGTCCCCCTTGCTCTGGGGACATGATCTGTGCCACCAATTATGGGGAAATCTGTGCTATGGGGATTCTGG
AGGGCCATTGGCTTGTGAAGTTAGGGCAGATGGATTCTGGCTGGGGTGTGTCTGGGAAAGGCTGCTGTG
AAGGCACAGATCCAGGTGTGATACCCCGCATACCAAAATACACCAATGGATCAAGAAGAACTGAGCAATGG
AGCCTCTCAGGTCCCTGTGCTCTGCCTGCCTCTGTTCTGTGCTGGCCGCTGCAGCCCGAGATGGGCTCCT
GA

>SGPR448_1_SEQID_25
ACCTCGCCACCGAGTGTGTGGCCGAACCCCTCTGAGAATCGTGGGAGGAGTGGACGCGGAGGAAGGGAGGTGG.
CCTCTGGCAGGTGAGCGGTGAGGACCAAAAGGCAGGCACATCTCGCGGGGCAACCTGGTCAACGCCACGTGGGTG
CTGACGGCAGGCCACATGCATTTCAGCGCGTTCCATTACAGTGTCAAGATGGGAGATCGGAGTGTCTATAATGAA
AACACAAGTGTGGTGGTCTCAGTCCAAAGAGCGCTTTGTCCACCTAAGTTCTCAACAGTTACACCACTCGAAAT
GACCTTGCCCTTCTCCAGCTCCAAACATCGTGTGAATTTACCTCAAACTCCAGGCTATCTGCATCCCTCAGGAG
AATTTCCAGGTGGAAAGTAGGACAGCGTCTGGGTGACCGGATGGGGCAAAACACGAGAAAGTGGAGAGAAAC
TGCACTAGAAATCTTCAGGATGTGGACCAATCATCATCTGCTGTAGGAATGTAAATGAAGTAATACAGAAAGGC
CTTGTCATCTACTAAGGATGTAATAATAAAAGGGATGGTCTGTGGCTATAAAGAAACAAGGAAAGGATTCTTGCAA
GGAGATTCTGGGGGGCCGCTTGCCCTGTGAATATAATGACACATGGGTCCAGGTAGGGATTGTGAGCTGGGGCA
TCCGGCTGTGGTGC

>SGPR447_SEQID_28
ATGGGCGCGCGCGGGCGCTGCTGCTGGCGCTGCTGCTGGCTCGGGCTGGACTCGGGAAGCCGGAGGCCTG
CGGACCACCGGGAATAACACGCGCTGGTGGCGGGCGGAGTGAGATCCGCGCGCGCGCTGCCATGCGCAGG
CCAGCCTCGCGCTGAGGAGACGCCACCGATGTGGAGGGAGCTGCTCAGCGCCGCTGGGTGGCTCTCGCGTG

- (57) Abstract:** The present invention relates to protease polypeptides, nucleotide sequences encoding the protease polypeptides, as well as various products and methods useful for the diagnosis and treatment of various protease-related diseases and conditions.

WO 02/00860 A2

**Published:**

- without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

What is claimed is:

CLAIMS

1. An isolated, enriched or purified nucleic acid molecule, wherein said nucleic acid molecule comprises a nucleotide sequence that:
- 5 (a) encodes a polypeptide having an amino acid sequence selected from the group consisting of those set forth in SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118 and biological domains thereof;
- 10 (b) is the complement of the nucleotide sequence of (a); or
- 15 (c) hybridizes under stringent conditions to the nucleotide molecule of (a) and encodes a protease polypeptide.
- 20
- 25
2. The nucleic acid molecule of claim 1, further comprising a vector or promoter operatively linked to the nucleotide sequence.

3. The nucleic acid molecule of claim 1, wherein said nucleic acid molecule is isolated, enriched, or purified from a mammal.
- 5 4. The nucleic acid molecule of claim 3, wherein said mammal is a human.
- 10 5. The nucleic acid molecule of claim 1 comprising a nucleic acid comprising a nucleotide sequence which hybridizes under stringent conditions to a nucleotide sequence encoding a protease polypeptide having an amino acid sequence selected from the group consisting of those set forth in SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118.
- 15
- 20
- 25

- 5
10
15
20
25
6. An isolated, enriched, or purified protease polypeptide, wherein said polypeptide comprises an amino acid sequence at least about 90% identical to a sequence selected from the group consisting of those set forth in SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118 and biological domains thereof.
 7. The protease polypeptide of claim 6, wherein said polypeptide is isolated, purified, or enriched from a mammal.
 8. The protease polypeptide of claim 7, wherein said mammal is a human.
 9. An antibody or antibody fragment having specific binding affinity to a protease polypeptide or to a domain of said polypeptide, wherein said polypeptide comprises an amino acid sequence selected from the

group consisting of those set forth in SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118.

10. A hybridoma which produces the antibody of Claim 9.
11. A kit comprising an antibody which binds to a polypeptide of claim 6 and a negative control antibody.
12. A method for identifying a substance that modulates the activity of a protease polypeptide comprising the steps of:
 - (a) contacting a protease polypeptide substantially identical to an amino acid sequence selected from the group consisting of those set forth in SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID

5 NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72,
SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID
NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79,
SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID
NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86,
SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID
NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93,
SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID
NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100,
10 SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID
NO:104, SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:107,
SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID
NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114,
SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ
15 ID NO:118 with a test substance;

(b) measuring the activity of said polypeptide; and

(c) determining whether said substance modulates the activity of
said polypeptide.

20 13. A method for identifying a substance that modulates the activity of a
protease polypeptide in a cell comprising the steps of:

(a) expressing a protease polypeptide in a cell, wherein said
polypeptide comprises a sequence substantially identical to an
amino acid sequence selected from the group consisting of
25 those set forth in SEQ ID NO:60, SEQ ID NO:61, SEQ ID
NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65,
SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID
NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72,
SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID

- 5 NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79,
SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID
NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86,
SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID
NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93,
SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID
NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100,
SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID
NO:104, SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:107,
10 SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID
NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114,
SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ
ID NO:118;
- 15 (b) adding a test substance to said cell; and
(c) monitoring a change in cell phenotype, cell proliferation, cell
differentiation or the interaction between said polypeptide and
a natural binding partner.
- 20 14. A method for treating a disease or disorder by administering to a
patient in need of such treatment a substance that modulates the
activity of a protease substantially identical to an amino acid
sequence selected from the group consisting of those set forth in SEQ
ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID
NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID
25 NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID
NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID
NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID
NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID
NO:84, SEQ ID NO:85, SEQ ID NO:86, SEQ ID NO:87, SEQ ID

5 NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118.

10 15. The method of claim 14, wherein said disease or disorder is selected from the group consisting of cancers, immune-related diseases and disorders, cardiovascular disease, brain or neuronal-associated diseases, metabolic disorders and inflammatory disorders.

15 16. The method of claim 15, wherein said disease or disorder is selected from the group consisting of cancers of tissues; cancers of blood or hematopoietic origin; cancers of the breast, colon, lung, prostate, cervical, brain, ovarian, bladder or kidney.

20 17. The method of claim 15, wherein said disease or disorder is selected from the group consisting of central or peripheral nervous system diseases, migraines; pain; sexual dysfunction; mood disorders; attention disorders; cognition disorders; hypotension; hypertension; psychotic disorders; neurological disorders and dyskinesias.

25 18. The method of claim 15, wherein said substance modulates protease activity *in vitro*.

19. The method of claim 18, wherein said substance is a protease inhibitor.
20. A method for detection of a protease polypeptide in a sample as a diagnostic tool for a disease or disorder, wherein said method comprises:
- (a) contacting said sample with a nucleic acid probe which hybridizes under hybridization assay conditions to a nucleic acid target region of a protease polypeptide having an amino acid sequence selected from the group consisting of those set forth in SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118, said probe comprising the nucleic acid sequence encoding the polypeptide, fragments thereof, and the complements of the sequences and fragments; and

(b) detecting the presence or amount of the probe:target region hybrid as an indication of the disease.

5 21. The method of claim 20, wherein said disease or disorder is selected from the group consisting of cancers, immune-related diseases and disorders, cardiovascular disease, brain or neuronal-associated diseases, metabolic disorders and inflammatory disorders.

10 22. The method of claim 21, wherein said disease or disorder is selected from the group consisting of cancers of tissues; cancers of hematopoietic; cancers of blood or hematopoietic origin; cancers of the breast, colon, lung, prostate, cervical, brain, ovarian, bladder or kidney.

15 23. The method of claim 21, wherein said disease or disorder is selected from the group consisting of central or peripheral nervous systems disease, migraines, pain; sexual dysfunction; mood disorders; attention disorders; cognition disorders; hypotension; hypertension; psychotic disorders; neurological disorders; and dyskinesias.

20 24. The isolated, enriched or purified nucleic acid molecule of claim 1 comprising a nucleic molecule encoding a biological domain of a protease polypeptide having a sequence selected from the group consisting of SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ
25 ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID

NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86, SEQ ID
 NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID
 NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID
 NO:95, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:98, SEQ ID
 5 NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ ID
 NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106, SEQ
 ID NO:107, SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110,
 SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID
 NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and
 10 SEQ ID NO:118.

25. The nucleic acid molecule of claim 1 comprising a nucleic acid
 sequence encoding a protease polypeptide having an amino acid
 sequence that has least 90 % identity to a polypeptide selected from
 15 the group consisting of those set forth in SEQ ID NO:60, SEQ ID
 NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID
 NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID
 NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID
 NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID
 20 NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID
 NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID
 NO:85, SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, SEQ ID
 NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID
 NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID
 25 NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID
 NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ
 ID NO:105, SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:108,
 SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:111, SEQ ID

NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118.

26. The nucleic acid molecule of Claim 1 wherein the molecule
5 comprises a nucleotide sequence substantially identical to a sequence
selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2,
SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID
NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID
NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID
10 NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID
NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID
NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID
NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID
NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID
15 NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID
NO:39, SEQ ID NO:40, SEQ ID NO:41, SEQ ID NO:42, SEQ ID
NO:43, SEQ ID NO:44, SEQ ID NO:45, SEQ ID NO:46, SEQ ID
NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID
NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:54, SEQ ID
20 NO:55, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:58, and SEQ
ID NO:59.

27. An isolated, enriched or purified nucleic acid molecule consisting
essentially of about 10-30 contiguous nucleotide bases of a nucleic
25 acid sequence that encodes a polypeptide that is selected from the
group consisting of SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62,
SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66,
SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70,
SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74,

5 SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78,
SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82,
SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86,
SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90,
SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94,
SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:98,
SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102,
SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID
10 NO:106, SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:109, SEQ
ID NO:110, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113,
SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID
NO:117 and SEQ ID NO:118.

28. The isolated, enriched or purified nucleic acid molecule of Claim 27
15 consisting essentially of about 10-30 contiguous nucleotide bases of a
nucleic acid sequence selected from the group consisting of SEQ ID
NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5,
SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID
NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID
20 NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID
NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID
NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID
NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID
NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID
25 NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID
NO:38, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:41, SEQ ID
NO:42, SEQ ID NO:43, SEQ ID NO:44, SEQ ID NO:45, SEQ ID
NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID
NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID

NO:54, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, SEQ ID
NO:58, and SEQ ID NO:59.

29. A recombinant cell comprising the nucleic acid molecule of claim 1.

5

30. A method for detecting the presence or amount of protease
polypeptide in a sample comprising

10

(a) contacting the sample with the antibody of Claim 9 under
conditions suitable for protease-antibody immunocomplex
formation; and

(b) detecting the presence or amount of the antibody conjugated to
the protease polypeptide.

THIS PAGE BLANK (USPTO)

cDNA Sources: ESTs gi|13745759, 7472044CB1, 7474338CB1, gi|13703426, gi|5392427, gi|2142177, gi|2103202, LIB4218-103-R1-K1-H5, LIB4218-085-Q1-K1-C6, LIB4752-019-R1-K1-H4

5 SGPr455, SEQ ID NO:42, SEQ ID NO:101

Genewise orthologs: gi|7512178, gi|7512176.

Genomic DNA sources:90000641321557

cDNA Sources: Incyte template 987279.1.

10 SGPr507_2, SEQ ID NO:43, SEQ ID NO:102

Genewise orthologs: gi|13385812, gi|12854692, gi|2499862.

Genomic DNA sources:90000642611957

SGPr559, SEQ ID NO:44, SEQ ID NO:103

15 Genewise orthologs: XP_016993, BAB20079

Genomic DNA sources:Celera_asm5h 335001064013332.

cDNA Sources:gi|13173471.

SGPr567_1, SEQ ID NO:45, SEQ ID NO:104

20 Genewise orthologs: NP_114435, Q9JIQ8

Genomic DNA sources:Celera_asm5h 90000642045213

cDNA Sources:gi|14042983|ref|NM_032046.1.

SGPr479_1, SEQ ID NO:46, SEQ ID NO:105

25 Genewise orthologs: NP_114154, NP_038949, NP_033382

Genomic DNA sources:90000624931837

cDNA Sources:EST gi|13997890 and Incyte EST 7480124CB1,.

SGPr489_1, SEQ ID NO:47, SEQ ID NO:106

THIS PAGE BLANK (USPTO)

SGPr520, SEQ ID NO:41, SEQ ID NO:100 is 1737 nucleotides long. The open reading frame starts at position 1 and ends at position 1737, giving an ORF length of 1737 nucleotides. The predicted protein is 578 amino acids long. This sequence codes for a full length protein. It is classified as (superfamily/group/family):

- 5 Protease, Serine, Trypsin. The cytogenetic position of this gene is 2q37. This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: none.

SGPr455, SEQ ID NO:42, SEQ ID NO:101 is 2913 nucleotides long. The open reading frame starts at position 1 and ends at position 2913, giving an ORF length of 2913 nucleotides. The predicted protein is 970 amino acids long. This sequence codes for a full length protein. It is classified as (superfamily/group/family):

- 10 Protease, Serine, Trypsin. The cytogenetic position of this gene is 12p11.2. This sequence is represented in the database of public ESTs (dbEST) by the following
15 ESTs: AW450155, AW995496.

SGPr507_2, SEQ ID NO:43, SEQ ID NO:102 is 798 nucleotides long. The open reading frame starts at position 1 and ends at position 798, giving an ORF length of 798 nucleotides. The predicted protein is 265 amino acids long. This sequence
20 codes for a full length protein. It is classified as (superfamily/group/family):
Protease, Serine, Trypsin. The cytogenetic position of this gene is 7q36. This sequence is represented in the database of public ESTs (dbEST) by the following
ESTs: BG217724, BG219738, BG192709.

- 25 SGPr559, SEQ ID NO:44, SEQ ID NO:103 is 1365 nucleotides long. The open reading frame starts at position 1 and ends at position 1365, giving an ORF length of 1365 nucleotides. The predicted protein is 454 amino acids long. This sequence codes for a full length protein. It is classified as (superfamily/group/family):
Protease, Serine, Trypsin. The cytogenetic position of this gene is 21q22. This

THIS PAGE BLANK (USPTO)

sequence is represented in the database of public ESTs (dbEST) by the following ESTs: AI978874, AI469095, BF435670

5 SGPr567_1, SEQ ID NO:45, SEQ ID NO:104 is 1614 nucleotides long. The open reading frame starts at position 1 and ends at position 1614, giving an ORF length of 1614 nucleotides. The predicted protein is 537 amino acids long. This sequence codes for a full length protein. It is classified as (superfamily/group/family):
Protease, Serine, Trypsin. The cytogenetic position of this gene is 11q23. This sequence is represented in the database of public ESTs (dbEST) by the following
10 ESTs: BE732381, R78581, AW845106.

15 SGPr479_1, SEQ ID NO:46, SEQ ID NO:105 is 981 nucleotides long. The open reading frame starts at position 1 and ends at position 981, giving an ORF length of 981 nucleotides. The predicted protein is 326 amino acids long. This sequence codes for a full length protein. It is classified as (superfamily/group/family):
Protease, Serine, Trypsin. The cytogenetic position of this gene is 1q42. This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: BG718703, AA401705, AA398170. The nucleic acid contains short
20 repetitive sequence (the position and sequence of the repeat): 780
tggaattgtgagctggggccg 800.

25 SGPr489_1, SEQ ID NO:47, SEQ ID NO:106 is 1671 nucleotides long. The open reading frame starts at position 1 and ends at position 1671, giving an ORF length of 1671 nucleotides. The predicted protein is 556 amino acids long. This sequence codes for a full length catalytic domain. It is classified as
(superfamily/group/family): Protease, Serine, Trypsin. The cytogenetic position of this gene is 11p15. This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: AW271430, AW237893.

THIS PAGE BLANK (USPTO)

domain(s) in this protein match the hidden Markov profile for a trypsin (PF00089), from amino acid 42 to amino acid 135. The positions within the HMMR profile that match the protein sequence are from profile position 35 to profile position 148. Other domains identified within this protein are: Trypsin domain from amino acid

5 247 to 258. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 2.40E-121; number of identical amino acids = 195; percent identity = 73%; percent similarity = 81%; the accession number of the most similar entry in NRAA is

10 NP_080593.1; the name or description, and species, of the most similar protein in NRAA is: RIKEN cDNA 1700016G05 gene [Mus musculus].

SGPr559, SEQ ID NO:44, SEQ ID NO:103 encodes a protein that is 454 amino acids long. It is classified as a Serine protease, of the trypsin family. The protease

15 domain(s) in this protein match the hidden Markov profile for a trypsin (PF00089), from amino acid 217 to amino acid 444. The positions within the HMMR profile that match the protein sequence are from profile position 1 to profile position 259. Other domains identified within this protein are: Low-density lipoprotein receptor domain class A (PF00057), from amino acid 71 to 109. LDL-receptors the class A

20 domains form the binding site for LDL and calcium. The acidic residues between the fourth and sixth cysteines are important for high-affinity binding of positively charged sequences in LDLR's ligands. The repeat has been shown to consist of a beta-hairpin structure followed by a series of beta turns (see <http://www.expasy.ch/cgi-bin/get-prodoc-entry?PDOC00929>). The results of a

25 Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 1.40E-288; number of identical amino acids = 454; percent identity = 100%; percent similarity = 100%; the accession number of the most similar entry in NRAA is NP_076927.1; the name or description, and

THIS PAGE BLANK (USPTO)

species, of the most similar protein in NRAA is: transmembrane protease, serine 3 [Homo sapiens]. This protein has a transmembrane domain from amino acid 49 to amino acid 71.

- 5 SGPr567_1, SEQ ID NO:45, SEQ ID NO:104 encodes a protein that is 537 amino acids long. It is classified as a Serine protease, of the trypsin family. The protease domain(s) in this protein match the hidden Markov profile for a trypsin (PF00089), from amino acid 296 to amino acid 524. The positions within the HMMR profile that match the protein sequence are from profile position 1 to profile position 259.
- 10 The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 1.70E-135; number of identical amino acids = 534; percent identity = 99%; percent similarity = 99%; the accession number of the most similar entry in NRAA is NP_114435.1; the name or
- 15 description, and species, of the most similar protein in NRAA is: mosaic serine protease [Homo sapiens].

- SGPr479_1, SEQ ID NO:46, SEQ ID NO:105 encodes a protein that is 326 amino acids long. It is classified as a Serine protease, of the trypsin family. The protease domain(s) in this protein match the hidden Markov profile for a trypsin (PF00089), from amino acid 60 to amino acid 288. The positions within the HMMR profile that match the protein sequence are from profile position 1 to profile position. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein
- 20 sequence yielded the following results: Pscore = 1.70E-39; number of identical amino acids = 107; percent identity = 42%; percent similarity = 57%; the accession number of the most similar entry in NRAA is NP_114154.1; the name or
- 25 description, and species, of the most similar protein in NRAA is: marapsin [Homo sapiens].

THIS PAGE BLANK (USPTO)

FIG. 1NN

>SGPR559_SEQID_44
ATGGGGGAAATGATCCGCTGCTGTGAAGCCCCCTTCTCATCCGATCGCTTTTGGCCCTGATGATTTGAAA
ATAAGTCTGTGACACAGATGCAGATGCTGTGCTGCACAGATCCTGTCACTGCTGCCATTGAAGTTTTTTC
ATCATCGTCATTGGGATCATTGCATTGATATTAGCACTGGCCATTGGTCTGGGCATCCACTTCGACTGCTCAGGG
AAGTACAGATGTCGCTCATCTTTAAGTGTATCGAGCTGATAGCTCGATGTGACGGAGTCTCGGATTGCAAGAC
GGGAGGACGAGTACCGCTGTGTCGGGTGGGTGTCAGAAATGCCGTGCTCCAGGTGTTCCAGAGTCTGCTTCGT
GGAAGACCATGTCTCCGATGACTGGAAGGTCACTACGCAATGTTGCCGTGTGCCAACTGGGTTTCCCAAGC
TATGTGAGTTCAGATAACCTCAGAGTGAGCTCGCTGGAGGGCAGTTCCGGGAGGAGTTTGTGTCCATCGATCA
CCTCTTGCCAGATGACAAGGTGACTGCATTACACCACCTCAGTATATGTAGGGAGGGATGTGCTCTGCCCACG
TGGTTACCTTGCAGTGCACAGCCTGTGTGTCATAGAAAGGGTACAGCTACGCATCGTGGTGGAAACATGTCC
TTGCTCTCGAGTGGCCCTGGCAGGCCAGCCTTCAGTCCAGGGCTACCACCTGTGCGGGGCTCTGTCTATCA
CGCCCTGTGGATCATCACTGCTGCACACTGTGTTTATGACTTGTACCTCCCAAGTCAATGGACCATCCAGGTGG
GTCTAGTTTCCCTGTTGGACAATCCAGCCCATCCACCTTGGTGGAGAAGATTGCTACCACAGCAAGTACAAGC
CAAAGAGGCTGGCAATGACATCGCCCTTATGAAGCTGGCCGGCCACTCAGCTTCAATGAAATGATCCAGCCT
GTGTCCTGCCCCAATCTGAAGAGAACTTCCCGATGGAAGTGTGCTGGACGTCAAGATGGGGGCCACAG
AGGATGGAGCAGGTGACGCTCCCTGTCTGAACCAACGCGGCTCCCTTTGATTTCCAAACAAGATCTGCAAC
CACAGGGACGTGTACGGTGGCATCATCTCCCTCCATGCTCTGCGGGGCTACCTGACGGGTGGCGTGGACA
GCTGCCAGGGGACAGCGGGGGCCCTGTTGTGTCAGAGAGGAGGCTGTGGAGTTAGTGGAGCGGACCA
GCTTTGGCATCGGCTGCGCAGAGGTGAACAAGCCTGGGGTGTACACCCGTGTACCTTCTTCTTCTGGACTGGAT
CCACGAGCAGATGGAGAGAGACCTTAAAAACCTGA

>SGPR567_1_SEQID_45
ATGGAGAGGACAGCCACGGGAATGCATCTCCAGCAAGAACAACCTTCAGCTGGAGCATCTCCAGCCCAGGCAT
CTCCAGCTGGACACCTCCAGGCGGGCATCTCCAGCCAGGCATCTCCAGCCCAGGCATCTCCAGCTGGGAC
ACCTCCGGGCGGGCATCTCCAGCCCAGGCATCTCCAGCTGGTACACCTCCAGCCGGGCATCTCCAGGCCG
GGCATCTCCAGCCCAGGCATCTCCAGCCCAGGCATCTCCAGCCCAGGCATCTCCAGCCCAGGCATCTCCGGCT
CTGGCATCACTTTCAGGTCTCATCCGGCAGGTCAATCCGCCAGGTCAAGCTCGGTGACAACTCCCCAAC
CAGAGTGTACCTTGTAGAGCAACACCAGTGGGGCTGTACCCATCCGATCATCTCTGCCAGGTCAAGCACCA
CAACCAGGGCCACAGGAGAGCCAGTCCAGTCTGGCAGGGCCACACAGGATCAGGTACAAGGAGCAGA
GGGAGAGCTGTCCCAAGCACGCTGTTCCGTGTGACGGGGTGGTGGACTGCAAGCTGAAGAGTGACGAGCTGG
GCTGCGTGAGGTTTACTGGGACAACTCTGCTTAAATCTACTCTGGTCTCTCCCATCAGTGGCTTCCCATCT
GTAGCAGCACTGGAATGACTCTACTCAGAGAAGACCTGCCAGCAGCTGGGTTTCGAGAGTGTCAACCGGACA

THIS PAGE BLANK (USPTO)

FIG. 20

LRNAQELLGPRRLRALPALPALRESPLHPARELRHLHSGSRAAGTRFPPKRRPEPRGEANGCPGLEPLRQKLA
ALQGAHAWILQVPSEHLAMNFHEVLADLGSKTLTGLFRAWVRAGLGGRHVAFSGLVGLEPATLARSLPRLVQALQA
FRVAALAEGEPEGPWMDVGQGPGLERKGGHPLNPQVPPARQP

>SGP455_SEQID_101
MSPDIALLYLKHVKFGNAVQPICLPDSDDKVEPGILCLSSGWGKISKTSEYSNVLQEMELPIMDDRACNTVLKSMNL
PLGRITMLCAGFPDWGMDACQDGGPLVCRGGGIWILAGITSWAGCAGSVPRNNHVKASLGIFSKVSELMDF
ITQNLFTGLDRGQPLSKVGSRYITKALSSVQEVNGSQDKILKFTSLDMEKQVGCDDHDYVLSRSSGVLFSKVCCKIL
PSPLLAETSEAMVPFVSDTEDSGSGFELTVAVQKSEAGSGCGLAILVEEGTNHSAKYPDLYPSNTRCHWFICAPEK
HIIKLTFEDFAVKFSPNCIYDAVVIYGDSEKHLAKL CGMLTITSSNNMTVIYFKSDGKNRLQGFKAFTILPSESLN
KFEKLPQNNPVSTV KAILHDVCGIPPFSPQWLSRRRAGGEEACPHCWQVGLRFLGDYQCGGAIINPVWILTAH
CVQLKNNPLSWTIIAGDHRNLKESTEQVRRRAKHIIVHEDFNTLSYDSDIALIQLSSPLEYNSVVRPVCLPHSAEPLFSS
EICAVTGWGSISAEISLVSSLDGGLASRLQQIQVHVLEREVEHTYYSAPHPGGITEKMICAGFAASGEKDFCQGDG
GPLVCRHENGPFVLYGIVSWGAGCVQWPKGVFARVMIFLDWIQSKINGKLFNSVIKTTITFFRVGLGTVSCCSEAELE
KPRGFFPTPRYLLDYRGRLECSWLVRSASSMAKFTIYLSLGGSPVCQDSVLIYEERHSKRKTAGGLHGRRLYSMT
FMSGPLVRVTFHALVRGAFGISYIVLVLPKPKDSKITRLSQSSNREHLVPCEDVLLTKPEGIMQIPRNSHRTTMCQW
RLVAPLNHIIQLNIINFPMKPTTTFVCHGHLRVYEGFGPGKLGRLMLMSTELSWFLSQFSTKTKTTASCGETAVSMKMMY
TSIFLALQNTCYHALPHEVLRK

>SGP507_2_SEQID_102
MKYVYLVGLAGTFFADSSVQKEDPAPYLVLKSHFNPCVGLKPSWWLAPAHCYLPNLKVMLGNFKSRVRDGT
QTINPIQVRYWNYSHSAPQDDMLIKLAKPAMLPKVQPLTLATTNVRPGTVCLLSGLDWSQENSGLWQLEPPGHLT
LHRGPAIPDWQRHNSHEQGRHPDLRQNLEAPVMSDRECQTEQKGKSHRNSLCVKFVKVFSRIFGEVAVATVICKDL
QGIEVGHFMGGDVGIYTNVYKYSWIENTAKDK

>SGP559_SEQID_103
MGENDPPAVEAPFSFRSLFGLDDLKISPVPADADAVAAQILSLLPLKFFPIVIGIALLALGIGHFDCSGKYRCRSSF
KCIELIARCDGVSDCKDGEDEYRCVVRGGQNAVQLQVFTAASWKTMCDDDWKGYANVACAQLGPPSYVSSDNLRV
SLEGGFREEFVSIHLLPDDKV TALHHSVYVREGCASGHVTLQCTACGHRRGYSSRVGGNMSLLSQWPWQASLQ
FQGYHLCGGSVITPLWIITAAHCVYDLYLPKSWTIQVGLVSLDNPAPSHLVEKIVYHSKYKPKRLGNDIALMKLAGPLT
FNEMIQVCLPNSEENFPDGKVCWTSWGATEDGAGDASPVLNHAAVPLISNKNHHRDVGGIISPSMLCAGYLTG
GVDSCQGDGSGGPLVCQERRLWKLVGATSFGIGCAEVNKPGVYTRVTSFLDWIHEQMERDLKT

THIS PAGE BLANK (USPTO)

5

10

NOVEL PROTEASES

The present invention claims priority to provisional application serial no. 60/214,047 filed June 26, 2000, which is hereby incorporated by reference in its entirety.

15

FIELD OF THE INVENTION

The present invention relates to protease polypeptides, nucleotide sequences encoding the protease polypeptides, as well as various products and methods useful for the diagnosis and treatment of various protease-related diseases and conditions.

20

BACKGROUND OF THE INVENTION

Proteases and Human Disease

"Protease," "proteinase," and "peptidase" are synonymous terms applying to all enzymes that hydrolyse peptide bonds, *i.e.* proteolytic enzymes. Proteases are an exceptionally important group of enzymes in medical research and biotechnology. They are necessary for the survival of all living creatures, and are encoded by 1-2% of all mammalian genes. Rawlings and Barrett (MEROPS: the peptidase database. *Nucleic Acids Res.*, 1999, 27:325-331) (<http://www.babraham.co.uk/Merops/Merops.htm> (Which is incorporated herein by reference in its entirety including any figures, tables, or drawings.) have classified peptidases into 157 families based on

structural similarity at the catalytic core sequence. These families are further classed into 26 clans, based on indications of common evolutionary relationship. Peptidases play key roles in both the normal physiology and disease-related pathways in mammalian cells. Examples include the modulation of apoptosis (caspases), control of blood pressure (renin, angiotensin-converting enzymes), tissue remodeling and tumor invasion (collagenase), the development of Alzheimer's Disease (β -secretase), protein turnover and cell-cycle regulation (proteasome), and inflammation (TNF- α convertase). (Barrett *et al.*, Handbook of Proteolytic Enzymes, 1998, Academic Press, San Diego which is incorporated herein by reference in its entirety including any figures, tables, or drawings.)

Peptidases are classed as either exopeptidases or endopeptidases. The exopeptidases act only near the ends of polypeptide chains: aminopeptidases act at the free N-terminus and carboxypeptidases at the free C-terminus. The endopeptidases are divided, on the basis of their mechanism of action, into six subclasses: aspartyl endopeptidases (3.4.23), cysteine endopeptidases (3.4.22), metalloendopeptidases (3.4.24), serine endopeptidases (3.4.21), threonine endopeptidases (3.4.25), and a final group that could not be assigned to any of the above classes (3.4.99). (Enzyme nomenclature and numbering are based on "Recommendations of the Nomenclature Committee of the International Union of Biochemistry and Molecular Biology (NC-IUBMB) 1992, (<http://www.chem.qmw.ac.uk/iubmb/enzyme/EC34/intro.html>).)

In serine-, threonine- and cysteine-type peptidases, the catalytic nucleophile is the reactive group of an amino acid side chain, either a hydroxyl group (serine- and threonine-type peptidases) or a sulfhydryl group (cysteine-type peptidases). In aspartic-type and metallopeptidases, the nucleophile is commonly an activated water molecule. In aspartic-type peptidases, the water molecule is directly bound by the side chains of aspartate residues. In metallopeptidases, one or two metal ions hold the water molecule in place, and charged amino acid side chains are ligands for the metal ions. The metal may be zinc, cobalt or manganese. One metal ion is usually

attached to three amino acid ligands. Families of peptidases are referred to by use of the numbering system of Rawlings & Barrett (Rawlings, N. D. & Barrett, A. J. MEROPS: the peptidase database. *Nucleic Acids Research* 27 (1999) 325-331, which is incorporated herein by reference in its entirety including any figures, tables, or drawings).). Enzyme nomenclature and numbering are based on "Recommendations of the Nomenclature Committee of the International Union of Biochemistry and Molecular Biology (NC-IUBMB) 1992, (<http://www.chem.qmw.ac.uk/iubmb/enzyme/EC34/intro.html>).

10

Protease Families

1. Aspartyl proteases (Prosit number PS00141)

Aspartyl proteases, also known as acid proteases, are a widely distributed family of proteolytic enzymes in vertebrates, fungi, plants, retroviruses and some plant viruses. Aspartate proteases of eukaryotes are monomeric enzymes which consist of two domains. Each domain contains an active site centered on a catalytic aspartyl residue. The two domains most probably evolved from the duplication of an ancestral gene encoding a primordial domain. Enzymes in this class include cathepsin E, renin, presenilin (PS1), and the APP secretases.

20

2. Cysteine proteases (Prosit PDOC00126)

Eukaryotic cysteine proteases are a family of proteolytic enzymes which contain an active site cysteine. Catalysis proceeds through a thioester intermediate and is facilitated by a nearby histidine side chain; an asparagine completes the essential catalytic triad. Peptidases in this family with important roles in disease include the caspases, calpain, hedgehog, ubiquitin hydrolases, and papain.

25

3. Metalloproteases (Prosit PDOC00129)

The metalloproteases are a class which includes matrix metalloproteases (MMPs), collagenase, stromelysin, gelatinase, neprylisin, carboxypeptidase, dipeptidase, and membrane-associated metalloproteases, such as those of the ADAM family. They require a metal co-factor for activity; frequently the required metal ion is zinc but some metalloproteases utilize cobalt and manganese.

Proteins of the extracellular matrix interact directly with cell surface receptors thereby initiating signal transduction pathways and modulating those triggered by growth factors, some of which may require binding to the extracellular matrix for optimal activity. Therefore the extracellular matrix has a profound effect on the cells encased by it and adjacent to it. Remodeling of the extracellular matrix requires protease of several families, including metalloproteases (MMPs).

4. Serine proteases (S1) (Prosite PS00134 trypsin-his; PS00135 trypsin-ser)

The catalytic activity of the serine proteases from the trypsin family is provided by a charge relay system involving an aspartic acid residue hydrogen-bonded to a histidine, which itself is hydrogen-bonded to a serine. The sequences in the vicinity of the active site serine and histidine residues are well conserved in this family of proteases. A partial list of proteases known to belong to this large and important family include: blood coagulation factors VII, IX, X, XI and XII; thrombin; plasminogen; complement components C1r, C1s, C2; complement factors B, D and I; complement-activating component of RA-reactive factor; elastases 1, 2, 3A, 3B (protease E); hepatocyte growth factor activator; glandular (tissue) kallikreins including EGF-binding protein types A, B, and C; NGF- γ hain, γ -renin, and prostate specific antigen (PSA); plasma kallikrein; mast cell proteases; myeloblastin (proteinase 3) (Wegener's autoantigen); plasminogen activators (urokinase-type, and tissue-type); and the trypsins I, II, III, and IV. These peptidases play key roles in coagulation, tumorigenesis, control of blood pressure, release of growth factors, and other roles.

5. Threonine peptidases (T1) – (Prosite PDOC00326/PDOC00668)

Threonine proteases are characterized by their use of a hydroxyl group of a threonine residue in the catalytic site of these enzymes. Only a few of these enzymes have been characterized thus far, such as the 20S proteasome from the archaeobacterium *Thermoplasma acidophilum* (Seemuller *et al.*, 1995, *Science*, 268:579-82, and chapter 167 of Barrett *et al.*, Handbook of Proteolytic Enzymes, 1998, Academic Press, San Diego).

SUMMARY OF THE INVENTION

10 This invention concerns the isolation and characterization of novel sequences of human proteases. These sequences are obtained via bioinformatics searching strategies on the predicted amino acid translations of new human genetic sequences. These sequences, now identified as proteases, are translated into polypeptides which are further characterized. Additionally, the nucleic acid sequences of these proteases
15 are used to obtain full-length cDNA clones of the proteases. The partial or complete sequences of these proteases are presented here, together with their classification, predicted or deduced protein structure.

Modulation of the activities of these proteases will prove useful therapeutically. Additionally, the presence or absence of these proteases or the
20 DNA sequence encoding them will prove useful in diagnosis or prognosis of a variety of diseases. In this regard, Example 8 describes the chromosomal localization of proteases of the present invention, and describes diseases mapping to the chromosomal locations of the proteases of the invention.

A first aspect of the invention features an identified, isolated, enriched, or
25 purified nucleic acid molecule having an amino acid sequence selected from the group consisting of those set forth in SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID

NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID
 NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID
 NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86, SEQ ID
 NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID
 5 NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID
 NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ
 ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106,
 SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID
 NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115,
 10 SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118 and biological domains
 thereof.

The term "identified" in reference to a nucleic acid is meant that a sequence
 was selected from a genomic, EST, or cDNA sequence database based on being
 predicted to encode a portion of a previously unknown or novel protease.

15 By "isolated" in reference to nucleic acid is meant a polymer of 10
 (preferably 21, more preferably 39, most preferably 75) or more nucleotides
 conjugated to each other, including DNA and RNA that is isolated from a natural
 source or that is synthesized as the sense or complementary antisense strand. In
 certain embodiments of the invention, longer nucleic acids are preferred, for
 20 example those of 300, 600, 900, 1200, 1500, or more nucleotides and/or those
 having at least 50%, 60%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%,
 97%, 98% or 99% identity to a sequence selected from the group consisting of those
 set forth in SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID
 NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10,
 25 SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15,
 SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20,
 SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25,
 SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30,
 SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35,

SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:40,
SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:44, SEQ ID NO:45,
SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50,
SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:55,
5 SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:58, and SEQ ID NO:59.

It is understood that by nucleic acid it is meant, without limitation, DNA, RNA or cDNA, and where the nucleic acid is RNA, the thymine (T) will be uracil (U).

The isolated nucleic acid of the present invention is unique in the sense that
10 it is not found in a pure or separated state in nature. Use of the term "isolated" indicates that a naturally occurring sequence has been removed from its normal cellular (*i.e.*, chromosomal) environment. Thus, the sequence may be in a cell-free solution or placed in a different cellular environment. The term does not imply that the sequence is the only nucleotide chain present, but that it is essentially free
15 (preferably about 90% pure, more preferably at least about 95% pure) of non-nucleotide material naturally associated with it, and thus is distinguished from isolated chromosomes.

By the use of the term "enriched" in reference to nucleic acid is meant that the specific DNA or RNA sequence constitutes a significantly higher fraction (2- to
20 5-fold) of the total DNA or RNA present in the cells or solution of interest than in normal or diseased cells or in the cells from which the sequence was taken. This could be caused by a person by preferential reduction in the amount of other DNA or RNA present, or by a preferential increase in the amount of the specific DNA or RNA sequence, or by a combination of the two. However, it should be noted that
25 enriched does not imply that there are no other DNA or RNA sequences present, just that the relative amount of the sequence of interest has been significantly increased. The term "significant" is used to indicate that the level of increase is useful to the person making such an increase, and generally means an increase relative to other nucleic acids of about at least 2-fold, more preferably at least 5-fold, more

preferably at least 10-fold or even more. The term also does not imply that there is no DNA or RNA from other sources. The DNA from other sources may, for example, comprise DNA from a yeast or bacterial genome, or a cloning vector such as pUC19. This term distinguishes from naturally occurring events, such as viral
5 infection, or tumor-type growths, in which the level of one mRNA may be naturally increased relative to other species of mRNA. That is, the term is meant to cover only those situations in which a person has intervened to elevate the proportion of the desired nucleic acid.

It is also advantageous for some purposes that a nucleotide sequence be in
10 purified form. The term "purified" in reference to nucleic acid does not require absolute purity (such as a homogeneous preparation). Instead, it represents an indication that the sequence is relatively more pure than in the natural environment (compared to the natural level this level should be at least 2- to 5-fold greater, *e.g.*, in terms of mg/mL). Individual clones isolated from a cDNA library may be
15 purified to electrophoretic homogeneity. The claimed DNA molecules obtained from these clones could be obtained directly from total DNA or from total RNA. The cDNA clones are not naturally occurring, but rather are preferably obtained via manipulation of a partially purified naturally occurring substance (messenger RNA). The construction of a cDNA library from mRNA involves the creation of a synthetic
20 substance (cDNA) and pure individual cDNA clones can be isolated from the synthetic library by clonal selection of the cells carrying the cDNA library. Thus, the process which includes the construction of a cDNA library from mRNA and isolation of distinct cDNA clones yields an approximately 10^6 -fold purification of the native message. Thus, purification of at least one order of magnitude, preferably
25 two or three orders, and more preferably four or five orders of magnitude is expressly contemplated.

By a "protease polypeptide" is meant 32 (preferably 40, more preferably 45, most preferably 55) or more contiguous amino acids in a polypeptide having an amino acid sequence selected from the group consisting of those set forth in SEQ ID

- NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118 and biological domains thereof. In certain aspects, polypeptides of 100, 200, 300, 400, 450, 500, 550, 600, 700, 800, 900 or more amino acids are preferred.
- The protease polypeptide can be encoded by a full-length nucleic acid sequence or any portion of the full-length nucleic acid sequence, so long as a functional activity of the polypeptide is retained. It is well known in the art that due to the degeneracy of the genetic code numerous different nucleic acid sequences can code for the same amino acid sequence. Equally, it is also well known in the art that conservative changes in amino acid can be made to arrive at a protein or polypeptide which retains the functionality of the original. Such substitutions may include the replacement of an amino acid by a residue having similar physicochemical properties, such as substituting one aliphatic residue (Ile, Val, Leu or Ala) for another, or substitution between basic residues Lys and Arg, acidic residues Glu and Asp, amide residues Gln and Asn, hydroxyl residues Ser and Tyr, or aromatic residues Phe and Tyr. Further information regarding making amino acid exchanges which have only slight, if any, effects on the overall protein can be found in Bowie *et al.*, *Science*, 1990, 247:1306-1310, which is incorporated herein by reference in its

entirety including any figures, tables, or drawings. In all cases, all permutations are intended to be covered by this disclosure.

The amino acid sequence of the protease peptide of the invention will be substantially similar to a sequence having an amino acid sequence selected from the group consisting of those set forth in SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118, or the corresponding full-length amino acid sequence, or fragments thereof.

A sequence that is substantially similar to a sequence selected from the group consisting of those set forth in SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106,

SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118 will preferably have at least 50%, 60%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identity to a sequence selected from the group consisting of SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118. Preferably the protease polypeptide will have at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identity to one of the aforementioned sequences.

By "identity" is meant a property of sequences that measures their similarity or relationship. Identity is measured by dividing the number of identical residues by the total number of residues and gaps and multiplying the product by 100. "Gaps" are spaces in an alignment that are the result of additions or deletions of amino acids. Thus, two copies of exactly the same sequence have 100% identity, but sequences that are less highly conserved, and have deletions, additions, or replacements, may have a lower degree of identity. Those skilled in the art will recognize that several computer programs are available for determining sequence identity using standard parameters, for example Gapped BLAST or PSI-BLAST (Altschul, *et al.* (1997) *Nucleic Acids Res.* 25:3389-3402), BLAST (Altschul, *et al.*

(1990) *J. Mol. Biol.* 215:403-410), and Smith-Waterman (Smith, *et al.* (1981) *J. Mol. Biol.* 147:195-197). Preferably, the default settings of these programs will be employed, but those skilled in the art recognize whether these settings need to be changed and know how to make the changes.

5 “Similarity” is measured by dividing the number of identical residues plus the number of conservatively substituted residues (see Bowie, *et al. Science*, 1999), 247:1306-1310, which is incorporated herein by reference in its entirety, including any drawings, figures, or tables) by the total number of residues and gaps and multiplying the product by 100.

10 In preferred embodiments, the invention features isolated, enriched, or purified nucleic acid molecules encoding a protease polypeptide comprising a nucleotide sequence that: (a) encodes a polypeptide having an amino acid sequence selected from the group consisting of those set forth in SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118 and biological domains thereof; (b) is the complement of the nucleotide sequence of (a); or (c) hybridizes under highly stringent conditions to the nucleotide molecule of (a) and encodes a naturally occurring protease polypeptide.

In preferred embodiments, the invention features isolated, enriched or purified nucleic acid molecules comprising a nucleotide sequence substantially identical to a sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:44, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:58, and SEQ ID NO:59. Preferably the sequence has at least 50%, 60%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or 100% identity to the above listed sequences.

The term "complement" refers to two nucleotides that can form multiple favorable interactions with one another. For example, adenine is complementary to thymine as they can form two hydrogen bonds. Similarly, guanine and cytosine are complementary since they can form three hydrogen bonds. A nucleotide sequence is the complement of another nucleotide sequence if all of the nucleotides of the first sequence are complementary to all of the nucleotides of the second sequence.

Various low or high stringency hybridization conditions may be used depending upon the specificity and selectivity desired. These conditions are well known to those skilled in the art. Under stringent hybridization conditions only highly complementary nucleic acid sequences hybridize. Preferably, such conditions prevent hybridization of nucleic acids having more than 1 or 2 mismatches out of 20 contiguous nucleotides, more preferably, such conditions

prevent hybridization of nucleic acids having more than 1 or 2 mismatches out of 50 contiguous nucleotides, most preferably, such conditions prevent hybridization of nucleic acids having more than 1 or 2 mismatches out of 100 contiguous nucleotides. In some instances, the conditions may prevent hybridization of nucleic acids having more than 5 mismatches in the full-length sequence.

By stringent hybridization assay conditions is meant hybridization assay conditions at least as stringent as the following: hybridization in 50% formamide, 5X SSC, 50 mM NaH_2PO_4 , pH 6.8, 0.5% SDS, 0.1 mg/mL sonicated salmon sperm DNA, and 5X Denhardt's solution at 42 °C overnight; washing with 2X SSC, 0.1% SDS at 45 °C; and washing with 0.2X SSC, 0.1% SDS at 45 °C. Under some of the most stringent hybridization assay conditions, the second wash can be done with 0.1X SSC at a temperature up to 70 °C (Berger *et al.* (1987) Guide to Molecular Cloning Techniques pg 421, hereby incorporated by reference herein in its entirety including any figures, tables, or drawings.). However, other applications may require the use of conditions falling between these sets of conditions. Methods of determining the conditions required to achieve desired hybridizations are well known to those with ordinary skill in the art, and are based on several factors, including but not limited to, the sequences to be hybridized and the samples to be tested. Washing conditions of lower stringency frequently utilize a lower temperature during the washing steps, such as 65 °C, 60 °C, 55 °C, 50 °C, or 42 °C.

The term "activity" means that the polypeptide hydrolyzes peptide bonds.

The term "catalytic activity", as used herein, defines the rate at which a protease catalytic domain cleaves a substrate. Catalytic activity can be measured, for example, by determining the amount of a substrate cleaved as a function of time. Catalytic activity can be measured by methods of the invention by holding time constant and determining the concentration of a cleaved substrate after a fixed period of time. Cleavage of a substrate occurs at the active site of the protease. The active site is normally a cavity in which the substrate binds to the protease and is cleaved.

The term "biological domain" means a domain or region of the protease polypeptide which has catalytic activity or which binds to the substrate of the protease.

5 The term "substrate" as used herein refers to a polypeptide or protein which is cleaved by a protease of the invention. The term "cleaved" refers to the severing of a covalent bond between amino acid residues of the backbone of the polypeptide or protein.

10 The term "insert" as used herein refers to a portion of a protease that is absent from a close homolog. Inserts may or may not be the product alternative splicing of exons. Inserts can be identified by using a Smith-Waterman sequence alignment of the protein sequence against the non-redundant protein database, or by means of a multiple sequence alignment of homologous sequences using the DNASTar program Megalign (Preferably, the default settings of this program will be used, but those skilled in the art will recognize whether these settings need to be changed and know how to make the changes.). Inserts may play a functional role by presenting a new interface for protein-protein interactions, or by interfering with such interactions.

20 In other preferred embodiments, the invention features isolated, enriched, or purified nucleic acid molecules encoding protease polypeptides, further comprising a vector or promoter operably linked to the nucleotide sequence. The invention also features recombinant nucleic acid, preferably in a cell or an organism. The recombinant nucleic acid may contain a sequence selected from the group consisting of those set forth in SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ

ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:44, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:58, and SEQ ID NO:59, or a functional derivative thereof and a vector or a promoter operably linked to the nucleotide sequence. The recombinant nucleic acid can alternatively contain a transcriptional initiation region functional in a cell, a sequence complementary to an RNA sequence encoding a protease polypeptide and a transcriptional termination region functional in a cell. Specific vectors and host cell combinations are discussed herein.

The term "vector" relates to a single or double-stranded circular nucleic acid molecule that can be transfected into cells and replicated within or independently of a cell genome. A circular double-stranded nucleic acid molecule can be cut and thereby linearized upon treatment with restriction enzymes. An assortment of nucleic acid vectors, restriction enzymes, and the knowledge of the nucleotide sequences cut by restriction enzymes are readily available to those skilled in the art. A nucleic acid molecule encoding a protease can be inserted into a vector by cutting the vector with restriction enzymes and ligating the two pieces together.

An operable linkage is a linkage in which the regulatory DNA sequences and the DNA sequence sought to be expressed are connected in such a way as to permit gene sequence expression. The precise nature of the regulatory regions needed for gene sequence expression may vary from organism to organism, but shall in general include a promoter region which, in prokaryotes, contains both the promoter (which directs the initiation of RNA transcription) as well as the DNA sequences which, when transcribed into RNA, will signal synthesis initiation.

The term "transfecting" defines a number of methods to insert a nucleic acid vector or other nucleic acid molecules into a cellular organism. These methods involve a variety of techniques, such as treating the cells with high concentrations of

salt, an electric field, detergent, or DMSO to render the outer membrane or wall of the cells permeable to nucleic acid molecules of interest or use of various viral transduction strategies.

The term "promoter" as used herein, refers to nucleic acid sequence needed for gene sequence expression. Promoter regions vary from organism to organism, but are well known to persons skilled in the art for different organisms. For example, in prokaryotes, the promoter region contains both the promoter (which directs the initiation of RNA transcription) as well as the DNA sequences which, when transcribed into RNA, will signal synthesis initiation. Such regions will normally include those 5'-non-coding sequences involved with initiation of transcription and translation, such as the TATA box, capping sequence, CAAT sequence, and the like.

In preferred embodiments, the isolated nucleic acid comprises, consists essentially of, or consists of a nucleic acid sequence selected from the group consisting of those set forth in SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:44, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:58, and SEQ ID NO:59 which encodes an amino acid sequence selected from the group consisting of those set forth in SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID

NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID
 NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID
 NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID
 NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86, SEQ ID NO:87, SEQ ID
 5 NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID
 NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97, SEQ ID
 NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ
 ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:107,
 SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:111, SEQ ID
 10 NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116,
 SEQ ID NO:117 and SEQ ID NO:118; a functional derivative thereof, or at least 35,
 40, 45, 50, 60, 75, 100, 200, or 300 contiguous amino acids selected from the group
 consisting of those set forth in SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62,
 SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67,
 15 SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72,
 SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77,
 SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82,
 SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86, SEQ ID NO:87,
 SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92,
 20 SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97,
 SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID
 NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106,
 SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID
 NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115,
 25 SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118. The nucleic acid may be
 isolated from a natural source by cDNA cloning or by subtractive hybridization.
 The natural source may be mammalian, preferably human, blood, semen, or tissue,
 and the nucleic acid may be synthesized by the triester method or by using an
 automated DNA synthesizer.

The term "mammal" refers preferably to such organisms as mice, rats, rabbits, guinea pigs, sheep, and goats, more preferably to cats, dogs, monkeys, and apes, and most preferably to humans.

In yet other preferred embodiments, the nucleic acid is a conserved or unique region, for example those useful for: the design of hybridization probes to facilitate identification and cloning of additional polypeptides, the design of PCR probes to facilitate cloning of additional polypeptides, obtaining antibodies to polypeptide regions, and designing antisense oligonucleotides.

By "conserved nucleic acid regions", are meant regions present on two or more nucleic acids encoding a protease polypeptide, to which a particular nucleic acid sequence can hybridize under lower stringency conditions. Examples of lower stringency conditions suitable for screening for nucleic acid encoding protease polypeptides are provided in Wahl *et al. Meth. Enzym.* 152:399-407 (1987) and in Wahl *et al. Meth. Enzym.* 152:415-423 (1987), which are hereby incorporated by reference herein in its entirety, including any drawings, figures, or tables.

Preferably, conserved regions differ by no more than 5 out of 20 nucleotides, even more preferably 2 out of 20 nucleotides or most preferably 1 out of 20 nucleotides.

By "unique nucleic acid region" is meant a sequence present in a nucleic acid coding for a protease polypeptide that is not present in a sequence coding for any other naturally occurring polypeptide. Such regions preferably encode 32 (preferably 40, more preferably 45, most preferably 55) or more contiguous amino acids set forth in a full-length amino acid sequence selected from the group consisting of those set forth in SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92,

SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97,
 SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID
 NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106,
 SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID
 5 NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115,
 SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118 in a sample. The nucleic
 acid probe contains a nucleotide base sequence that will hybridize to the sequence
 selected from the group consisting of those set forth in SEQ ID NO:1, SEQ ID
 NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7,
 10 SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12,
 SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17,
 SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22,
 SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27,
 SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32,
 15 SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37,
 SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:41, SEQ ID NO:42,
 SEQ ID NO:43, SEQ ID NO:44, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47,
 SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52,
 SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57,
 20 SEQ ID NO:58, and SEQ ID NO:59, or a functional derivative thereof.

In preferred embodiments, the nucleic acid probe hybridizes to nucleic acid
 encoding at least 12, 32, 75, 90, 105, 120, 150, 200, 250, 300 or 350 contiguous
 amino acids of a full-length sequence selected from the group consisting of those set
 forth in SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID
 25 NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID
 NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID
 NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID
 NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID
 NO:84, SEQ ID NO:85, SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, SEQ ID

NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:108, 5 SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118, or a functional derivative thereof.

Methods for using the probes include detecting the presence or amount of protease RNA in a sample by contacting the sample with a nucleic acid probe under 10 conditions such that hybridization occurs and detecting the presence or amount of the probe bound to protease RNA. The nucleic acid duplex formed between the probe and a nucleic acid sequence coding for a protease polypeptide may be used in the identification of the sequence of the nucleic acid detected (Nelson *et al.*, in Nonisotopic DNA Probe Techniques, Academic Press, San Diego, Kricka, ed., p. 15 275, 1992, hereby incorporated by reference herein in its entirety, including any drawings, figures, or tables). Kits for performing such methods may be constructed to include a container means having disposed therein a nucleic acid probe.

Methods for using the probes also include using these probes to find the full-length clone of each of the predicted proteases by techniques known to one skilled in 20 the art. These clones will be useful for screening for small molecule compounds that inhibit the catalytic activity of the encoded protease with potential utility in treating cancers, immune-related diseases and disorders, cardiovascular disease, brain or neuronal-associated diseases, and metabolic disorders. More specifically disorders including cancers of tissues, blood, or hematopoietic origin, particularly those 25 involving breast, colon, lung, prostate, cervical, brain, ovarian, bladder, or kidney; central or peripheral nervous system diseases and conditions including migraine, pain, sexual dysfunction, mood disorders, attention disorders, cognition disorders, hypotension, and hypertension; psychotic and neurological disorders, including anxiety, schizophrenia, manic depression, delirium, dementia, severe mental

retardation and dyskinesias, such as Huntington's disease or Tourette's Syndrome; neurodegenerative diseases including Alzheimer's, Parkinson's, multiple sclerosis, and amyotrophic lateral sclerosis; viral or non-viral infections caused by HIV-1, HIV-2 or other viral- or prion-agents or fungal- or bacterial- organisms; metabolic disorders including Diabetes and obesity and their related syndromes, among others; cardiovascular disorders including reperfusion restenosis, coronary thrombosis, clotting disorders, unregulated cell growth disorders, atherosclerosis; ocular disease including glaucoma, retinopathy, and macular degeneration; inflammatory disorders including rheumatoid arthritis, chronic inflammatory bowel disease, chronic inflammatory pelvic disease, multiple sclerosis, asthma, osteoarthritis, psoriasis, atherosclerosis, rhinitis, autoimmunity, and organ transplant rejection.

In another aspect, the invention describes a recombinant cell or tissue comprising a nucleic acid molecule encoding a protease polypeptide having an amino acid sequence selected from the group consisting of those set forth in SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118. In such cells, the nucleic acid may be under the control of the genomic regulatory elements, or may be under the control of exogenous regulatory elements including an exogenous promoter. By "exogenous" it is meant a promoter that is not

normally coupled *in vivo* transcriptionally to the coding sequence for the protease polypeptides.

The polypeptide is preferably a fragment of the protein encoded by a full-length amino acid sequence selected from the group consisting of those set forth in

5 SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84,

10 SEQ ID NO:85, SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:108,

15 SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118. By "fragment," is meant an amino acid sequence present in a protease polypeptide. Preferably, such a sequence comprises at least 32, 45, 50, 60, 100, 200, or 300 contiguous amino acids of a full-length sequence selected from the

20 group consisting of those set forth in SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106,

25

SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118.

- In another aspect, the invention features an isolated, enriched, or purified protease polypeptide having a sequence substantially identical to an amino acid sequence selected from the group consisting of those set forth in SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118 and biological domains thereof. Preferable the polypeptide sequence has at least 50%, 60%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or 100% identity to the above listed sequences.

- By "isolated" in reference to a polypeptide is meant a polymer of 6 (preferably 12, more preferably 18, most preferably 25, 32, 40, or 50) or more amino acids conjugated to each other, including polypeptides that are isolated from a natural source or that are synthesized. In certain aspects longer polypeptides are preferred, such as those with 100, 200, 300, 400, 450, 500, 550, 600, 700, 800, 900 or more contiguous amino acids of a full-length sequence selected from the group consisting of those set forth in SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67,

SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72,
 SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77,
 SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82,
 SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86, SEQ ID NO:87,
 5 SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92,
 SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97,
 SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID
 NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106,
 SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID
 10 NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115,
 SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118, and/or those polypeptides
 having at least 50%, 60%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%,
 97%, 98% or 99% identity to a sequence selected from the group consisting of SEQ
 ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ
 15 ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ
 ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ
 ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ
 ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ
 ID NO:85, SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ
 20 ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ
 ID NO:95, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ
 ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104,
 SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:108, SEQ ID
 NO:109, SEQ ID NO:110, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113,
 25 SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ ID
 NO:118.

The isolated polypeptides of the present invention are unique in the sense
 that they are not found in a pure or separated state in nature. Use of the term
 "isolated" indicates that a naturally occurring sequence has been removed from its

normal cellular environment. Thus, the sequence may be in a cell-free solution or placed in a different cellular environment. The term does not imply that the sequence is the only amino acid chain present, but that it is essentially free (at least about 90% pure, more preferably at least about 95% pure or more) of non-amino acid-based material naturally associated with it.

By the use of the term "enriched" in reference to a polypeptide is meant that the specific amino acid sequence constitutes a significantly higher fraction (2- to 5-fold) of the total amino acid sequences present in the cells or solution of interest than in normal or diseased cells or in the cells from which the sequence was taken. This could be caused by a person by preferential reduction in the amount of other amino acid sequences present, or by a preferential increase in the amount of the specific amino acid sequence of interest, or by a combination of the two. However, it should be noted that enriched does not imply that there are no other amino acid sequences present, just that the relative amount of the sequence of interest has been significantly increased. The term significant here is used to indicate that the level of increase is useful to the person making such an increase, and generally means an increase relative to other amino acid sequences of about at least 2-fold, more preferably at least 5- to 10-fold or even more. The term also does not imply that there is no amino acid sequence from other sources. The other source of amino acid sequences may, for example, comprise amino acid sequence encoded by a yeast or bacterial genome, or a cloning vector such as pUC19. The term is meant to cover only those situations in which man has intervened to increase the proportion of the desired amino acid sequence.

It is also advantageous for some purposes that an amino acid sequence be in purified form. The term "purified" in reference to a polypeptide does not require absolute purity (such as a homogeneous preparation); instead, it represents an indication that the sequence is relatively purer than in the natural environment. Compared to the natural level this level should be at least 2-to 5-fold greater (e.g., in terms of mg/mL). Purification of at least one order of magnitude, preferably two or

three orders, and more preferably four or five orders of magnitude is expressly contemplated. The substance is preferably free of contamination at a functionally significant level, for example 90%, 95%, or 99% pure.

In preferred embodiments, the protease polypeptide is a fragment of the
5 protein encoded by a full-length amino acid sequence selected from the group
consisting of those set forth in SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62,
SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67,
SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72,
SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77,
10 SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82,
SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86, SEQ ID NO:87,
SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92,
SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97,
SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID
15 NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106,
SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID
NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115,
SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118. Preferably, the protease
polypeptide contains at least 32, 45, 50, 60, 100, 200, or 300 contiguous amino acids
20 of a full-length sequence selected from the group consisting of those set forth in
SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64,
SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69,
SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74,
SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79,
25 SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84,
SEQ ID NO:85, SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89,
SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94,
SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99,
SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID

NO:104, SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118, or a functional derivative thereof.

5 In preferred embodiments, the protease polypeptide comprises an amino acid sequence having an amino acid sequence selected from the group consisting of those set forth in SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118.

20 The polypeptide can be isolated from a natural source by methods well-known in the art. The natural source may be mammalian, preferably human, blood, semen, or tissue, and the polypeptide may be synthesized using an automated polypeptide synthesizer.

 In some embodiments the invention includes a recombinant protease polypeptide having (a) an amino acid sequence selected from the group consisting of those set forth in SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78,

SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83,
SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88,
SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93,
SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:98,
5 SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ ID
NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:107,
SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:111, SEQ ID
NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116,
SEQ ID NO:117 and SEQ ID NO:118. By "recombinant protease polypeptide" is
10 meant a polypeptide produced by recombinant DNA techniques such that it is
distinct from a naturally occurring polypeptide either in its location (*e.g.*, present in
a different cell or tissue than found in nature), purity or structure. Generally, such a
recombinant polypeptide will be present in a cell in an amount different from that
normally observed in nature.

15 The polypeptides to be expressed in host cells may also be fusion proteins
which include regions from heterologous proteins. Such regions may be included to
allow, *e.g.*, secretion, improved stability, or facilitated purification of the
polypeptide. For example, a sequence encoding an appropriate signal peptide can be
incorporated into expression vectors. A DNA sequence for a signal peptide
20 (secretory leader) may be fused in-frame to the polynucleotide sequence so that the
polypeptide is translated as a fusion protein comprising the signal peptide. A signal
peptide that is functional in the intended host cell promotes extracellular secretion of
the polypeptide. Preferably, the signal sequence will be cleaved from the
polypeptide upon secretion of the polypeptide from the cell. Thus, preferred fusion
25 proteins can be produced in which the N-terminus of a protease polypeptide is fused
to a carrier peptide.

In one embodiment, the polypeptide comprises a fusion protein which
includes a heterologous region used to facilitate purification of the polypeptide.
Many of the available peptides used for such a function allow selective binding of

the fusion protein to a binding partner. A preferred binding partner includes one or more of the IgG binding domains of protein A are easily purified to homogeneity by affinity chromatography on, for example, IgG-coupled Sepharose. Alternatively, many vectors have the advantage of carrying a stretch of histidine residues that can
5 be expressed at the N-terminal or C-terminal end of the target protein, and thus the protein of interest can be recovered by metal chelation chromatography. A nucleotide sequence encoding a recognition site for a proteolytic enzyme such as enterokinase, factor X procollagenase or thrombin may immediately precede the sequence for a protease polypeptide to permit cleavage of the fusion protein to
10 obtain the mature protease polypeptide. Additional examples of fusion-protein binding partners include, but are not limited to, the yeast I-factor, the honeybee melatin leader in sf9 insect cells, 6-His tag, thioredoxin tag, hemagglutinin tag, GST tag, and OmpA signal sequence tag. As will be understood by one of skill in the art, the binding partner which recognizes and binds to the peptide may be any ion,
15 molecule or compound including metal ions (e.g., metal affinity columns), antibodies, or fragments thereof, and any protein or peptide which binds the peptide, such as the FLAG tag.

Antibodies

20 In another aspect, the invention features an antibody (e.g., a monoclonal or polyclonal antibody) having specific binding affinity to a protease polypeptide or a protease polypeptide domain or fragment where the polypeptide is selected from the group having a sequence at least about 90% identical to an amino acid sequence selected from the group consisting of those set forth in SEQ ID NO:60, SEQ ID
25 NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID

NO:86, SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID
NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID
NO:96, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID
NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105,
5 SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:109, SEQ ID
NO:110, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114,
SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118.

Preferably the polypeptide is has at least about 90%, 91%, 92%, 93%, 94%, 95%,
96%, 97%, 98% 99% or 100% identity with the sequences listed above. By
10 "specific binding affinity" is meant that the antibody binds to the target protease
polypeptide with greater affinity than it binds to other polypeptides under specified
conditions. Antibodies or antibody fragments are polypeptides that contain regions
that can bind other polypeptides. The term "specific binding affinity" describes an
antibody that binds to a protease polypeptide with greater affinity than it binds to
15 other polypeptides under specified conditions. Antibodies can be used to identify an
endogenous source of protease polypeptides, to monitor cell cycle regulation, and
for immuno-localization of protease polypeptides within the cell.

The term "polyclonal" refers to antibodies that are heterogenous populations
of antibody molecules derived from the sera of animals immunized with an antigen
20 or an antigenic functional derivative thereof. For the production of polyclonal
antibodies, various host animals may be immunized by injection with the antigen.
Various adjuvants may be used to increase the immunological response, depending
on the host species.

"Monoclonal antibodies" are substantially homogenous populations of
25 antibodies to a particular antigen. They may be obtained by any technique which
provides for the production of antibody molecules by continuous cell lines in
culture. Monoclonal antibodies may be obtained by methods known to those skilled
in the art (Kohler *et al.*, *Nature*, 1975, 256:495-497, and U.S. Patent No. 4,376,110,

both of which are hereby incorporated by reference herein in their entirety including any figures, tables, or drawings).

An antibody of the present invention includes "humanized" monoclonal and polyclonal antibodies. Humanized antibodies are recombinant proteins in which non-human (typically murine) complementarity determining regions of an antibody have been transferred from heavy and light variable chains of the non-human (e.g. murine) immunoglobulin into a human variable domain, followed by the replacement of some human residues in the framework regions of their murine counterparts. Humanized antibodies in accordance with this invention are suitable for use in therapeutic methods. General techniques for cloning murine immunoglobulin variable domains are described, for example, by the publication of Orlandi *et al.*, *Proc. Nat'l Acad. Sci. USA* 86: 3833 (1989). Techniques for producing humanized monoclonal antibodies are described, for example, by Jones *et al.*, *Nature* 321:522 (1986), Riechmann *et al.*, *Nature* 332:323 (1988), Verhoeven *et al.*, *Science* 239:1534 (1988), Carter *et al.*, *Proc. Nat'l Acad. Sci. USA* 89:4285 (1992), Sandhu, *Crit. Rev. Biotech.* 12:437 (1992), and Singer *et al.*, *J. Immun.* 150:2844 (1993).

The term "antibody fragment" refers to a portion of an antibody, often the hypervariable region and portions of the surrounding heavy and light chains, that displays specific binding affinity for a particular molecule. A hypervariable region is a portion of an antibody that physically binds to the polypeptide target.

An antibody fragment of the present invention includes a "single-chain antibody," a phrase used in this description to denote a linear polypeptide that binds antigen with specificity and that comprises variable or hypervariable regions from the heavy and light chain chains of an antibody. Such single chain antibodies can be produced by conventional methodology. The Vh and Vl regions of the Fv fragment can be covalently joined and stabilized by the insertion of a disulfide bond. See Glockshuber, *et al.*, *Biochemistry* 1362 (1990). Alternatively, the Vh and Vl regions can be joined by the insertion of a peptide linker. A gene encoding the Vh, Vl and

peptide linker sequences can be constructed and expressed using a recombinant expression vector. See Colcher, *et al.*, *J. Nat'l Cancer Inst.* 82: 1191 (1990). Amino acid sequences comprising hypervariable regions from the Vh and VI antibody chains can also be constructed using disulfide bonds or peptide linkers.

5 Antibodies or antibody fragments having specific binding affinity to a protease polypeptide of the invention may be used in methods for detecting the presence and/or amount of protease polypeptide in a sample by probing the sample with the antibody under conditions suitable for protease-antibody immunocomplex formation and detecting the presence and/or amount of the antibody conjugated to
10 the protease polypeptide. Diagnostic kits for performing such methods may be constructed to include antibodies or antibody fragments specific for the protease as well as a conjugate of a binding partner of the antibodies or the antibodies themselves.

 An antibody or antibody fragment with specific binding affinity to a protease
15 polypeptide of the invention can be isolated, enriched, or purified from a prokaryotic or eukaryotic organism. Routine methods known to those skilled in the art enable production of antibodies or antibody fragments, in both prokaryotic and eukaryotic organisms. Purification, enrichment, and isolation of antibodies, which are polypeptide molecules, are described above.

20 Antibodies having specific binding affinity to a protease polypeptide of the invention may be used in methods for detecting the presence and/or amount of protease polypeptide in a sample by contacting the sample with the antibody under conditions such that an immunocomplex forms and detecting the presence and/or amount of the antibody conjugated to the protease polypeptide. Diagnostic kits for
25 performing such methods may be constructed to include a first container containing the antibody and a second container having a conjugate of a binding partner of the antibody and a label, such as, for example, a radioisotope. The diagnostic kit may also include notification of an FDA approved use and instructions therefor.

In another aspect, the invention features a hybridoma which produces an antibody having specific binding affinity to a protease polypeptide or a protease polypeptide domain, where the polypeptide is selected from the group consisting of those set forth in SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, 5 SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, 10 SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, 15 SEQ ID NO:117 and SEQ ID NO:118. By "hybridoma" is meant an immortalized cell line that is capable of secreting an antibody, for example an antibody to a protease of the invention. In preferred embodiments, the antibody to the protease comprises a sequence of amino acids that is able to specifically bind a protease polypeptide of the invention. 20

In another aspect, the present invention is also directed to kits comprising antibodies that bind to a polypeptide encoded by any of the nucleic acid molecules described above, and a negative control antibody.

The term "negative control antibody" refers to an antibody derived from 25 similar source as the antibody having specific binding affinity, but where it displays no binding affinity to a polypeptide of the invention.

In another aspect, the invention features a protease polypeptide binding agent able to bind to a protease polypeptide selected from the group having an amino acid sequence selected from the group consisting of those set forth in SEQ ID NO:60,

SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65,
SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70,
SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75,
SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80,
5 SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85,
SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90,
SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95,
SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100,
SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID
10 NO:105, SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:109,
SEQ ID NO:110, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID
NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118.

The binding agent is preferably a purified antibody that recognizes an epitope
present on a protease polypeptide of the invention. Other binding agents include
15 molecules that bind to protease polypeptides and analogous molecules that bind to a
protease polypeptide. Such binding agents may be identified by using assays that
measure protease binding partner activity, or they may be identified using assays
that measure protease activity, such as the release of a fluorogenic or radioactive
marker attached to a substrate molecule.

20

Screening Methods to Detect Protease Polypeptides

The invention also features a method for screening for human cells
containing a protease polypeptide of the invention or an equivalent sequence. The
method involves identifying the novel polypeptide in human cells using techniques
25 that are routine and standard in the art, such as those described herein for identifying
the proteases of the invention (*e.g.*, cloning, Southern or Northern blot analysis, *in situ*
hybridization, PCR amplification, etc.).

Screening Methods to Identify Substances that Modulate Protease

Activity

In another aspect, the invention features methods for identifying a substance that modulates protease activity comprising the steps of: (a) contacting a protease polypeptide comprising an amino acid sequence substantially identical to a sequence selected from the group consisting of those set forth in SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118 with a test substance; (b) measuring the activity of said polypeptide; and (c) determining whether said substance modulates the activity of said polypeptide. More preferably the sequence is at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the listed sequences.

The term "modulates" refers to the ability of a compound to alter the function of a protease of the invention. A modulator preferably activates or inhibits the activity of a protease of the invention depending on the concentration of the compound exposed to the protease.

The term "modulates" also refers to altering the function of proteases of the invention by increasing or decreasing the probability that a complex forms between the protease and a natural binding partner. A modulator preferably increases the

probability that such a complex forms between the protease and the natural binding partner, more preferably increases or decreases the probability that a complex forms between the protease and the natural binding partner depending on the concentration of the compound exposed to the protease, and most preferably decreases the probability that a complex forms between the protease and the natural binding partner.

The term "activates" refers to increasing the cellular activity of the protease. The term "inhibits" refers to decreasing the cellular activity of the protease.

The term "complex" refers to an assembly of at least two molecules bound to one another. Signal transduction complexes often contain at least two protein molecules bound to one another. For instance, a protein tyrosine receptor protein kinase, GRB2, SOS, RAF, and RAS assemble to form a signal transduction complex in response to a mitogenic ligand. Similarly, the proteases involved in blood coagulation and their cofactors are known to form macromolecular complexes on cellular membranes. Additionally, proteases involved in modification of the extracellular matrix are known to form complexes with their inhibitors and also with components of the extracellular matrix.

The term "natural binding partner" refers to polypeptides, lipids, small molecules, or nucleic acids that bind to proteases in cells. A change in the interaction between a protease and a natural binding partner can manifest itself as an increased or decreased probability that the interaction forms, or an increased or decreased concentration of protease/natural binding partner complex.

The term "contacting" as used herein refers to mixing a solution comprising the test compound with a liquid medium bathing the cells of the methods. The solution comprising the compound may also comprise another component, such as dimethyl sulfoxide (DMSO), which facilitates the uptake of the test compound or compounds into the cells of the methods. The solution comprising the test compound may be added to the medium bathing the cells by utilizing a delivery apparatus, such as a pipette-based device or syringe-based device.

In another aspect, the invention features methods for identifying a substance that modulates protease activity in a cell comprising the steps of: (a) expressing a protease polypeptide in a cell, wherein said polypeptide has a sequence substantially identical to an amino acid sequence selected from the group consisting of those set forth in SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118; (b) adding a test substance to said cell; and (c) monitoring a change in cell phenotype, cell proliferation, cell differentiation or the interaction between said polypeptide and a natural binding partner.

The term "expressing" as used herein refers to the production of proteases of the invention from a nucleic acid vector containing protease genes within a cell. The nucleic acid vector is transfected into cells using well known techniques in the art as described herein.

Another aspect of the instant invention is directed to methods of identifying compounds that bind to protease polypeptides of the present invention, comprising contacting the protease polypeptides with a compound, and determining whether the compound binds the protease polypeptides. Binding can be determined by binding assays which are well known to the skilled artisan, including, but not limited to, gel-shift assays, Western blots, radiolabeled competition assay, phage-based expression

cloning, co-fractionation by chromatography, co-precipitation, cross linking, interaction trap/two-hybrid analysis, southwestern analysis, ELISA, and the like, which are described in, for example, *Current Protocols in Molecular Biology*, 1999, John Wiley & Sons, NY, which is incorporated herein by reference in its entirety.

- 5 The compounds to be screened include, but are not limited to, compounds of extracellular, intracellular, biological or chemical origin.

The methods of the invention also embrace compounds that are attached to a label, such as a radiolabel (e.g., ^{125}I , ^{35}S , ^{32}P , ^{33}P , ^3H), a fluorescence label, a chemiluminescent label, an enzymic label and an immunogenic label. The protease polypeptides employed in such a test may either be free in solution, attached to a solid support, borne on a cell surface, located intracellularly or associated with a portion of a cell. One skilled in the art can, for example, measure the formation of complexes between a protease polypeptide and the compound being tested. Alternatively, one skilled in the art can examine the diminution in complex formation between a protease polypeptide and its substrate caused by the compound being tested.

Other assays can be used to examine enzymatic activity including, but not limited to, photometric, radiometric, HPLC, electrochemical, and the like, which are described in, for example, Enzyme Assays: A Practical Approach, eds. R. Eisenthal and M. J. Danson, 1992, Oxford University Press, which is incorporated herein by reference in its entirety.

Another aspect of the present invention is directed to methods of identifying compounds which modulate (i.e., increase or decrease) activity of a protease polypeptide comprising contacting the protease polypeptide with a compound, and determining whether the compound modifies activity of the protease polypeptide. These compounds are also referred to as "modulators of proteases." The activity in the presence of the test compound is measured to the activity in the absence of the test compound. Where the activity of a sample containing the test compound is higher than the activity in a sample lacking the test compound, the compound will

have increased the activity. Similarly, where the activity of a sample containing the test compound is lower than the activity in the sample lacking the test compound, the compound will have inhibited the activity.

The present invention is particularly useful for screening compounds by using a protease polypeptide in any of a variety of drug screening techniques. The compounds to be screened include, but are not limited to, extracellular, intracellular, biological or chemical origin. The protease polypeptide employed in such a test may be in any form, preferably, free in solution, attached to a solid support, borne on a cell surface or located intracellularly. One skilled in the art can measure the change in rate that a protease of the invention cleaves a substrate polypeptide. One skilled in the art can also, for example, measure the formation of complexes between a protease polypeptide and the compound being tested. Alternatively, one skilled in the art can examine the diminution in complex formation between a protease polypeptide and its substrate caused by the compound being tested.

The activity of protease polypeptides of the invention can be determined by, for example, examining the ability to bind or be activated by chemically synthesised peptide ligands. Alternatively, the activity of the protease polypeptides can be assayed by examining their ability to bind metal ions such as calcium, hormones, chemokines, neuropeptides, neurotransmitters, nucleotides, lipids, odorants, and photons. Thus, modulators of the protease polypeptide's activity may alter a protease function, such as a binding property of a protease or an activity such as cleaving protein substrates or polypeptide substrates, or membrane localization.

In various embodiments of the method, the assay may take the form of a yeast growth assay, an Aequorin assay, a Luciferase assay, a mitogenesis assay, a MAP Kinase activity assay, as well as other binding or function-based assays of protease activity that are generally known in the art. In several of these embodiments, the invention includes any of the serine proteases, cysteine proteases, aspartyl proteases, metalloproteases, threonine proteases, and other proteases. Biological activities of proteases according to the invention include, but are not

limited to, the binding of a natural or a synthetic ligand, as well as any one of the functional activities of proteases known in the art. Non-limiting examples of protease activities include cleavage of polypeptide chains, processing the pro-form of a polypeptide chain to the active product, transmembrane signaling of various forms, and/or the modification of the extracellular matrix.

The modulators of the invention exhibit a variety of chemical structures, which can be generally grouped into mimetics of natural protease ligands, and peptide and non-peptide allosteric effectors of proteases. The invention does not restrict the sources for suitable modulators, which may be obtained from natural sources such as plant, animal or mineral extracts, or non-natural sources such as small molecule libraries, including the products of combinatorial chemical approaches to library construction, and peptide libraries.

The use of cDNAs encoding proteins in drug discovery programs is well-known; assays capable of testing thousands of unknown compounds per day in high-throughput screens (HTSs) are thoroughly documented. The literature is replete with examples of the use of radiolabelled ligands in HTS binding assays for drug discovery (see, Williams, *Medicinal Research Reviews*, 1991, 11:147-184.; Sweetnam, *et al.*, *J. Natural Products*, 1993, 56:441-455 for review). Recombinant proteins are preferred for binding assay HTS because they allow for better specificity (higher relative purity), provide the ability to generate large amounts of receptor material, and can be used in a broad variety of formats (see Hodgson, *Bio/Technology*, 1992, 10:973-980 which is incorporated herein by reference in its entirety). A variety of heterologous systems is available for functional expression of recombinant proteins that are well known to those skilled in the art. Such systems include bacteria (Strosberg, *et al.*, *Trends in Pharmacological Sciences*, 1992, 13:95-98), yeast (Pausch, *Trends in Biotechnology*, 1997, 15:487-494), several kinds of insect cells (Vanden Broeck, *Int. Rev. Cytology*, 1996, 164:189-268), amphibian cells (Jayawickreme *et al.*, *Current Opinion in Biotechnology*, 1997, 8:629-634) and several mammalian cell lines (CHO, HEK293, COS, etc.; see, Gerhardt, *et al.*, *Eur.*

J. Pharmacology, 1997, 334:1-23). These examples do not preclude the use of other possible cell expression systems, including cell lines obtained from nematodes (PCT application WO 98/37177).

5 An expressed protease can be used for HTS binding assays in conjunction with its defined ligand, in this case the corresponding peptide that activates it. The identified peptide is labeled with a suitable radioisotope, including, but not limited to, ^{125}I , ^3H , ^{35}S or ^{32}P , by methods that are well known to those skilled in the art. Alternatively, the peptides may be labeled by well-known methods with a suitable fluorescent derivative (Baindur, *et al.*, *Drug Dev. Res.*, 1994, 33:373-398; Rogers, 10 *Drug Discovery Today*, 1997, 2:156-160). Radioactive ligand specifically bound to the receptor in membrane preparations made from the cell line expressing the recombinant protein can be detected in HTS assays in one of several standard ways, including filtration of the receptor-ligand complex to separate bound ligand from unbound ligand (Williams, *Med. Res. Rev.*, 1991, 11:147-184.; Sweetnam, *et al.*, *J.* 15 *Natural Products*, 1993, 56:441-455). Alternative methods include a scintillation proximity assay (SPA) or a FlashPlate format in which such separation is unnecessary (Nakayama, *Cur. Opinion Drug Disc. Dev.*, 1998, 1:85-91 Bossé, *et al.*, *J. Biomolecular Screening*, 1998, 3:285-292.). Binding of fluorescent ligands can be detected in various ways, including fluorescence energy transfer (FRET), direct 20 spectrophotofluorometric analysis of bound ligand, or fluorescence polarization (Rogers, *Drug Discovery Today*, 1997, 2:156-160; Hill, *Cur. Opinion Drug Disc. Dev.*, 1998, 1:92-97).

The proteases and natural binding partners required for functional expression of heterologous protease polypeptides can be native constituents of the host cell or 25 can be introduced through well-known recombinant technology. The protease polypeptides can be intact or chimeric. The protease activation may result in the stimulation or inhibition of other native proteins, events that can be linked to a measurable response.

Examples of such biological responses include, but are not limited to, the following: the ability to survive in the absence of a limiting nutrient in specifically engineered yeast cells (Pausch, *Trends in Biotechnology*, 1997, 15:487-494); changes in intracellular Ca^{2+} concentration as measured by fluorescent dyes (Murphy, *et al.*, *Cur. Opinion Drug Disc. Dev.*, 1998, 1:192-199). Fluorescence changes can also be used to monitor ligand-induced changes in membrane potential or intracellular pH; an automated system suitable for HTS has been described for these purposes (Schroeder, *et al.*, *J. Biomolecular Screening*, 1996, 1:75-80). Assays are also available for the measurement of common second but these are not generally preferred for HTS.

The invention contemplates a multitude of assays to screen and identify inhibitors of ligand binding to protease polypeptides or of substrate cleavage by protease polypeptides. In one example, the protease polypeptide is immobilized and interaction with a binding partner or substrate is assessed in the presence and absence of a candidate modulator such as an inhibitor compound. In another example, interaction between the protease polypeptide and its binding partner or a substrate is assessed in a solution assay, both in the presence and absence of a candidate inhibitor compound. In either assay, an inhibitor is identified as a compound that decreases binding between the protease polypeptide and its natural binding partner or the activity of a protease polypeptide in cleaving a substrate molecule. Another contemplated assay involves a variation of the di-hybrid assay wherein an inhibitor of protein/protein interactions is identified by detection of a positive signal in a transformed or transfected host cell, as described in PCT publication number WO 95/20652, published August 3, 1995 and is included by reference herein including any figures, tables, or drawings.

Candidate modulators contemplated by the invention include compounds selected from libraries of either potential activators or potential inhibitors. There are a number of different libraries used for the identification of small molecule modulators, including: (1) chemical libraries, (2) natural product libraries, and (3)

combinatorial libraries comprised of random peptides, oligonucleotides or organic molecules. Chemical libraries consist of random chemical structures, some of which are analogs of known compounds or analogs of compounds that have been identified as "hits" or "leads" in other drug discovery screens, while others are derived from

5 natural products, and still others arise from non-directed synthetic organic chemistry. Natural product libraries are collections of microorganisms, animals, plants, or marine organisms which are used to create mixtures for screening by: (1) fermentation and extraction of broths from soil, plant or marine microorganisms or (2) extraction of plants or marine organisms. Natural product libraries include

10 polyketides, non-ribosomal peptides, and variants (non-naturally occurring) thereof. For a review, *see, Science* 282:63-68 (1998). Combinatorial libraries are composed of large numbers of peptides, oligonucleotides, or organic compounds as a mixture. These libraries are relatively easy to prepare by traditional automated synthesis methods, PCR, cloning, or proprietary synthetic methods. Of particular interest are

15 non-peptide combinatorial libraries. Still other libraries of interest include peptide, protein, peptidomimetic, multiparallel synthetic collection, recombinatorial, and polypeptide libraries. For a review of combinatorial chemistry and libraries created therefrom, *see, Myers, Curr. Opin. Biotechnol.* 8:701-707 (1997). Identification of modulators through use of the various libraries described herein permits

20 modification of the candidate "hit" (or "lead") to optimize the capacity of the "hit" to modulate activity.

Still other candidate inhibitors contemplated by the invention can be designed and include soluble forms of binding partners, as well as such binding partners as chimeric, or fusion, proteins. A "binding partner" as used herein broadly

25 encompasses both natural binding partners as described above as well as chimeric polypeptides, peptide modulators other than natural ligands, antibodies, antibody fragments, and modified compounds comprising antibody domains that are immunospecific for the expression product of the identified protease gene.

Other assays may be used to identify specific peptide ligands of a protease polypeptide, including assays that identify ligands of the target protein through measuring direct binding of test ligands to the target protein, as well as assays that identify ligands of target proteins through affinity ultrafiltration with ion spray mass spectroscopy/HPLC methods or other physical and analytical methods.

Alternatively, such binding interactions are evaluated indirectly using the yeast two-hybrid system described in Fields *et al.*, *Nature*, 340:245-246 (1989), and Fields *et al.*, *Trends in Genetics*, 10:286-292 (1994), both of which are incorporated herein by reference. The two-hybrid system is a genetic assay for detecting interactions between two proteins or polypeptides. It can be used to identify proteins that bind to a known protein of interest, or to delineate domains or residues critical for an interaction. Variations on this methodology have been developed to clone genes that encode DNA binding proteins, to identify peptides that bind to a protein, and to screen for drugs. The two-hybrid system exploits the ability of a pair of interacting proteins to bring a transcription activation domain into close proximity with a DNA binding domain that binds to an upstream activation sequence (UAS) of a reporter gene, and is generally performed in yeast. The assay requires the construction of two hybrid genes encoding (1) a DNA-binding domain that is fused to a first protein and (2) an activation domain fused to a second protein. The DNA-binding domain targets the first hybrid protein to the UAS of the reporter gene; however, because most proteins lack an activation domain, this DNA-binding hybrid protein does not activate transcription of the reporter gene. The second hybrid protein, which contains the activation domain, cannot by itself activate expression of the reporter gene because it does not bind the UAS. However, when both hybrid proteins are present, the noncovalent interaction of the first and second proteins tethers the activation domain to the UAS, activating transcription of the reporter gene. For example, when the first protein is a protease gene product, or fragment thereof, that is known to interact with another protein or nucleic acid, this assay can be used to detect agents that interfere with the binding interaction. Expression of the reporter

gene is monitored as different test agents are added to the system. The presence of an inhibitory agent results in lack of a reporter signal.

When the function of the protease polypeptide gene product is unknown and no ligands are known to bind the gene product, the yeast two-hybrid assay can also
5 be used to identify proteins that bind to the gene product. In an assay to identify proteins that bind to a protease polypeptide, or fragment thereof, a fusion polynucleotide encoding both a protease polypeptide (or fragment) and a UAS binding domain (*i.e.*, a first protein) may be used. In addition, a large number of hybrid genes each encoding a different second protein fused to an activation domain
10 are produced and screened in the assay. Typically, the second protein is encoded by one or more members of a total cDNA or genomic DNA fusion library, with each second protein coding region being fused to the activation domain. This system is applicable to a wide variety of proteins, and it is not even necessary to know the identity or function of the second binding protein. The system is highly sensitive
15 and can detect interactions not revealed by other methods; even transient interactions may trigger transcription to produce a stable mRNA that can be repeatedly translated to yield the reporter protein.

Other assays may be used to search for agents that bind to the target protein. One such screening method to identify direct binding of test ligands to a target
20 protein is described in U.S. Patent No. 5,585,277, incorporated herein by reference. This method relies on the principle that proteins generally exist as a mixture of folded and unfolded states, and continually alternate between the two states. When a test ligand binds to the folded form of a target protein (*i.e.*, when the test ligand is a ligand of the target protein), the target protein molecule bound by the ligand remains
25 in its folded state. Thus, the folded target protein is present to a greater extent in the presence of a test ligand which binds the target protein, than in the absence of a ligand. Binding of the ligand to the target protein can be determined by any method which distinguishes between the folded and unfolded states of the target protein. The function of the target protein need not be known in order for this assay to be

performed. Virtually any agent can be assessed by this method as a test ligand, including, but not limited to, metals, polypeptides, proteins, lipids, polysaccharides, polynucleotides and small organic molecules.

Another method for identifying ligands of a target protein is described in
5 Wieboldt *et al.*, *Anal. Chem.*, 69:1683-1691 (1997), incorporated herein by
reference. This technique screens combinatorial libraries of 20-30 agents at a time
in solution-phase for binding to the target protein. Agents that bind to the target
protein are separated from other library components by simple membrane washing.
The specifically selected molecules that are retained on the filter are subsequently
10 liberated from the target protein and analyzed by HPLC and pneumatically assisted
electrospray (ion spray) ionization mass spectroscopy. This procedure selects
library components with the greatest affinity for the target protein, and is particularly
useful for small molecule libraries.

In preferred embodiments of the invention, methods of screening for
15 compounds which modulate protease activity comprise contacting test compounds
with protease polypeptides and assaying for the presence of a complex between the
compound and the protease polypeptide. In such assays, the ligand is typically
labelled. After suitable incubation, free ligand is separated from that present in
bound form, and the amount of free or uncomplexed label is a measure of the ability
20 of the particular compound to bind to the protease polypeptide.

In another embodiment of the invention, high throughput screening for
compounds having suitable binding affinity to protease polypeptides is employed.
Briefly, large numbers of different small peptide test compounds are synthesised on
a solid substrate. The peptide test compounds are contacted with the protease
25 polypeptide and washed. Bound protease polypeptide is then detected by methods
well known in the art. Purified polypeptides of the invention can also be coated
directly onto plates for use in the aforementioned drug screening techniques. In
addition, non-neutralizing antibodies can be used to capture the protein and
immobilize it on the solid support.

Other embodiments of the invention comprise using competitive screening assays in which neutralizing antibodies capable of binding a polypeptide of the invention specifically compete with a test compound for binding to the polypeptide. In this manner, the antibodies can be used to detect the presence of any peptide that shares one or more antigenic determinants with a protease polypeptide.

Radiolabeled competitive binding studies are described in A.H. Lin *et al.*

Antimicrobial Agents and Chemotherapy, 1997, vol. 41, no. 10. pp. 2127-2131, the disclosure of which is incorporated herein by reference in its entirety.

Therapeutic Methods

The invention includes methods for treating a disease or disorder by administering to a patient in need of such treatment a protease polypeptide substantially identical to an amino acid sequence selected from the group consisting of those set forth in SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118, and any other protease polypeptide of the present invention. As discussed in the section "Gene Therapy," a protease polypeptide of the invention may also be administered indirectly by via administration of suitable polynucleotide means for *in vivo* expression of the protease polypeptide. Preferably the protease polypeptide will have at least about

90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or 100% identity to one of the aforementioned sequences.

In another aspect, the invention provides methods for treating a disease or disorder by administering to a patient in need of such treatment a substance that modulates the activity of a protease substantially identical to a sequence selected from the group consisting of those set forth in SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118.

Preferably the disease is selected from the group consisting of cancers, immune-related diseases and disorders, cardiovascular disease, brain or neuronal-associated diseases, and metabolic disorders. More specifically these diseases include cancer of tissues, blood, or hematopoietic origin, particularly those involving breast, colon, lung, prostate, cervical, brain, ovarian, bladder, or kidney; central or peripheral nervous system diseases and conditions including migraine, pain, sexual dysfunction, mood disorders, attention disorders, cognition disorders, hypotension, and hypertension; psychotic and neurological disorders, including anxiety, schizophrenia, manic depression, delirium, dementia, severe mental retardation and dyskinesias, such as Huntington's disease or Tourette's Syndrome; neurodegenerative diseases including Alzheimer's, Parkinson's, Multiple sclerosis,

- and Amyotrophic lateral sclerosis; viral or non-viral infections caused by HIV-1, HIV-2 or other viral- or prion-agents or fungal- or bacterial- organisms; metabolic disorders including Diabetes and obesity and their related syndromes, among others; cardiovascular disorders including reperfusion restenosis, coronary thrombosis,
- 5 clotting disorders, unregulated cell growth disorders, atherosclerosis; ocular disease including glaucoma, retinopathy, and macular degeneration; inflammatory disorders including rheumatoid arthritis, chronic inflammatory bowel disease, chronic inflammatory pelvic disease, multiple sclerosis, asthma, osteoarthritis, psoriasis, atherosclerosis, rhinitis, autoimmunity, and organ transplant rejection.
- 10 In preferred embodiments, the invention provides methods for treating or preventing a disease or disorder by administering to a patient in need of such treatment a substance that modulates the activity of a protease polypeptide having an amino acid sequence selected from the group consisting of those set forth in SEQ ID
- 15 NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID
- 20 NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113,
- 25 SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118.

Preferably the disease is selected from the group consisting of cancers, immune-related diseases and disorders, cardiovascular disease, brain or neuronal-associated diseases, and metabolic disorders. More specifically these diseases

include cancer of tissues, blood, or hematopoietic origin, particularly those involving breast, colon, lung, prostate, cervical, brain, ovarian, bladder, or kidney; central or peripheral nervous system diseases and conditions including migraine, pain, sexual dysfunction, mood disorders, attention disorders, cognition disorders, hypotension, and hypertension; psychotic and neurological disorders, including anxiety, schizophrenia, manic depression, delirium, dementia, severe mental retardation and dyskinesias, such as Huntington's disease or Tourette's Syndrome; neurodegenerative diseases including Alzheimer's, Parkinson's, Multiple sclerosis, and Amyotrophic lateral sclerosis; viral or non-viral infections caused by HIV-1, HIV-2 or other viral- or prion-agents or fungal- or bacterial- organisms; metabolic disorders including Diabetes and obesity and their related syndromes, among others; cardiovascular disorders including reperfusion restenosis, coronary thrombosis, clotting disorders, unregulated cell growth disorders, atherosclerosis; ocular disease including glaucoma, retinopathy, and macular degeneration; inflammatory disorders including rheumatoid arthritis, chronic inflammatory bowel disease, chronic inflammatory pelvic disease, multiple sclerosis, asthma, osteoarthritis, psoriasis, atherosclerosis, rhinitis, autoimmunity, and organ transplant rejection.

Preferably the disease is selected from the group consisting of immune-related diseases and disorders, cardiovascular disease, and cancer. Most preferably, the immune-related diseases and disorders are selected from the group consisting of rheumatoid arthritis, chronic inflammatory bowel disease, chronic inflammatory pelvic disease, multiple sclerosis, asthma, osteoarthritis, psoriasis, atherosclerosis, rhinitis, autoimmunity, and organ transplantation.

Substances useful for treatment of protease-related disorders or diseases preferably show positive results in one or more *in vitro* assays for an activity corresponding to treatment of the disease or disorder in question (Examples of such assays are provided herein, including Example 7). Examples of substances that can be screened for favorable activity are provided and referenced throughout the specification, including this section (Screening Methods to Identify Substances

that Modulate Protease Activity). The substances that modulate the activity of the proteases preferably include, but are not limited to, antisense oligonucleotides, ribozymes, and other inhibitors of proteases, as determined by methods and screens referenced this section and in Example 7, below, and any other suitable methods.

- 5 The use of antisense oligonucleotides and ribozymes are discussed more fully in the Section "Gene Therapy," below.

The term "preventing" refers to decreasing the probability that an organism contracts or develops an abnormal condition.

- 10 The term "treating" refers to having a therapeutic effect and at least partially alleviating or abrogating an abnormal condition in the organism.

- The term "therapeutic effect" refers to the inhibition or activation factors causing or contributing to the abnormal condition. A therapeutic effect relieves to some extent one or more of the symptoms of the abnormal condition. In reference to the treatment of abnormal conditions, a therapeutic effect can refer to one or more of the following: (a) an increase or decrease in the proliferation, growth, and/or
- 15 differentiation of cells; (b) activation or inhibition (*i.e.*, slowing or stopping) of cell death; (c) inhibition of degeneration; (d) relieving to some extent one or more of the symptoms associated with the abnormal condition; and (e) enhancing the function of the affected population of cells. Compounds demonstrating efficacy against
- 20 abnormal conditions can be identified as described herein.

The term "abnormal condition" refers to a function in the cells or tissues of an organism that deviates from their normal functions in that organism. An abnormal condition can relate to cell proliferation, cell differentiation, or cell survival.

- 25 Abnormal cell proliferative conditions include cancers such as fibrotic and mesangial disorders, abnormal angiogenesis and vasculogenesis, wound healing, psoriasis, diabetes mellitus, and inflammation.

Abnormal differentiation conditions include, but are not limited to neurodegenerative disorders, slow wound healing rates, and slow tissue grafting healing rates.

Abnormal cell survival conditions relate to conditions in which programmed cell death (apoptosis) pathways are activated or abrogated. A number of proteases are associated with the apoptosis pathways. Aberrations in the function of any one of the proteases could lead to cell immortality or premature cell death.

The term "aberration", in conjunction with the function of a protease in a signal transduction process, refers to a protease that is over- or under-expressed in an organism, mutated such that its catalytic activity is lower or higher than wild-type protease activity, mutated such that it can no longer interact with a natural binding partner, is no longer modified by another protein, or no longer interacts with a natural binding partner.

The term "administering" relates to a method of incorporating a compound into cells or tissues of an organism. The abnormal condition can be prevented or treated when the cells or tissues of the organism exist within the organism or outside of the organism. Cells existing outside the organism can be maintained or grown in cell culture dishes. For cells harbored within the organism, many techniques exist in the art to administer compounds, including (but not limited to) oral, parenteral, dermal, injection, and aerosol applications. For cells outside of the organism, multiple techniques exist in the art to administer the compounds, including (but not limited to) cell microinjection techniques, transformation techniques, and carrier techniques.

The abnormal condition can also be prevented or treated by administering a compound to a group of cells having an aberration in a signal transduction pathway to an organism. The effect of administering a compound on organism function can then be monitored. The organism is preferably a mouse, rat, rabbit, guinea pig, or goat, more preferably a monkey or ape, and most preferably a human.

In another aspect, the invention features methods for detection of a protease polypeptide in a sample as a diagnostic tool for diseases or disorders, wherein the method comprises the steps of: (a) contacting the sample with a nucleic acid probe which hybridizes under hybridization assay conditions to a nucleic acid target region of a protease polypeptide having an amino acid sequence selected from the group consisting of those set forth in SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118, said probe comprising the nucleic acid sequence encoding the polypeptide, fragments thereof, and the complements of the sequences and fragments; and (b) detecting the presence or amount of the probe:target region hybrid as an indication of the disease.

In preferred embodiments of the invention, the disease or disorder is selected from the group consisting of rheumatoid arthritis, arteriosclerosis, autoimmune disorders, organ transplantation, myocardial infarction, cardiomyopathies, stroke, renal failure, oxidative stress-related neurodegenerative disorders, and cancer. Preferably the disease is selected from the group consisting of cancers, immune-related diseases and disorders, cardiovascular disease, brain or neuronal-associated diseases, and metabolic disorders. More specifically these diseases include cancer of tissues, blood, or hematopoietic origin, particularly those involving breast, colon,

lung, prostate, cervical, brain, ovarian, bladder, or kidney; central or peripheral nervous system diseases and conditions including migraine, pain, sexual dysfunction, mood disorders, attention disorders, cognition disorders, hypotension, and hypertension; psychotic and neurological disorders, including anxiety, schizophrenia, manic depression, delirium, dementia, severe mental retardation and dyskinesias, such as Huntington's disease or Tourette's Syndrome; neurodegenerative diseases including Alzheimer's, Parkinson's, Multiple sclerosis, and Amyotrophic lateral sclerosis; viral or non-viral infections caused by HIV-1, HIV-2 or other viral- or prion-agents or fungal- or bacterial- organisms; metabolic disorders including Diabetes and obesity and their related syndromes, among others; cardiovascular disorders including reperfusion restenosis, coronary thrombosis, clotting disorders, unregulated cell growth disorders, atherosclerosis; ocular disease including glaucoma, retinopathy, and macular degeneration; inflammatory disorders including rheumatoid arthritis, chronic inflammatory bowel disease, chronic inflammatory pelvic disease, multiple sclerosis, asthma, osteoarthritis, psoriasis, atherosclerosis, rhinitis, autoimmunity, and organ transplant rejection.

The protease "target region" is the nucleotide base sequence selected from the group consisting of those set forth in SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:44, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:58,

and SEQ ID NO:59, or the corresponding full-length sequences, a functional derivative thereof, or a fragment thereof or a domain thereof to which the nucleic acid probe will specifically hybridize. Specific hybridization indicates that in the presence of other nucleic acids the probe only hybridizes detectably with the nucleic acid target region of the protease of the invention. Putative target regions can be identified by methods well known in the art consisting of alignment and comparison of the most closely related sequences in the database.

In preferred embodiments the nucleic acid probe hybridizes to a protease target region encoding at least 6, 12, 75, 90, 105, 120, 150, 200, 250, 300 or 350 contiguous amino acids of a sequence selected from the group consisting of those set forth in SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118, or the corresponding full-length amino acid sequence, or a functional derivative thereof. Hybridization conditions should be such that hybridization occurs only with the protease genes in the presence of other nucleic acid molecules. Under stringent hybridization conditions only highly complementary nucleic acid sequences hybridize. Preferably, such conditions prevent hybridization of nucleic acids having more than 1 or 2 mismatches out of 20 contiguous nucleotides. Such conditions are defined in Berger *et al.* (1987) (Guide

to Molecular Cloning Techniques pg 421, hereby incorporated by reference herein in its entirety including any figures, tables, or drawings.).

The diseases for which detection of protease genes in a sample could be diagnostic include diseases in which protease nucleic acid (DNA and/or RNA) is amplified in comparison to normal cells. By "amplification" is meant increased numbers of protease DNA or RNA in a cell compared with normal cells. In normal cells, proteases may be found as single copy genes. In selected diseases, the chromosomal location of the protease genes may be amplified, resulting in multiple copies of the gene, or amplification. Gene amplification can lead to amplification of protease RNA, or protease RNA can be amplified in the absence of protease DNA amplification.

"Amplification" as it refers to RNA can be the detectable presence of protease RNA in cells, since in some normal cells there is no basal expression of protease RNA. In other normal cells, a basal level of expression of protease exists, therefore in these cases amplification is the detection of at least 1-2-fold, and preferably more, protease RNA, compared to the basal level.

The diseases that could be diagnosed by detection of protease nucleic acid in a sample preferably include cancers. The test samples suitable for nucleic acid probing methods of the present invention include, for example, cells or nucleic acid extracts of cells, or biological fluids. The samples used in the above-described methods will vary based on the assay format, the detection method and the nature of the tissues, cells or extracts to be assayed. Methods for preparing nucleic acid extracts of cells are well known in the art and can be readily adapted in order to obtain a sample that is compatible with the method utilized.

In a final aspect, the invention features a method for detection of a protease polypeptide in a sample as a diagnostic tool for a disease or disorder, wherein the method comprises: (a) comparing a nucleic acid target region encoding the protease polypeptide in a sample, where the protease polypeptide has an amino acid sequence selected from the group consisting those set forth in SEQ ID NO:60, SEQ ID

- NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118, or one or more fragments thereof, with a control nucleic acid target region encoding the protease polypeptide, or one or more fragments thereof; and (b) detecting differences in sequence or amount between the target region and the control target region, as an indication of the disease or disorder. Preferably the disease is selected from the group consisting of cancers, immune-related diseases and disorders, cardiovascular disease, brain or neuronal-associated diseases, and metabolic disorders.
- More specifically these diseases include cancer of tissues, blood, or hematopoietic origin, particularly those involving breast, colon, lung, prostate, cervical, brain, ovarian, bladder, or kidney; central or peripheral nervous system diseases and conditions including migraine, pain, sexual dysfunction, mood disorders, attention disorders, cognition disorders, hypotension, and hypertension; psychotic and neurological disorders, including anxiety, schizophrenia, manic depression, delirium, dementia, severe mental retardation and dyskinesias, such as Huntington's disease or Tourette's Syndrome; neurodegenerative diseases including Alzheimer's, Parkinson's, Multiple sclerosis, and Amyotrophic lateral sclerosis; viral or non-viral infections caused by HIV-1, HIV-2 or other viral- or prion-agents

or fungal- or bacterial- organisms; metabolic disorders including Diabetes and obesity and their related syndromes, among others; cardiovascular disorders including reperfusion restenosis, coronary thrombosis, clotting disorders, unregulated cell growth disorders, atherosclerosis; ocular disease including
5 glaucoma, retinopathy, and macular degeneration; inflammatory disorders including rheumatoid arthritis, chronic inflammatory bowel disease, chronic inflammatory pelvic disease, multiple sclerosis, asthma, osteoarthritis, psoriasis, atherosclerosis, rhinitis, autoimmunity, and organ transplant rejection.

The term "comparing" as used herein refers to identifying discrepancies
10 between the nucleic acid target region isolated from a sample, and the control nucleic acid target region. The discrepancies can be in the nucleotide sequences, e.g. insertions, deletions, or point mutations, or in the amount of a given nucleotide sequence. Methods to determine these discrepancies in sequences are well-known to one of ordinary skill in the art. The "control" nucleic acid target region refers to the
15 sequence or amount of the sequence found in normal cells, e.g. cells that are not diseased as discussed previously.

The term "domain" refers to a region of a polypeptide which serves a particular function. For instance, N-terminal or C-terminal domains of signal transduction proteins can serve functions including, but not limited to, binding
20 molecules that localize the signal transduction molecule to different regions of the cell or binding other signaling molecules directly responsible for propagating a particular cellular signal. Some domains can be expressed separately from the rest of the protein and function by themselves, while others must remain part of the intact protein to retain function. The latter are termed functional regions of proteins
25 and also relate to domains.

The expression of proteases can be modulated by signal transduction pathways such as the Ras/MAP kinase signaling pathways. Additionally, the activity of proteases can modulate the activity of the MAP kinase signal transduction

pathway. Furthermore, proteases can be shown to be instrumental in the communication between disparate signal transduction pathways.

The term "signal transduction pathway" refers to the molecules that propagate an extracellular signal through the cell membrane to become an intracellular signal. This signal can then stimulate a cellular response. The polypeptide molecules involved in signal transduction processes are typically receptor and non-receptor protein tyrosine kinases, receptor and non-receptor protein phosphatases, polypeptides containing SRC homology 2 and 3 domains, phosphotyrosine binding proteins (SRC homology 2 (SH2) and phosphotyrosine binding (PTB and PH) domain containing proteins), proline-rich binding proteins (SH3 domain containing proteins), GTPases, phosphodiesterases, phospholipases, prolyl isomerases, proteases, Ca^{2+} binding proteins, cAMP binding proteins, guanyl cyclases, adenylyl cyclases, NO generating proteins, nucleotide exchange factors, and transcription factors.

The summary of the invention described above is not limiting and other features and advantages of the invention will be apparent from the following detailed description of the invention, and from the claims.

BRIEF DESCRIPTION OF THE FIGURES

Figures 1A-WW shows the nucleotide sequences for human proteases oriented in a 5' to 3' direction (SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39,

SEQ ID NO:40, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:44,
SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49,
SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:54,
SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:58, and SEQ ID
5 NO:59). In the sequences, N means any nucleotide.

Figure 2A-S shows the amino acid sequences for the human proteases
encoded by SEQ ID No. 1-59 in the direction of translation (SEQ ID NO:60, SEQ
ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ
ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70 SEQ
10 ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ
ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ
ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ
ID NO:86, SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ
ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ
15 ID NO:96, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ
ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105,
SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:109, SEQ ID
NO:110, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114,
SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118). In the
20 sequences, X means any amino acid.

DETAILED DESCRIPTION OF THE INVENTION

The following description of the background of the invention is provided to
aid in understanding the invention, but is not admitted to be or to describe prior art
25 to the invention.

Proteases are enzymes capable of severing the amino acid backbone of other
proteins, and are involved in a large number of diverse processes within the body.
Their normal functions include modulation of apoptosis (caspases) (Salvesen and

Dixon, *Cell*, 1997, 91:443-46), control of blood pressure (renin, angiotensin-converting enzymes) (van Hooft *et al.*, 1991, *N Engl J Med.* 324(19):1305-11, and chapters 254 and 359 in Barrett *et al.*, Handbook of Proteolytic Enzymes, 1998, Academic Press, San Diego), tissue remodeling and tumor invasion (collagenase) (Vu *et al.*, 1998, *Cell* 93:411-22, Werb, 1997, *Cell*, 91:439-442), development of Alzheimer's Disease (β -secretase) (De Strooper *et al.*, 1999, *Nature* 398:518-22), protein turnover and cell-cycle regulation (proteosome) (Bastians *et al.*, 1999, *Mol. Biol. Cell.* 10:3927-41, Gottesman, *et al.*, 1997, *Cell*, 91:435-38, Larsen *et al.*, 1997, *Cell*, 91:431-34), inflammation (TNF- α convertase) (Black *et al.*, *Nature*, 1997, 385:729-33); and protein turnover (Bochtler *et al.*, 1999, *Annu. Rev. Biophys Biomol Struct.* 28:295-317). Proteases may be classified into several major groups including serine proteases, cysteine proteases, aspartyl proteases, metalloproteases, threonine proteases, and other proteases.

1. Aspartyl proteases (A1; Prosite number PS00141):

Aspartyl proteases, also known as acid proteases, are a widely distributed family of proteolytic enzymes in vertebrates, fungi, plants, retroviruses and some plant viruses. Aspartate proteases of eukaryotes are monomeric enzymes which consist of two domains. Each domain contains an active site centered on a catalytic aspartyl residue. The two domains most probably evolved from the duplication of an ancestral gene encoding a primordial domain. Enzymes in this class include cathepsin E, renin, presenilin (PS1), and the APP secretases.

Cathepsin E

Cathepsin E is an immunologically discrete aspartic protease found in the gastrointestinal tract (Azuma *et al.*, 1992 *J. Biol. Chem.*, 267:1609-1614). Cathepsin E is an intracellular proteinase that does not appear to be involved in the digestion of

dietary protein. It is found in highest concentration in the surface of epithelial mucus-producing cells of the stomach. It is the first aspartic proteinase expressed in the fetal stomach and is found in more than half of gastric cancers. It appears, therefore, to be an 'oncofetal' antigen. Its association with stomach cancers suggests
5 it may play a role in the development of this disease.

Renin

Released by the juxtaglomerular cells of the kidney, renin catalyzes the first step in the activation pathway of angiotensinogen--a cascade that can result in
10 aldosterone release, vasoconstriction, and increase in blood pressure. Renin cleaves angiotensinogen to form angiotensin I, which is converted to angiotensin II by angiotensin I converting enzyme, an important regulator of blood pressure and electrolyte balance. Renin occurs in other organs than the kidney, e.g., in the brain, where it is implicated in the regulation of numerous activities.

15

Presenilin proteins

Alzheimer's disease (AD) patients with an inherited form of the disease carry mutations in the presenilin proteins (PSEN1; PSEN2) or the amyloid precursor protein (APP). These disease-linked mutations result in increased production of the
20 longer form of amyloid-beta (main component of amyloid deposits found in AD brains) (Saftig et al., *Eur. Arch. Psychiatry Clin. Neurosci.*, 1999, 249:271-79). Presenilins are postulated to regulate APP processing through their effects on gamma-secretase, an enzyme that cleaves APP (Cruts et al., 1998, *Hum. Mutat.*, 11:183-190, Haass et al., *Science*, 1999, 286:916-19). Also, it is thought that the
25 presenilins are involved in the cleavage of the Notch receptor, such that that they either directly regulate gamma-secretase activity or themselves are protease enzymes

(De Strooper et al., *Nature*, 1999, 398:518-22). Two alternative transcripts of PSEN2 have been identified (Sato et al., 1999, *J. Neurochem.* 72(6):2498-505). Point mutations in the PS1 gene result in a selective increase in the production of the amyloidogenic peptide amyloid-beta (1-42) by proteolytic processing of the amyloid precursor protein (APP) (Lemere et al., 1996, *Nat Med* 2(10):1146-50). The possible role of PS1 in normal APP processing was studied by De Strooper et al. (*Nature* 391: 387-390, 1998) in neuronal cultures derived from PS1-deficient mouse embryos. They found that cleavage by alpha- and beta-secretase of the extracellular domain of APP was not affected by the absence of PS1, whereas cleavage by gamma-secretase of the transmembrane domain of APP was prevented, causing C-terminal fragments of APP to accumulate and a 5-fold drop in the production of amyloid peptide. Pulse-chase experiments indicated that PS1 deficiency specifically decreased the turnover of the membrane-associated fragments of APP. Thus, PS1 appears to facilitate a proteolytic activity that cleaves the integral membrane domain of APP. The results indicated to the authors that mutations in PS1 that manifest clinically cause a gain of function, and that inhibition of PS1 activity is a potential target for anti-amyloidogenic therapy in Alzheimer disease.

Beta-secretase

Beta-secretase, expressed specifically in the brain, is responsible for the proteolytic processing of the amyloid precursor protein (APP) associated with Alzheimer's disease (Potter et al., 2000, *Nat. Biotechnol* 18(2):125-26). It cleaves at the amino terminus of the beta peptide sequence, between residues 671 and 672 of APP, leading to the generation and extracellular release of beta-cleaved soluble APP, and a carboxyterminal fragment that is later released by gamma-secretase (Kinberly et al., 2000 *J. Biol. Chem.* 275(5):3173-78). Yan et al. (*Nature*, 1999, 402:533-37) identified a new membrane-bound aspartyl protease (Asp2) with beta-secretase activity. The Asp2 gene is expressed widely in brain and other tissues. Decreasing the expression of Asp2 in cells reduces amyloid beta-peptide production

and blocks the accumulation of the carboxy-terminal APP fragment that is created by beta-secretase cleavage. Asp2 is a new protein target for drugs that are designed to block the production of amyloid beta-peptide peptide and the consequent formation of amyloid plaque in Alzheimer's disease.

5 Two aspartyl proteases involved in human placentation have recently been isolated: decidual aspartyl protease (DAP-1) and DAP-2 (Moses et al., *Mol. Hum Reprod.* 1999, 5:983-89).

Another member of the aspartyl peptidase family is HIV-1 retropepsin, from the human immunodeficiency virus type 1. This enzyme is vital for processing of
10 the viral polyprotein and maturation of the mature virion.

2. Cysteine proteases

Another class of proteases which perform a wide variety of functions within the body are the cysteine proteases. Among their roles are the processing of
15 precursor proteins, and intracellular degradation of proteins marked for disposal via the ubiquitin pathway. Eukaryotic cysteine proteases are a family of proteolytic enzymes which contain an active site cysteine. Catalysis proceeds through a thioester intermediate and is facilitated by a nearby histidine side chain; an asparagine completes the essential catalytic triad. Peptidases in this family with
20 important roles in disease include the caspases, calpain, hedgehog, and Ubiquitin hydrolases.

Cysteine proteases are produced by a large number of cells including those of the immune system (macrophages, monocytes, etc.). These immune cells exercise their protective role in the body, in part, by migrating to sites of
25 inflammation and secreting molecules, among the secreted molecules are cysteine proteases.

Under some conditions, the inappropriate regulation of cysteine proteases of the immune system can lead to autoimmune diseases such as rheumatoid arthritis. For example, the over-secretion of the cysteine protease cathepsin C causes the

degradation of elastin, collagen, laminin, and other structural proteins found in bones. Bone subjected to this inappropriate digestion is more susceptible to metastasis.

Caspase (C14) - apoptosis

5 A cascade of protease reactions is believed to be responsible for the apoptotic changes observed in mammalian cells undergoing programmed cell death. This cascade involves many members of the aspartate-specific cysteine proteases of the caspase family, including caspases 2, 3, 6, 7, 8 and 10 (Salvesen and Dixit, *Cell* 1997, 91:443-446). Cancer cells that escape apoptotic signals, generated by
10 cytotoxic chemotherapeutics or loss of normal cellular survival signals (as in metastatic cells), can go on to develop palpable tumors.

Other caspases are also involved in the activation of pro-inflammatory cytokines. Caspase 1 specifically processes the precursors of IL-1 β , and IL-18 (interferon- γ -inducing factor)(Salvesen and Dixit *Cell* 1997).

15 Calpain (C2) - axonal death, dystrophies

Calcium-dependent cysteine proteases, collectively called calpain, are widely distributed in mammalian cells (Wang, 2000, *Trends Neurosci.* 23(1):20-26). The calpains are nonlysosomal intracellular cysteine proteases. The mammalian calpains
20 include 2 ubiquitous proteins, CAPN1 and CAPN2, as well as 2 stomach-specific proteins, and CAPN3, which is muscle-specific (Hérasse *et al.*, 1999, *Mol. Cell. Biol.* 19(6):4047-55). The ubiquitous enzymes consist of heterodimers with distinct large subunits associated with a common small subunit, all of which are encoded by different genes. The large subunits of calpains can be subdivided into 4 domains;
25 domains I and III, whose functions remain unknown, show no homology with known proteins. The former, however, may be important for the regulation of the proteolytic activity. Domain II shows similarity with other cysteine proteases, which share histidine, cysteine, and asparagine residues at their active sites. Domain IV is calmodulin-like. CAPN5 and CAPN6 differ from previously identified

vertebrate calpains in that they lack a calmodulin-like domain IV (Ohno *et al.*, 1990, *Cytogenet. Cell Genet.* 53(4):225-29).

Mutations in the CAPN3 gene have been associated with limb-girdle muscular dystrophy, type 2A (LGMD2A) (Allamand *et al.*, 1995, *Hum. Molec. Genet.* 4:459-463). The slowly progressive muscle weakness associated with this disease is usually first evident in the pelvic girdle and then spreads to the upper limbs while sparing facial muscles. Calpain has also been implicated in the development of hyperactive Cdk5 leading to neuronal cell death associated with Alzheimer's disease (Patrick *et al.*, 1999, *Nature* 402:615-622).

Hedgehog (C46) – Cancer

The organization and morphology of the developing embryo are established through a series of inductive interactions. One family of vertebrate genes has been described related to the *Drosophila* gene 'hedgehog' (hh) that encodes inductive signals during embryogenesis (Johnson and Tabin, 1997, *Cell* 90:979-990). 'Hedgehog' encodes a secreted protein that is involved in establishing cell fates at several points during *Drosophila* development (Marigo *et al.*, 1995, *Genomics* 28:44-51). There are 3 known mammalian homologs of hh: Sonic hedgehog (Shh), Indian hedgehog (Ihh), and desert hedgehog (Dhh) (Johnson and Tabin, 1997, *Cell* 90:979-990). Like its *Drosophila* cognate, Shh encodes a signal that is instrumental in patterning the early embryo. It is expressed in Hensen's node, the floorplate of the neural tube, the early gut endoderm, the posterior of the limb buds, and throughout the notochord (Chiang *et al.*, 1996, *Nature* 383:407-413). It has been implicated as the key inductive signal in patterning of the ventral neural tube, the anterior-posterior limb axis, and the ventral somites. Oro *et al.* ("Basal cell carcinomas in mice overexpressing sonic hedgehog." *Science* 276: 817-821, 1997) showed that transgenic mice overexpressing SHH in the skin developed many features of the basal cell nevus syndrome, demonstrating that SHH is sufficient to induce basal cell carcinomas (BCCs) in mice. The data suggested that SHH may

have a role in human tumorigenesis. Activating mutations of SHH or another 'hedgehog' gene may be an alternative pathway for BCC formation in humans. The human mutation his133tyr (his134tyr in mouse) is a candidate. It is distinct from loss-of-function mutations reported for individuals with holoprosencephaly (Oro *et al.*, 1997, *Science* 276:817-821). His133 lies adjacent in the catalytic site to his134, one of the conserved residues thought to be necessary for catalysis. SHH may be a dominant oncogene in multiple human tumors, a mirror of the tumor suppressor activity of the opposing 'patched' (PTCH) gene (Aszterbaum *et al.*, 1998, *J. Invest. Derm.* 110:885-888). The rapid and frequent appearance of Shh-induced tumors in the mice suggested that disruption of the SHH-PTC pathway is sufficient to create BCCs.

Members of the vertebrate hedgehog family (Sonic, Indian, and Desert) have been shown to be essential for the development of various organ systems, including neural, somite, limb, skeletal, and for male gonad morphogenesis. Desert hedgehog is expressed in the developing retina, whereas Indian hedgehog (Ihh) is expressed in the developing and mature retinal pigmented epithelium beginning at embryonic day 13 (Levine *et al.*, *J. Neurosci.*, 1997, 17(16):6277-88). Dhh has also been implicated in having a role in the regulation of spermatogenesis. Sertoli cell precursors express Sry, sex determining gene, which leads to testis development in mammals. Dhh expression is initiated in Sertoli cell precursors shortly after the activation of Sry and persists in the testis into the adult. Bitgood *et al.* (*Curr. Biol.*, 1996, 6(3):298-304) disclose that female mice homozygous for a Dhh-null mutation show no obvious phenotype, whereas males are viable but infertile having a complete absence of mature sperm, demonstrating that Dhh signaling plays an essential role in the regulation of mammalian spermatogenesis. Dhh has also been found to have a role in the and maintenance of protective nerve sheaths endo-, peri- and epineurium. In Dhh knockout mice, the connective tissue sheaths in adult nerves appear highly abnormal by electron microscopy. Mirsky *et al.*, (*Ann. N.Y. Acad. Sci.*, 1999, 883:196-202) demonstrate that Dhh signaling from Schwann cells to the

mesenchyme is involved in the formation of a morphologically and functionally normal perineurium.

Recent advances in developmental and molecular biology during embryogenesis and organogenesis have provided new insights into the mechanism of bone formation. Iwasaki *et al.*, (*J. Bone Joint Surg. Br.*, 1999, 81(6):1076-82) demonstrate that Indian Hedgehog (Ihh) is expressed in cartilage cell precursors and later in mature and hypertrophic chondrocytes. Ihh plays a critical role in the morphogenesis of the vertebrate skeleton. Becker *et al.* (*Dev. Biol.*, 1997, 187(2):298-310) provide data which suggests that Ihh is also involved in mediating differentiation of extraembryonic endoderm during early mouse embryogenesis. Short limbed dwarfism, with decreased chondrocyte proliferation and extensive hypertrophy are the results of targeted deletion of Ihh (Karp *et al.*, 2000, Development 127(3):543-48). The expression of Ihh mRNA and protein is unregulated dramatically as F9 cells differentiate in response to retinoic acid, into either parietal endoderm or embryoid bodies, containing an outer visceral endoderm layer. RT-PCR analysis of blastocyst outgrowth cultures demonstrates that whereas little or no Ihh message is present in blastocysts, significant levels appear upon subsequent days of culture, coincident with the emergence of parietal endoderm cells.

20

Ubiquitin hydrolases (C12) - apoptosis, checkpoint integrity

14 genes in this patent belong to the ubiquitin hydrolase family, SEQID:5, SEQID:6, SEQID:7, SEQID:8, SEQID:9, SEQID:10, SEQID:11, SEQID:12, SEQID:13, SEQID:14, SEQID:15, SEQID:16, SEQID:17, SEQID:18. The polypeptides encoded by these genes may have one or more of the following activities.

25

Ubiquitin carboxyl-terminal hydrolases (3.1.2.15) (deubiquitinating enzymes) are thiol proteases that recognize and hydrolyze the peptide bond at the C-terminal glycine of ubiquitin. These enzymes are involved in the processing of

poly-ubiquitin precursors as well as that of ubiquitinated proteins. In eukaryotic cells, the covalent attachment of ubiquitin to proteins plays a role in a variety of cellular processes. In many cases, ubiquitination leads to protein degradation by the 26S proteasome. Protein ubiquitination is reversible, and the removal of ubiquitin is
5 catalyzed by deubiquitinating enzymes, or DUBs. A defect in these enzymes, catalyzing the removal of ubiquitin from ubiquitinated proteins, may be characteristic of neurodegenerative diseases such as Alzheimer's, Parkinson's, progressive supranuclear palsy, and Pick's and Kuf's disease.

10 Papain (C1) – cathepsins K, S and B, - bone resorption, Ag processing (Prosite PS00139).

One gene in this patent belongs to the Papain family, SEQID:4. The polypeptide encoded by this gene may have one or more of the following activities.

Cathepsin K, a member of the papain family of peptidases, is involved in
15 osteoclastic resorption. It plays an important role in extracellular degradation and may have a role in disorders of bone remodeling, such as pycnodysostosis, an autosomal recessive osteochondrodysplasia characterized by osteosclerosis and short stature. Antigen presentation by major histocompatibility complex (MHC) class II molecules requires the participation of different proteases in the endocytic route to
20 degrade endocytosed antigens as well as the MHC class II-associated invariant chain. Only cathepsin S, a member of the papain family, appears to be essential for complete destruction of the invariant chain. Cathepsin B is overexpressed in tumors of the lung, prostate, colon, breast, and stomach. Hughes et al. (*Proc. Nat. Acad. Sci.* 95: 12410-12415, 1998) found an amplicon at 8p23-p22 that resulted in cathepsin B
25 overexpression in esophageal adenocarcinoma. Abundant extracellular expression of CTSB protein was found in 29 of 40 (72.5%) of esophageal adenocarcinoma specimens by use of immunohistochemical analysis. The findings were thought to support an important role for CTSB in esophageal adenocarcinoma and possibly in other tumors.

Cathepsin B, a lysosomal protease, is being studied as a prognostic marker in various cancers (breast, pulmonary adenocarcinomas).

Cysteine Protease AEP

- 5 The cysteine protease AEP plays another role in the immune functions. It has been implicated in the protease step required for antigen processing in B cells. (Manoury *et al. Nature* 396:695-699 (1998))

Hepatitis A viral protease (C3E)

- 10 The Hepatitis A genome encodes a cysteine protease required for enzymatic cleavages *in vivo* to yield mature proteins (Wang, 1999, *Prog. Drug Res.* 52:197-219). This enzyme and its homologs in other viruses (such as hepatitis E virus) are potential targets for chemotherapeutic intervention.

15 3. Metalloproteases

Collagenase (M10) – invasion

Two genes in this patent are members of the M10 family, SEQID:19 and SEQID:20. The polypeptides encoded by these genes may have one or more of the following activities.

- 20 Matrix degradation is an essential step in the spread of cancer. The 72- and 92-kD type IV collagenases are members of a group of secreted zinc metalloproteases which, in mammals, degrade the collagens of the extracellular matrix. Other members of this group include interstitial collagenase and stromelysin (Nagase *et al.*, 1992, *Matrix Suppl.* 1:421-424). By targeted disruption in embryonic stem cells, Vu *et al.* (*Cell*, 1998, 934:11-22) created homozygous mice with a null
25 mutation in the MMP9/gelatinase B gene. These mice exhibited an abnormal pattern of skeletal growth plate vascularization and ossification. Growth plates from MMP9-null mice in culture showed a delayed release of an angiogenic activator, establishing a role for this proteinase in controlling angiogenesis.

MMP2 (gelatinase A) have been associated with the aggressiveness of human cancers (Chenard *et al.*, 1999, *Int. J. Cancer*, 82:208-12). In a study comparing basal cell carcinomas (BCC) with the more aggressive squamous cell carcinomas (SCC), both MMP2 and MMP9 were expressed at a higher level in SCC (Dumas *et al.*, 1999, *Anticancer Res.*, 19(4B):2929-38). Additionally, expression of MMP2 and MMP9 in T lymphocytes has recently been shown to be modulated by the Ras/MAP kinase signaling pathways (Esparza *et al.*, 1999, *Blood*, 94:2754-66) (see also, Li *et al.*, 1998, *Biochim. Biophys. Acta*, 1405:110-20).

10 ADAMs (M12) - TNF, inflammation, growth factor processing

The ADAM peptidases are a family of proteins containing a disintegrin and metalloproteinase (ADAM) domain (Werb and Yan, *Science*, 1998, 282:1279-1280). Members of this family are cell surface proteins with a unique structure possessing both potential adhesion and protease domains (Primakoff and Myles, *Trends in Genet.*, 2000, 16:83-87). Activity of these proteases can be linked to TNF, inflammation, and/or growth factor processing.

ADAM proteases have also been characterized as having a pro- and metalloproteinase domain, a disintegrin domain, a cysteine-rich region and an EGF repeat (Blobel, 1997, *Cell*, 90:589-592 which is hereby incorporated herein by reference in its entirety including any figures, tables, or drawings). They have been associated with the release from the plasma membrane of numerous proteins including Tumor Necrosis Factor- α (TNF- α), kit-ligand, TGF α , Fas-ligand, cytokine receptors such as the Il-6 receptor and the NGF receptor, as well as adhesion proteins such as L-selectin, and the b amyloid precursor proteins (Blobel, 1997, *Cell*, 90:589-592).

Tumor necrosis factor- α is synthesized as a proinflammatory cytokine from a 233-amino acid precursor. Conversion of the membrane-bound precursor to a secreted mature protein is mediated by a protease termed TNF- α convertase. TNF- α

is involved in a variety of diseases. ADAM17, which contains a disintegrin and metalloproteinase domains, is also called 'tumor necrosis factor- α converting enzyme' (TACE) (Black *et al.*, *Nature*, 1997, 385:729-33). The gene encodes an 824-amino acid polypeptide containing the features of the ADAM family: a
5 secretory signal sequence, a disintegrin domain, and a metalloprotease domain. Expression studies showed that the encoded protein cleaves precursor tumor necrosis factor- α to its mature form. This enzyme may also play a role in the processing of Transforming Growth Factor- α (TGF- α), as mice which lack the gene are similar in phenotype to those that lack TGF- α (Peschon *et al.*, *Science*,
10 282:1281-1284).

Neprylisin (M13) - Endothelin-converting enzyme

One gene in this patent, SEQID:21, is a member of this family. The polypeptide encoded by this gene may have one or more of the following activities.
15 Neprylisin, a metallopeptidase active in degradation of enkephalins and other bioactive peptides, is a drug target in hypertension and renal disease (Oefner, *et al.*, *J. Mol. Biol.*, 2000, 296:341-49).

Carboxypeptidase (M14) - Neurotransmitter processing

20 Three genes in this application are Zn carboxypeptidases, SEQID:1, SEQID:2, and SEQID:3. The polypeptides encoded by these genes may have one or more of the following activities.

Carboxypeptidases specifically remove COOH-terminal basic amino acids (arginine or lysine). They have important functions in many biologic processes,
25 including activation, inactivation, or modulation of peptide hormone activity, neurotransmitter processing, and alteration of physical properties of proteins and enzymes.

Dipeptidase (M2) – ACE

One protease in this patent is a member of the M2 family: SEQID:22. The polypeptide encoded by this gene may have one or more of the following activities.

- 5 Angiotensin I converting enzyme (EC 3.4.15.1), or kininase II, is adipeptidyl carboxypeptidase that plays an important role in blood pressure regulation and electrolyte balance by hydrolyzing angiotensin I into angiotensin II, a potent vasopressor, and aldosterone-stimulating peptide. The enzyme is also able to inactivate bradykinin, a potent vasodilator. Although angiotensin-converting
10 enzyme has been studied primarily in the context of its role in blood pressure regulation, this widely distributed enzyme has many other physiologic functions. There are two forms of ACE: a testis-specific isozyme and a somatic isozyme which has two active centers.

15 Matrix metalloproteases (M10B) – tissue remodeling and inflammation

- The matrix metalloproteases (MMPs) are a family of related matrix-degrading enzymes that are important in tissue remodeling and repair during development and inflammation. Abnormal expression is associated with various diseases such as tumor invasiveness, arthritis, and atherosclerosis. MMP activity
20 may also be related to tobacco-induced pulmonary emphysema.

- The matrix metalloproteases (MMPs) are a family of related matrix-degrading enzymes that are important in tissue remodeling and repair during development and inflammation (Belotti *et al.*, 1999, *Int. J. Biol. Markers* 14(4):232-38). Abnormal expression is associated with various diseases such as tumor
25 invasiveness (Johansson and Kahari, 2000, *Histol. Histopathol.* 15(1):225-37), arthritis (Malemud *et al.*, 1999, *Front. Biosci.* 4:D762-71), and atherosclerosis (Nagase, 1997, *Biol. Chem.* 378(3-4):151-60). MMP activity may also be related to

tobacco-induced pulmonary emphysema (Dhami *et al.*, *Am. J. Respir. Cell Mol. Biol.*, 2000, 22:244-52).

SREBP Protease (M50)

5 The sterol regulatory element-binding proteins protease functions in the intra-membrane proteolysis and release of sterol-regulatory binding proteins (SREBPs) (Duncan *et al.*, 1997, *J. Biol. Chem.* 272:12778-85). SREBPs activate genes of cholesterol and fatty acid metabolism, making the SREBP protease an attractive target for therapeutic modulation (Brown *et al.*, 1997, *Cell* 89:331-340).

10

Metalloprotease processing of growth factors

 In addition to the processing of TGF- α described above, metalloproteases have been directly demonstrated to be active in the processing of the precursor of other growth factors such as heparin-binding EGF (proHB-EGF) (Izumi *et al.*,
15 *EMBO J*, 1998,17:7260-72), and amphiregulin (Brown *et al.*, 1998, *J. Biol. Chem.*, 27:17258-68).

 Additionally, metalloproteases have recently been shown to be instrumental in the communication whereby stimulation of a GPCR pathway results in stimulation of the MAP kinase pathway (Prenzel *et al.*, 1999, *Nature*, 402:884-888).
20 The growth factor intermediate in the pathway, HB-EGF is released by the cell in a proteolytic step regulated by the GPCR pathway involving an uncharacterized metalloprotease. After release, the HB-EGF is bound by the extracellular matrix and then presented to the EGF receptors on the surface, resulting in the activation of the MAP kinase pathway (Prenzel *et al.*, 1999, *Nature*, 402:884-888).

25 A recent study by Gallea-Robache *et al.* (1997) has also implicated a metalloprotease family displaying different substrate specificities in the shedding of other growth factors including macrophage colony-stimulating factor (M-CSF) and stem cell factor (SCF) (Gallea-Robache *et al.*, 1997, *Cytokine* 9:340-46). The

shedding of M-CSF (also known as CSF-1) has been linked to activation of Protein Kinase C by phorbol esters (Stein *et al.*, 1991, *Oncogene*, 6:601-05).

4. Serine Proteases

5 The serine proteases are a class which includes trypsin, kallikrein, chymotrypsin, elastase, thrombin, tissue plasminogen activator (tPA), urokinase plasminogen activator (uPA), plasmin (Werb, *Cell*, 1997, 91:439-442), kallikrein (Clements, *Biol. Res.*, 1998, 31:151-59), and cathepsin G (Shamamian *et al.*, *Surgery*, 2000, 127:142-47). These proteases have in common a well-conserved
10 catalytic triad of amino acid residues in their active site consisting of histidine-57, aspartic acid-102, and serine-195 (using the chymotrypsin numbering system). Serine protease activity has been linked to coagulation and they may have use as tumor markers.

 Serine proteases can be further subclassified by their specificity in substrates.
15 The elastases prefer to cleave substrates adjacent to small aliphatic residues such as valine, chymases prefer to cleave near large aromatic hydrophobic residues, and tryptases prefer positively charged residues. One additional class of serine protease has been described recently which prefers to cleave adjacent to a proline. This prolyl endopeptidase has been implicated in the progression of memory loss in
20 Alzheimer's patients (Toide *et al.*, 1998, *Rev. Neurosci.* 9(1):17-29).

 A partial list of proteases known to belong to this large and important family include: blood coagulation factors VII, IX, X, XI and XII; thrombin; plasminogen; complement components C1r, C1s, C2; complement factors B, D and I; complement-activating component of RA-reactive factor; elastases 1, 2, 3A, 3B
25 (protease E); hepatocyte growth factor activator; glandular (tissue) kallikreins including EGF-binding protein types A, B, and C; NGF- γ chain, γ -renin, and prostate specific antigen (PSA); plasma kallikrein; mast cell proteases; myeloblastin (proteinase 3) (Wegener's autoantigen); plasminogen activators (urokinase-type, and tissue-type); and the tryptases I, II, III, and IV. These peptidases play key roles in

coagulation, tumorigenesis, control of blood pressure, release of growth factors, and other roles. (<http://www.babraham.co.uk/Merops/Merops.htm>).

Proteases of the trypsin family in this patent include SGPr434, SEQID:24; SGPr446_1, SEQID:25; SGPr447, SEQID:26; SGPr432_1, SEQID:27; SGPr529, SEQID:28; SGPr428_1, SEQID:29; SGPr425, SEQID:30; SGPr548, SEQID:31; SGPr396, SEQID:32; SGPr426, SEQID:33; SGPr552, SEQID:34; SGPr405, SEQID:35; SGPr485_1, SEQID:36; SGPr534, SEQID:37; SGPr390, SEQID:38; SGPr521, SEQID:39; SGPr530_1, SEQID:40; SGPr520, SEQID:41; SGPr455, SEQID:42; SGPr507_2, SEQID:43; SGPr559, SEQID:44; SGPr567_1, SEQID:45; SGPr479_1, SEQID:46; SGPr489_1, SEQID:47; SGPr465_1, SEQID:48; SGPr524_1, SEQID:49; SGPr422, SEQID:50; SGPr538, SEQID:51; SGPr527_1, SEQID:52; SGPr542, SEQID:53; SGPr551, SEQID:54; SGPr451, SEQID:55; SGPr452_1, SEQID:56; SGPr504, SEQID:57; SGPr469, SEQID:58; SGPr400, SEQID:59. SEQID:23 is a serine protease of the subtilase sub-family. Limited proteolysis of most large protein precursors is carried out in vivo by the subtilisin-like pro-protein convertases. Many important biological processes such as peptide hormone synthesis, viral protein processing and receptor maturation involve proteolytic processing by these enzymes, making them potential targets for the development of novel therapeutic agents (Bergeron F, J Mol Endocrinol 2000 Feb;24(1):1-22)

5. Threonine peptidases (T1) – (Prosite PDOC00326/PDOC00668)

Proteasomal subunits (T1A)

The proteasome is a multicatalytic threonine proteinase complex involved in ATP/ubiquitin dependent non-lysosomal proteolysis of cellular substrates. It is responsible for selective elimination of proteins with aberrant structures, as well as naturally occurring short-lived proteins related to metabolic regulation and cell-cycle progression (Momand *et al.*, 2000, *Gene* 242(1-2):15-29, Bochtler *et al.*, 1999, *Annu. Rev. Biophys Biomol Struct.* 28:295-317). The proteasome inhibitor

lactacystin reversibly inhibits proliferation of human endothelial cells, suggesting a role for proteasomes in angiogenesis (Kumeda, *et al.*, *Anticancer Res.* 1999 Sep-Oct;19(5B):3961-8). Another important function of the proteasome in higher vertebrates is to generate the peptides presented on MHC-class 1 molecules to circulating lymphocytes (Castelli *et al.*, 1997, *Int. J. Clin. Lab. Res.* 27(2):103-10). The proteasome has a sedimentation coefficient of 26S and is composed of a 20S catalytic core and a 22S regulatory complex. Eukaryotic 20S proteasomes have a molecular mass of 700 to 800 kD and consist of a set of over 15 kinds of polypeptides of 21 to 32 kD. All eukaryotic 20S proteasome subunits can be classified grossly into 2 subfamilies, α and β , by their high similarity with either the α or β subunits of the archaeobacterium *Thermoplasma acidophilum* (Mayr *et al.*, 1999, *Biol. Chem.* 380(10):1183-92). Several of the components have been identified as threonine peptidases, suggesting that this class of peptidases plays a key role in regulating metabolic pathways and cell-cycle progression, among other functions (Yorgin *et al.*, 2000, *J. Immunol.* 164(6):2915-23).

6. Peptidases of Unknown Catalytic Mechanism

The prenyl-protein specific protease responsible for post-translational processing of the Ras proto-oncogene and other prenylated proteins falls into this class. This class also includes several viral peptidases that may play a role in mammalian infection, including cardiovirus endopeptidase 2A (encephalomyocarditis virus) (Molla *et al.*, 1993, *J. Virol.* 67(8):4688-95), NS2-3 protease (hepatitis C virus) (Blight *et al.*, 1998, *Antivir. Ther.* 3(Suppl 3):71-81), endopeptidase (infectious pancreatic necrosis virus) (Lejal *et al.*, *J. Gen. Virol.*, 2000, 81:983-992), and the Npro endopeptidase (hog cholera virus) (Tratschin *et al.*, 1998, *J. Virol.* 72(9):7681-84).

Nucleic Acid Probes, Methods, and Kits for Detection of Proteases

A nucleic acid probe of the present invention may be used to probe an appropriate chromosomal or cDNA library by usual hybridization methods to obtain other nucleic acid molecules of the present invention. A chromosomal DNA or
5 cDNA library may be prepared from appropriate cells according to recognized methods in the art (*cf.* "Molecular Cloning: A Laboratory Manual", second edition, Cold Spring Harbor Laboratory, Sambrook, Fritsch, & Maniatis, eds., 1989).

In the alternative, chemical synthesis can be carried out in order to obtain nucleic acid probes having nucleotide sequences which correspond to N-terminal
10 and C-terminal portions of the amino acid sequence of the polypeptide of interest. The synthesized nucleic acid probes may be used as primers in a polymerase chain reaction (PCR) carried out in accordance with recognized PCR techniques, essentially according to PCR Protocols, "A Guide to Methods and Applications", - Academic Press, Michael, *et al.*, eds., 1990, utilizing the appropriate chromosomal
15 or cDNA library to obtain the fragment of the present invention.

One skilled in the art can readily design such probes based on the sequence disclosed herein using methods of computer alignment and sequence analysis known in the art ("Molecular Cloning: A Laboratory Manual", 1989, *supra*). The hybridization probes of the present invention can be labeled by standard labeling
20 techniques such as with a radiolabel, enzyme label, fluorescent label, biotin-avidin label, chemiluminescence, and the like. After hybridization, the probes may be visualized using known methods.

The nucleic acid probes of the present invention include RNA, as well as DNA probes, such probes being generated using techniques known in the art. The
25 nucleic acid probe may be immobilized on a solid support. Examples of such solid supports include, but are not limited to, plastics such as polycarbonate, complex carbohydrates such as agarose and sepharose, and acrylic resins, such as polyacrylamide and latex beads. Techniques for coupling nucleic acid probes to such solid supports are well known in the art.

The test samples suitable for nucleic acid probing methods of the present invention include, for example, cells or nucleic acid extracts of cells, or biological fluids. The samples used in the above-described methods will vary based on the assay format, the detection method and the nature of the tissues, cells or extracts to be assayed. Methods for preparing nucleic acid extracts of cells are well known in the art and can be readily adapted in order to obtain a sample which is compatible with the method utilized.

One method of detecting the presence of nucleic acids of the invention in a sample comprises (a) contacting said sample with the above-described nucleic acid probe under conditions such that hybridization occurs, and (b) detecting the presence of said probe bound to said nucleic acid molecule. One skilled in the art would select the nucleic acid probe according to techniques known in the art as described above. Samples to be tested include but should not be limited to RNA samples of human tissue.

A kit for detecting the presence of nucleic acids of the invention in a sample comprises at least one container means having disposed therein the above-described nucleic acid probe. The kit may further comprise other containers comprising one or more of the following: wash reagents and reagents capable of detecting the presence of bound nucleic acid probe. Examples of detection reagents include, but are not limited to radiolabelled probes, enzymatic labeled probes (horseradish peroxidase, alkaline phosphatase), and affinity labeled probes (biotin, avidin, or streptavidin). Preferably, the kit further comprises instructions for use.

In detail, a compartmentalized kit includes any kit in which reagents are contained in separate containers. Such containers include small glass containers, plastic containers or strips of plastic or paper. Such containers allow the efficient transfer of reagents from one compartment to another compartment such that the samples and reagents are not cross-contaminated and the agents or solutions of each container can be added in a quantitative fashion from one compartment to another. Such containers will include a container which will accept the test sample, a

container which contains the probe or primers used in the assay, containers which contain wash reagents (such as phosphate buffered saline, Tris-buffers, and the like), and containers which contain the reagents used to detect the hybridized probe, bound antibody, amplified product, or the like. One skilled in the art will readily recognize that the nucleic acid probes described in the present invention can readily be
5 incorporated into one of the established kit formats which are well known in the art.

DNA Constructs Comprising a Protease Nucleic Acid Molecule and Cells
Containing These Constructs.

10 The present invention also relates to a recombinant DNA molecule comprising, 5' to 3', a promoter effective to initiate transcription in a host cell and the above-described nucleic acid molecules. In addition, the present invention relates to a recombinant DNA molecule comprising a vector and an above-described nucleic acid molecule. The present invention also relates to a nucleic acid molecule
15 comprising a transcriptional region functional in a cell, a sequence complementary to an RNA sequence encoding an amino acid sequence corresponding to the above-described polypeptide, and a transcriptional termination region functional in said cell. The above-described molecules may be isolated and/or purified DNA molecules.

20 The present invention also relates to a cell or organism that contains an above-described nucleic acid molecule and thereby is capable of expressing a polypeptide. The polypeptide may be purified from cells which have been altered to express the polypeptide. A cell is said to be "altered to express a desired polypeptide" when the cell, through genetic manipulation, is made to produce a
25 protein which it normally does not produce or which the cell normally produces at lower levels. One skilled in the art can readily adapt procedures for introducing and expressing either genomic, cDNA, or synthetic sequences into either eukaryotic or prokaryotic cells.

A nucleic acid molecule, such as DNA, is said to be "capable of expressing" a polypeptide if it contains nucleotide sequences which contain transcriptional and translational regulatory information and such sequences are "operably linked" to nucleotide sequences which encode the polypeptide. An operable linkage is a
5 linkage in which the regulatory DNA sequences and the DNA sequence sought to be expressed are connected in such a way as to permit gene sequence expression. The precise nature of the regulatory regions needed for gene sequence expression may vary from organism to organism, but shall in general include a promoter region which, in prokaryotes, contains both the promoter (which directs the initiation of
10 RNA transcription) as well as the DNA sequences which, when transcribed into RNA, will signal synthesis initiation. Such regions will normally include those 5'-non-coding sequences involved with initiation of transcription and translation, such as the TATA box, capping sequence, CAAT sequence, and the like.

If desired, the non-coding region 3' to the sequence encoding a protease of
15 the invention may be obtained by the above-described methods. This region may be retained for its transcriptional termination regulatory sequences, such as termination and polyadenylation. Thus, by retaining the 3'-region naturally contiguous to the DNA sequence encoding a protease of the invention, the transcriptional termination signals may be provided. Where the transcriptional termination signals are not
20 satisfactorily functional in the expression host cell, then a 3' region functional in the host cell may be substituted.

Two DNA sequences (such as a promoter region sequence and a sequence encoding a protease of the invention) are said to be operably linked if the nature of the linkage between the two DNA sequences allows the protease sequence to be
25 transcribed, i.e., where the linkage does not (1) result in the introduction of a frame-shift mutation, (2) interfere with the ability of the promoter region sequence to direct the transcription of a gene sequence encoding a protease of the invention, or (3) interfere with the ability of the gene sequence of a protease of the invention to be

transcribed by the promoter region sequence. Thus, a promoter region would be operably linked to a DNA sequence if the promoter were capable of effecting transcription of that DNA sequence. Thus, to express a gene encoding a protease of the invention, transcriptional and translational signals recognized by an appropriate host are necessary.

The present invention encompasses the expression of a gene encoding a protease of the invention (or a functional derivative thereof) in either prokaryotic or eukaryotic cells. Prokaryotic hosts are, generally, very efficient and convenient for the production of recombinant proteins and are, therefore, one type of preferred expression system for proteases of the invention. Prokaryotes most frequently are represented by various strains of *E. coli*. However, other microbial strains may also be used, including other bacterial strains.

In prokaryotic systems, plasmid vectors that contain replication sites and control sequences derived from a species compatible with the host may be used. Examples of suitable plasmid vectors may include pBR322, pUC118, pUC119 and the like; suitable phage or bacteriophage vectors may include λ gt10, λ gt11 and the like; and suitable virus vectors may include pMAM-neo, pKRC and the like. Preferably, the selected vector of the present invention has the capacity to replicate in the selected host cell.

Recognized prokaryotic hosts include bacteria such as *E. coli*, *Bacillus*, *Streptomyces*, *Pseudomonas*, *Salmonella*, *Serratia*, and the like. However, under such conditions, the polypeptide will not be glycosylated. The prokaryotic host must be compatible with the replicon and control sequences in the expression plasmid.

To express a protease of the invention (or a functional derivative thereof) in a prokaryotic cell, it is necessary to operably link the sequence encoding the protease of the invention to a functional prokaryotic promoter. Such promoters may be either constitutive or, more preferably, regulatable (*i.e.*, inducible or derepressible). Examples of constitutive promoters include the *int* promoter of bacteriophage λ , the

- bla* promoter of the β -lactamase gene sequence of pBR322, and the *cat* promoter of the chloramphenicol acetyl transferase gene sequence of pPR325, and the like. Examples of inducible prokaryotic promoters include the major right and left promoters of bacteriophage λ (P_L and P_R), the *trp*, *recA*, *lacZ*, *lacI*, and *gal*
- 5 promoters of *E. coli*, the α -amylase (Ulmanen *et al.*, *J. Bacteriol.* 162:176-182, 1985) and the ζ -28-specific promoters of *B. subtilis* (Gilman *et al.*, *Gene Sequence* 32:11-20, 1984), the promoters of the bacteriophages of *Bacillus* (Gryczan, in: The Molecular Biology of the Bacilli, Academic Press, Inc., NY, 1982), and *Streptomyces* promoters (Ward *et al.*, *Mol. Gen. Genet.* 203:468-478, 1986).
- 10 Prokaryotic promoters are reviewed by Glick (*Ind. Microbiot.* 1:277-282, 1987), Cenatiempo (*Biochimie* 68:505-516, 1986), and Gottesman (*Ann. Rev. Genet.* 18:415-442, 1984).

Proper expression in a prokaryotic cell may also require the presence of a ribosome-binding site upstream of the gene sequence-encoding sequence. Such

15 ribosome-binding sites are disclosed, for example, by Gold *et al.* (*Ann. Rev. Microbiol.* 35:365-404, 1981). The selection of control sequences, expression vectors, transformation methods, and the like, are dependent on the type of host cell used to express the gene. As used herein, "cell", "cell line", and "cell culture" may be used interchangeably and all such designations include progeny. Thus, the words

20 "transformants" or "transformed cells" include the primary subject cell and cultures derived therefrom, without regard to the number of transfers. It is also understood that all progeny may not be precisely identical in DNA content, due to deliberate or inadvertent mutations. However, as defined, mutant progeny have the same functionality as that of the originally transformed cell.

25 Host cells which may be used in the expression systems of the present invention are not strictly limited, provided that they are suitable for use in the expression of the protease polypeptide of interest. Suitable hosts may often include eukaryotic cells. Preferred eukaryotic hosts include, for example, yeast, fungi,

insect cells, mammalian cells either *in vivo*, or in tissue culture. Mammalian cells which may be useful as hosts include HeLa cells, cells of fibroblast origin such as VERO or CHO-K1, or cells of lymphoid origin and their derivatives. Preferred mammalian host cells include SP2/0 and J558L, as well as neuroblastoma cell lines
5 such as IMR 332, which may provide better capacities for correct post-translational processing.

In addition, plant cells are also available as hosts, and control sequences compatible with plant cells are available, such as the cauliflower mosaic virus 35S and 19S, and nopaline synthase promoter and polyadenylation signal sequences.
10 Another preferred host is an insect cell, for example the *Drosophila* larvae. Using insect cells as hosts, the *Drosophila* alcohol dehydrogenase promoter can be used (Rubin, *Science* 240:1453-1459, 1988). Alternatively, baculovirus vectors can be engineered to express large amounts of proteases of the invention in insect cells (Jasny, *Science* 238:1653, 1987; Miller *et al.*, in: *Genetic Engineering*, Vol. 8,
15 Plenum, Setlow *et al.*, eds., pp. 277-297, 1986).

Any of a series of yeast expression systems can be utilized which incorporate promoter and termination elements from the actively expressed sequences coding for glycolytic enzymes that are produced in large quantities when yeast are grown in mediums rich in glucose. Known glycolytic gene sequences can also provide very
20 efficient transcriptional control signals. Yeast provides substantial advantages in that it can also carry out post-translational modifications. A number of recombinant DNA strategies exist utilizing strong promoter sequences and high copy number plasmids which can be utilized for production of the desired proteins in yeast. Yeast recognizes leader sequences on cloned mammalian genes and secretes peptides
25 bearing leader sequences (*i.e.*, pre-peptides). Several possible vector systems are available for the expression of proteases of the invention in a mammalian host.

A wide variety of transcriptional and translational regulatory sequences may be employed, depending upon the nature of the host. The transcriptional and translational regulatory signals may be derived from viral sources, such as

adenovirus, bovine papilloma virus, cytomegalovirus, simian virus, or the like, where the regulatory signals are associated with a particular gene sequence which has a high level of expression. Alternatively, promoters from mammalian expression products, such as actin, collagen, myosin, and the like, may be employed.

- 5 Transcriptional initiation regulatory signals may be selected which allow for repression or activation, so that expression of the gene sequences can be modulated. Of interest are regulatory signals which are temperature-sensitive so that by varying the temperature, expression can be repressed or initiated, or are subject to chemical (such as metabolite) regulation.

- 10 Expression of proteases of the invention in eukaryotic hosts requires the use of eukaryotic regulatory regions. Such regions will, in general, include a promoter region sufficient to direct the initiation of RNA synthesis. Preferred eukaryotic promoters include, for example, the promoter of the mouse metallothionein I gene sequence (Hamer *et al.*, *J. Mol. Appl. Gen.* 1:273-288, 1982); the TK promoter of
15 Herpes virus (McKnight, *Cell* 31:355-365, 1982); the SV40 early promoter (Benoist *et al.*, *Nature* (London) 290:304-31, 1981); and the yeast gal4 gene sequence promoter (Johnston *et al.*, *Proc. Natl. Acad. Sci. (USA)* 79:6971-6975, 1982; Silver *et al.*, *Proc. Natl. Acad. Sci. (USA)* 81:5951-5955, 1984).

- 20 Translation of eukaryotic mRNA is initiated at the codon which encodes the first methionine. For this reason, it is preferable to ensure that the linkage between a eukaryotic promoter and a DNA sequence which encodes a protease of the invention (or a functional derivative thereof) does not contain any intervening codons which are capable of encoding a methionine (*i.e.*, AUG). The presence of such codons results either in the formation of a fusion protein (if the AUG codon is in the same
25 reading frame as the protease of the invention coding sequence) or a frame-shift mutation (if the AUG codon is not in the same reading frame as the protease of the invention coding sequence).

A nucleic acid molecule encoding a protease of the invention and an operably linked promoter may be introduced into a recipient prokaryotic or

eukaryotic cell either as a nonreplicating DNA or RNA molecule, which may either be a linear molecule or, more preferably, a closed covalent circular molecule. Since such molecules are incapable of autonomous replication, the expression of the gene may occur through the transient expression of the introduced sequence.

- 5 Alternatively, permanent expression may occur through the integration of the introduced DNA sequence into the host chromosome.

A vector may be employed which is capable of integrating the desired gene sequences into the host cell chromosome. Cells which have stably integrated the introduced DNA into their chromosomes can be selected by also introducing one or
10 more markers which allow for selection of host cells which contain the expression vector. The marker may provide for prototrophy to an auxotrophic host, biocide resistance, *e.g.*, antibiotics, or heavy metals, such as copper, or the like. The selectable marker gene sequence can either be directly linked to the DNA gene sequences to be expressed, or introduced into the same cell by co-transfection.

- 15 Additional elements may also be needed for optimal synthesis of mRNA. These elements may include splice signals, as well as transcription promoters, enhancers, and termination signals. cDNA expression vectors incorporating such elements include those described by Okayama (*Mol. Cell. Biol.* 3:280-289, 1983).

The introduced nucleic acid molecule can be incorporated into a plasmid or
20 viral vector capable of autonomous replication in the recipient host. Any of a wide variety of vectors may be employed for this purpose. Factors of importance in selecting a particular plasmid or viral vector include: the ease with which recipient cells that contain the vector may be recognized and selected from those recipient cells which do not contain the vector; the number of copies of the vector which are
25 desired in a particular host; and whether it is desirable to be able to "shuttle" the vector between host cells of different species.

Preferred prokaryotic vectors include plasmids such as those capable of replication in *E. coli* (such as, for example, pBR322, ColEI, pSC101, pACYC 184, π VX; "Molecular Cloning: A Laboratory Manual", 1989, *supra*). Bacillus plasmids

include pC194, pC221, pT127, and the like (Gryczan, In: The Molecular Biology of the Bacilli, Academic Press, NY, pp. 307-329, 1982). Suitable *Streptomyces* plasmids include p1J101 (Kendall *et al.*, *J. Bacteriol.* 169:4177-4183, 1987), and streptomyces bacteriophages such as ϕ C31 (Chater *et al.*, In: Sixth International Symposium on Actinomycetales Biology, Akademiai Kiado, Budapest, Hungary, pp. 45-54, 1986). *Pseudomonas* plasmids are reviewed by John *et al.* (*Rev. Infect. Dis.* 8:693-704, 1986), and Izaki (*Jpn. J. Bacteriol.* 33:729-742, 1978).

Preferred eukaryotic plasmids include, for example, BPV, vaccinia, SV40, 2-micron circle, and the like, or their derivatives. Such plasmids are well known in the art (Botstein *et al.*, *Miami Wntr. Symp.* 19:265-274, 1982; Broach, In: The Molecular Biology of the Yeast *Saccharomyces*: Life Cycle and Inheritance, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY, p. 445-470, 1981; Broach, *Cell* 28:203-204, 1982; Bollon *et al.*, *J. Clin. Hematol. Oncol.* 10:39-48, 1980; Maniatis, In: Cell Biology: A Comprehensive Treatise, Vol. 3, *Gene Sequence Expression*, Academic Press, NY, pp. 563-608, 1980).

Once the vector or nucleic acid molecule containing the construct(s) has been prepared for expression, the DNA construct(s) may be introduced into an appropriate host cell by any of a variety of suitable means, *i.e.*, transformation, transfection, conjugation, protoplast fusion, electroporation, particle gun technology, calcium phosphate-precipitation, direct microinjection, and the like. After the introduction of the vector, recipient cells are grown in a selective medium, which selects for the growth of vector-containing cells. Expression of the cloned gene(s) results in the production of a protease of the invention, or fragments thereof. This can take place in the transformed cells as such, or following the induction of these cells to differentiate (for example, by administration of bromodeoxyuracil to neuroblastoma cells or the like). A variety of incubation conditions can be used to form the peptide of the present invention. The most preferred conditions are those which mimic physiological conditions.

Antibodies, Hybridomas, Methods of Use and Kits for Detection of Proteases

The present invention relates to an antibody having binding affinity to a protease of the invention. The protease polypeptide may have the amino acid sequence selected from the group consisting of those set forth in SEQ ID NO:60,
5 SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85,
10 SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:109,
15 SEQ ID NO:110, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118, or a functional derivative thereof, or at least 9 contiguous amino acids thereof (preferably, at least 20, 30, 35, or 40 contiguous amino acids thereof).

The present invention also relates to an antibody having specific binding
20 affinity to a protease of the invention. Such an antibody may be isolated by comparing its binding affinity to a protease of the invention with its binding affinity to other polypeptides. Those which bind selectively to a protease of the invention would be chosen for use in methods requiring a distinction between a protease of the invention and other polypeptides. Such methods could include, but should not be
25 limited to, the analysis of altered protease expression in tissue containing other polypeptides.

The proteases of the present invention can be used in a variety of procedures and methods, such as for the generation of antibodies, for use in identifying pharmaceutical compositions, and for studying DNA/protein interaction.

The proteases of the present invention can be used to produce antibodies or hybridomas. One skilled in the art will recognize that if an antibody is desired, such a peptide could be generated as described herein and used as an immunogen. The antibodies of the present invention include monoclonal and polyclonal antibodies, as well fragments of these antibodies, and humanized forms. Humanized forms of the antibodies of the present invention may be generated using one of the procedures known in the art such as chimerization or CDR grafting.

The present invention also relates to a hybridoma which produces the above-described monoclonal antibody, or binding fragment thereof. A hybridoma is an immortalized cell line which is capable of secreting a specific monoclonal antibody.

In general, techniques for preparing monoclonal antibodies and hybridomas are well known in the art (Campbell, Monoclonal Antibody Technology: Laboratory Techniques in Biochemistry and Molecular Biology, Elsevier Science Publishers, Amsterdam, The Netherlands, 1984; St. Groth *et al.*, *J. Immunol. Methods* 35:1-21, 1980). Any animal (mouse, rabbit, and the like) which is known to produce antibodies can be immunized with the selected polypeptide. Methods for immunization are well known in the art. Such methods include subcutaneous or intraperitoneal injection of the polypeptide. One skilled in the art will recognize that the amount of polypeptide used for immunization will vary based on the animal which is immunized, the antigenicity of the polypeptide and the site of injection.

The polypeptide may be modified or administered in an adjuvant in order to increase the peptide antigenicity. Methods of increasing the antigenicity of a polypeptide are well known in the art. Such procedures include coupling the antigen with a heterologous protein (such as globulin or β -galactosidase) or through the inclusion of an adjuvant during immunization.

For monoclonal antibodies, spleen cells from the immunized animals are removed, fused with myeloma cells, such as SP2/0-Ag14 myeloma cells, and allowed to become monoclonal antibody producing hybridoma cells. Any one of a number of methods well known in the art can be used to identify the hybridoma cell

which produces an antibody with the desired characteristics. These include screening the hybridomas with an ELISA assay, western blot analysis, or radioimmunoassay (Lutz *et al.*, *Exp. Cell Res.* 175:109-124, 1988). Hybridomas secreting the desired antibodies are cloned and the class and subclass are determined
5 using procedures known in the art (Campbell, "Monoclonal Antibody Technology: Laboratory Techniques in Biochemistry and Molecular Biology", *supra*, 1984).

For polyclonal antibodies, antibody-containing antisera is isolated from the immunized animal and is screened for the presence of antibodies with the desired specificity using one of the above-described procedures. The above-described
10 antibodies may be detectably labeled. Antibodies can be detectably labeled through the use of radioisotopes, affinity labels (such as biotin, avidin, and the like), enzymatic labels (such as horseradish peroxidase, alkaline phosphatase, and the like) fluorescent labels (such as FITC or rhodamine, and the like), paramagnetic atoms, and the like. Procedures for accomplishing such labeling are well-known in the art,
15 for example, *see* Stemberger *et al.*, *J. Histochem. Cytochem.* 18:315, 1970; Bayer *et al.*, *Meth. Enzym.* 62:308, 1979; Engval *et al.*, *Immunol.* 109:129, 1972; Goding, *J. Immunol. Meth.* 13:215, 1976. The antibodies of the present invention may be indirectly labelled by the use of secondary labelled antibodies, such as labelled anti-rabbit antibodies. The labeled antibodies of the present invention can be used for *in*
20 *vitro*, *in vivo*, and *in situ* assays to identify cells or tissues which express a specific peptide.

The above-described antibodies may also be immobilized on a solid support. Examples of such solid supports include plastics such as polycarbonate, complex carbohydrates such as agarose and sepharose, acrylic resins such as polyacrylamide
25 and latex beads. Techniques for coupling antibodies to such solid supports are well known in the art (Weir *et al.*, "Handbook of Experimental Immunology" 4th Ed., Blackwell Scientific Publications, Oxford, England, Chapter 10, 1986; Jacoby *et al.*, *Meth. Enzym.* 34, Academic Press, N.Y., 1974). The immobilized antibodies of the

present invention can be used for *in vitro*, *in vivo*, and *in situ* assays as well as in immunochromotography.

Furthermore, one skilled in the art can readily adapt currently available procedures, as well as the techniques, methods and kits disclosed herein with regard
5 to antibodies, to generate peptides capable of binding to a specific peptide sequence in order to generate rationally designed anti-peptide peptides (Hurby *et al.*, "Application of Synthetic Peptides: Antisense Peptides", In Synthetic Peptides, A User's Guide, W.H. Freeman, NY, pp. 289-307, 1992; Kaspczak *et al.*, *Biochemistry* 28:9230-9238, 1989).

10 Anti-peptide peptides can be generated by replacing the basic amino acid residues found in the peptide sequences of the proteases of the invention with acidic residues, while maintaining hydrophobic and uncharged polar groups. For example, lysine, arginine, and/or histidine residues are replaced with aspartic acid or glutamic acid and glutamic acid residues are replaced by lysine, arginine or histidine.

15 The present invention also encompasses a method of detecting a protease polypeptide in a sample, comprising: (a) contacting the sample with an above-described antibody, under conditions such that immunocomplexes form, and (b) detecting the presence of said antibody bound to the polypeptide. In detail, the methods comprise incubating a test sample with one or more of the antibodies of the
20 present invention and assaying whether the antibody binds to the test sample. Altered levels of a protease of the invention in a sample as compared to normal levels may indicate disease.

Conditions for incubating an antibody with a test sample vary. Incubation conditions depend on the format employed in the assay, the detection methods
25 employed, and the type and nature of the antibody used in the assay. One skilled in the art will recognize that any one of the commonly available immunological assay formats (such as radioimmunoassays, enzyme-linked immunosorbent assays, diffusion-based Ouchterlony, or rocket immunofluorescent assays) can readily be adapted to employ the antibodies of the present invention. Examples of such assays

can be found in Chard ("An Introduction to Radioimmunoassay and Related Techniques") Elsevier Science Publishers, Amsterdam, The Netherlands, 1986), Bullock *et al.* ("Techniques in Immunocytochemistry," Academic Press, Orlando, FL Vol. 1, 1982; Vol. 2, 1983; Vol. 3, 1985), Tijssen ("Practice and Theory of Enzyme Immunoassays: Laboratory Techniques in Biochemistry and Molecular Biology," Elsevier Science Publishers, Amsterdam, The Netherlands, 1985).

The immunological assay test samples of the present invention include cells, protein or membrane extracts of cells, or biological fluids such as blood, serum, plasma, or urine. The test samples used in the above-described method will vary based on the assay format, nature of the detection method and the tissues, cells or extracts used as the sample to be assayed. Methods for preparing protein extracts or membrane extracts of cells are well known in the art and can readily be adapted in order to obtain a sample which is testable with the system utilized.

A kit contains all the necessary reagents to carry out the previously described methods of detection. The kit may comprise: (i) a first container means containing an above-described antibody, and (ii) second container means containing a conjugate comprising a binding partner of the antibody and a label. In another preferred embodiment, the kit further comprises one or more other containers comprising one or more of the following: wash reagents and reagents capable of detecting the presence of bound antibodies.

Examples of detection reagents include, but are not limited to, labeled secondary antibodies, or in the alternative, if the primary antibody is labeled, the chromophoric, enzymatic, or antibody binding reagents which are capable of reacting with the labeled antibody. The compartmentalized kit may be as described above for nucleic acid probe kits. One skilled in the art will readily recognize that the antibodies described in the present invention can readily be incorporated into one of the established kit formats which are well known in the art.

Isolation of Compounds Which Interact with Proteases

The present invention also relates to a method of detecting a compound capable of binding to a protease of the invention comprising incubating the compound with a protease of the invention and detecting the presence of the compound bound to the protease. The compound may be present within a complex mixture, for example, serum, body fluid, or cell extracts.

The present invention also relates to a method of detecting an agonist or antagonist of protease activity or protease binding partner activity comprising incubating cells that produce a protease of the invention in the presence of a compound and detecting changes in the level of protease activity or protease binding partner activity. The compounds thus identified would produce a change in activity indicative of the presence of the compound. The compound may be present within a complex mixture, for example, serum, body fluid, or cell extracts. Once the compound is identified it can be isolated using techniques well known in the art.

The present invention also encompasses a method of modulating protease associated activity in a mammal comprising administering to said mammal an agonist or antagonist to a protease of the invention in an amount sufficient to effect said modulation. A method of treating diseases in a mammal with an agonist or antagonist of the activity of one of the proteases of the invention comprising administering the agonist or antagonist to a mammal in an amount sufficient to agonize or antagonize protease-associated functions is also encompassed in the present application.

In an effort to discover novel treatments for diseases, biomedical researchers and chemists have designed, synthesized, and tested molecules that inhibit the function of proteases. Some small organic molecules form a class of compounds that modulate the function of protein proteases.

Examples of molecules that have been reported to inhibit the function of protein proteases include, but are not limited to, phenylmethylsulfonyl fluoride (PMSF), diisopropylfluorophosphate (DFP) (chapter 3, Barrett *et al.*, Handbook of

Proteolytic Enzymes, 1998, Academic Press, San Diego), 3,4-dichloroisocoumarin (DCI) (*Id.*, chapter 16), serpins (*Id.*, chapter 37), E-64 (*trans*-epoxysuccinyl L-leucylamido-(4-guanidino) butane) (*Id.*, chapter 188), peptidyl-diazomethanes, peptidyl-*O*-acyl-hydroxamates, epoxysuccinyl-peptides (*Id.*, chapter 210), DAN, 5 EPNP (1,2-epoxy-3(p-nitrophenoxy)propane) (*Id.*, chapter 298), thiorphan (dl-3-Mercapto-2-benzylpropanoyl-glycine) (*Id.*, chapter 362), CGS 26303, PD 069185 (*Id.*, chapter 363), and COT989-00 (N-4-hydroxy-N1-[1-(s)-(4-aminosulfonyl)phenylethyl-aminocarboxyl-2-cyclohexylethyl)-2R-[4-methyl)phenylpropyl]succinamide) (*Id.*, chapter 401). Other protease inhibitors 10 include, but are not limited to, aprotinin, amastatin, antipain, calcineurin autoinhibitory fragment, and histatin 5 (*Id.*). Preferably, these inhibitors will have molecular weights from 100 to 200 daltons, from 200 to 300 daltons, from 300 to 400 daltons, from 400 to 600 daltons, from 600 to 1000 daltons, from 1000 to 2000 daltons, from 2000 to 4000 daltons, and from 4000 to 8000 daltons.

15 Compounds that can traverse cell membranes and are resistant to acid hydrolysis are potentially advantageous as therapeutics as they can become highly bioavailable after being administered orally to patients. However, many of these protease inhibitors only weakly inhibit the function of proteases. In addition, many inhibit a variety of proteases and will therefore cause multiple side-effects as 20 therapeutics for diseases.

Transgenic Animals.

A variety of methods are available for the production of transgenic animals associated with this invention. DNA can be injected into the pronucleus of a 25 fertilized egg before fusion of the male and female pronuclei, or injected into the nucleus of an embryonic cell (*e.g.*, the nucleus of a two-cell embryo) following the initiation of cell division (Brinster *et al.*, *Proc. Nat. Acad. Sci. USA* 82:4438-4442, 1985). Embryos can be infected with viruses, especially retroviruses, modified to carry inorganic-ion receptor nucleotide sequences of the invention.

Pluripotent stem cells derived from the inner cell mass of the embryo and stabilized in culture can be manipulated in culture to incorporate nucleotide sequences of the invention. A transgenic animal can be produced from such cells through implantation into a blastocyst that is implanted into a foster mother and
5 allowed to come to term. Animals suitable for transgenic experiments can be obtained from standard commercial sources such as Charles River (Wilmington, MA), Taconic (Germantown, NY), Harlan Sprague Dawley (Indianapolis, IN), etc.

The procedures for manipulation of the rodent embryo and for microinjection of DNA into the pronucleus of the zygote are well known to those of ordinary skill
10 in the art (Hogan *et al.*, *supra*). Microinjection procedures for fish, amphibian eggs and birds are detailed in Houdebine and Chourrout (*Experientia* 47:897-905, 1991). Other procedures for introduction of DNA into tissues of animals are described in U.S. Patent No. 4,945,050 (Sanford *et al.*, July 30, 1990).

By way of example only, to prepare a transgenic mouse, female mice are
15 induced to superovulate. Females are placed with males, and the mated females are sacrificed by CO₂ asphyxiation or cervical dislocation and embryos are recovered from excised oviducts. Surrounding cumulus cells are removed. Pronuclear embryos are then washed and stored until the time of injection. Randomly cycling adult female mice are paired with vasectomized males. Recipient females are mated
20 at the same time as donor females. Embryos then are transferred surgically. The procedure for generating transgenic rats is similar to that of mice (Hammer *et al.*, *Cell* 63:1099-1112, 1990).

Methods for the culturing of embryonic stem (ES) cells and the subsequent production of transgenic animals by the introduction of DNA into ES cells using
25 methods such as electroporation, calcium phosphate/DNA precipitation and direct injection also are well known to those of ordinary skill in the art (Teratocarcinomas and Embryonic Stem Cells, A Practical Approach, E.J. Robertson, ed., IRL Press, 1987).

In cases involving random gene integration, a clone containing the sequence(s) of the invention is co-transfected with a gene encoding resistance. Alternatively, the gene encoding neomycin resistance is physically linked to the sequence(s) of the invention. Transfection and isolation of desired clones are carried
5 out by any one of several methods well known to those of ordinary skill in the art (E.J. Robertson, *supra*).

DNA molecules introduced into ES cells can also be integrated into the chromosome through the process of homologous recombination (Capecchi, *Science* 244:1288-1292, 1989). Methods for positive selection of the recombination event
10 (*i.e.*, neo resistance) and dual positive-negative selection (*i.e.*, neo resistance and gancyclovir resistance) and the subsequent identification of the desired clones by PCR have been described by Capecchi, *supra* and Joyner *et al.* (*Nature* 338:153-156, 1989), the teachings of which are incorporated herein in their entirety including any drawings. The final phase of the procedure is to inject targeted ES cells into
15 blastocysts and to transfer the blastocysts into pseudopregnant females. The resulting chimeric animals are bred and the offspring are analyzed by Southern blotting to identify individuals that carry the transgene. Procedures for the production of non-rodent mammals and other animals have been discussed by others (Houdebine and Chourrout, *supra*; Pursel *et al.*, *Science* 244:1281-1288, 1989; and
20 Simms *et al.*, *Bio/Technology* 6:179-183, 1988).

Thus, the invention provides transgenic, nonhuman mammals containing a transgene encoding a protease of the invention or a gene affecting the expression of the protease. Such transgenic nonhuman mammals are particularly useful as an *in vivo* test system for studying the effects of introduction of a protease, or regulating
25 the expression of a protease (*i.e.*, through the introduction of additional genes, antisense nucleic acids, or ribozymes).

A "transgenic animal" is an animal having cells that contain DNA which has been artificially inserted into a cell, which DNA becomes part of the genome of the animal which develops from that cell. Preferred transgenic animals are primates,

mice, rats, cows, pigs, horses, goats, sheep, dogs and cats. The transgenic DNA may encode human proteases. Native expression in an animal may be reduced by providing an amount of antisense RNA or DNA effective to reduce expression of the receptor.

5

Gene Therapy

Proteases or their genetic sequences will also be useful in gene therapy (reviewed in Miller, *Nature* 357:455-460, 1992). Miller states that advances have resulted in practical approaches to human gene therapy that have demonstrated
10 positive initial results. The basic science of gene therapy is described in Mulligan (*Science* 260:926-931, 1993).

In one preferred embodiment, an expression vector containing a protease coding sequence is inserted into cells, the cells are grown *in vitro* and then infused in large numbers into patients. In another preferred embodiment, a DNA segment
15 containing a promoter of choice (for example a strong promoter) is transferred into cells containing an endogenous gene encoding proteases of the invention in such a manner that the promoter segment enhances expression of the endogenous protease gene (for example, the promoter segment is transferred to the cell such that it becomes directly linked to the endogenous protease gene).

20 The gene therapy may involve the use of an adenovirus containing protease cDNA targeted to a tumor, systemic protease increase by implantation of engineered cells, injection with protease-encoding virus, or injection of naked protease DNA into appropriate tissues.

Target cell populations may be modified by introducing altered forms of one
25 or more components of the protein complexes in order to modulate the activity of such complexes. For example, by reducing or inhibiting a complex component activity within target cells, an abnormal signal transduction event(s) leading to a condition may be decreased, inhibited, or reversed. Deletion or missense mutants of a component, that retain the ability to interact with other components of the protein

complexes but cannot function in signal transduction, may be used to inhibit an abnormal, deleterious signal transduction event.

Expression vectors derived from viruses such as retroviruses, vaccinia virus, adenovirus, adeno-associated virus, herpes viruses, several RNA viruses, or bovine papilloma virus, may be used for delivery of nucleotide sequences (*e.g.*, cDNA) encoding recombinant protease of the invention protein into the targeted cell population (*e.g.*, tumor cells). Methods which are well known to those skilled in the art can be used to construct recombinant viral vectors containing coding sequences (Maniatis *et al.*, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, N.Y., 1989; Ausubel *et al.*, Current Protocols in Molecular Biology, Greene Publishing Associates and Wiley Interscience, N.Y., 1989). Alternatively, recombinant nucleic acid molecules encoding protein sequences can be used as naked DNA or in a reconstituted system *e.g.*, liposomes or other lipid systems for delivery to target cells (*e.g.*, Felgner *et al.*, *Nature* 337:387-8, 1989). Several other methods for the direct transfer of plasmid DNA into cells exist for use in human gene therapy and involve targeting the DNA to receptors on cells by complexing the plasmid DNA to proteins (Miller, *supra*).

In its simplest form, gene transfer can be performed by simply injecting minute amounts of DNA into the nucleus of a cell, through a process of microinjection (Capecchi, *Cell* 22:479-88, 1980). Once recombinant genes are introduced into a cell, they can be recognized by the cell's normal mechanisms for transcription and translation, and a gene product will be expressed. Other methods have also been attempted for introducing DNA into larger numbers of cells. These methods include: transfection, wherein DNA is precipitated with calcium phosphate and taken into cells by pinocytosis (Chen *et al.*, *Mol. Cell Biol.* 7:2745-52, 1987); electroporation, wherein cells are exposed to large voltage pulses to introduce holes into the membrane (Chu *et al.*, *Nucleic Acids Res.* 15:1311-26, 1987); lipofection/liposome fusion, wherein DNA is packaged into lipophilic vesicles which fuse with a target cell (Felgner *et al.*, *Proc. Natl. Acad. Sci. USA.* 84:7413-

7417, 1987); and particle bombardment using DNA bound to small projectiles (Yang *et al.*, *Proc. Natl. Acad. Sci.* 87:9568-9572, 1990). Another method for introducing DNA into cells is to couple the DNA to chemically modified proteins.

5 It has also been shown that adenovirus proteins are capable of destabilizing endosomes and enhancing the uptake of DNA into cells. The admixture of adenovirus to solutions containing DNA complexes, or the binding of DNA to polylysine covalently attached to adenovirus using ~~protein~~ crosslinking agents substantially improves the uptake and expression of the recombinant gene (Curiel *et al.*, *Am. J. Respir. Cell. Mol. Biol.*, 6:247-52, 1992).

0 As used herein "gene transfer" means the process of introducing a foreign nucleic acid molecule into a cell. Gene transfer is commonly performed to enable the expression of a particular product encoded by the gene. The product may include a protein, polypeptide, anti-sense DNA or RNA, or enzymatically active RNA. Gene transfer can be performed in cultured cells or by direct administration
5 into animals. Generally gene transfer involves the process of nucleic acid contact with a target cell by non-specific or receptor mediated interactions, uptake of nucleic acid into the cell through the membrane or by endocytosis, and release of nucleic acid into the cyto-plasm from the plasma membrane or endosome. Expression may require, in addition, movement of the nucleic acid into the nucleus of the cell and
10 binding to appropriate nuclear factors for transcription.

As used herein "gene therapy" is a form of gene transfer and is included within the definition of gene transfer as used herein and specifically refers to gene transfer to express a therapeutic product from a cell *in vivo* or *in vitro*. Gene transfer can be performed *ex vivo* on cells which are then transplanted into a patient, or can
15 be performed by direct administration of the nucleic acid or nucleic acid-protein complex into the patient.

In another preferred embodiment, a vector having nucleic acid sequences encoding a protease polypeptide is provided in which the nucleic acid sequence is expressed only in specific tissue. Methods of achieving tissue-specific gene

expression are set forth in International Publication No. WO 93/09236, filed November 3, 1992 and published May 13, 1993.

In all of the preceding vectors set forth above, a further aspect of the invention is that the nucleic acid sequence contained in the vector may include
5 additions, deletions or modifications to some or all of the sequence of the nucleic acid, as defined above.

Expression, including over-expression, of a protease polypeptide of the invention can be inhibited by administration of an antisense molecule that binds to and inhibits expression of the mRNA encoding the polypeptide. Alternatively, expression
10 can be inhibited in an analogous manner using a ribozyme that cleaves the mRNA. General methods of using antisense and ribozyme technology to control gene expression, or of gene therapy methods for expression of an exogenous gene in this manner are well known in the art. Each of these methods utilizes a system, such as a vector, encoding either an antisense or ribozyme transcript of a protease polypeptide of
15 the invention.

The term "*ribozyme*" refers to an RNA structure of one or more RNAs having catalytic properties. Ribozymes generally exhibit endonuclease, ligase or polymerase activity. Ribozymes are structural RNA molecules which mediate a number of RNA self-cleavage reactions. Various types of trans-acting ribozymes,
20 including "hammerhead" and "hairpin" types, which have different secondary structures, have been identified. A variety of ribozymes have been characterized. See, for example, U.S. Pat. Nos. 5,246,921, 5,225,347, 5,225,337 and 5,149,796. Mixed ribozymes comprising deoxyribo and ribooligonucleotides with catalytic activity have been described. Perreault, *et al.*, *Nature*, 344:565-567 (1990).

25 As used herein, "antisense" refers of nucleic acid molecules or their derivatives which specifically hybridize, *e.g.*, bind, under cellular conditions, with the genomic DNA and/or cellular mRNA encoding a protease polypeptide of the invention, so as to inhibit expression of that protein, for example, by inhibiting transcription and/or translation. The binding may be by conventional base pair

complementarity, or, for example, in the case of binding to DNA duplexes, through specific interactions in the major groove of the double helix.

In one aspect, the antisense construct is an nucleic acid which is generated *ex vivo* and that, when introduced into the cell, can inhibit gene expression by, without
5 limitation, hybridizing with the mRNA and/or genomic sequences of a protease polynucleotide of the invention.

Antisense approaches can involve the design of oligonucleotides (either DNA or RNA) that are complementary to protease polypeptide mRNA and are based on the protease polynucleotides of the invention, including SEQ ID NO:1,
10 SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID
15 NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:44, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:58, and SEQ ID NO:59. The antisense oligonucleotides will bind to the protease polypeptide mRNA transcripts and prevent translation.

Although absolute complementarity is preferred, it is not required. A sequence "complementary" to a portion of an RNA, as referred to herein, means a
25 sequence having sufficient complementarity to be able to hybridize with the RNA, forming a stable duplex; in the case of double-stranded antisense nucleic acids, a single strand of the duplex DNA may thus be tested, or triplex formation may be assayed. The ability to hybridize will depend on both the degree of complementarity and the length of the antisense nucleic acid. Generally, the longer the hybridizing

nucleic acid, the more base mismatches with an RNA it may contain and still form a stable duplex (or triplex, as the case may be). One skilled in the art can ascertain a tolerable degree of mismatch by use of standard procedures to determine the melting point of the hybridized complex.

- 5 In general, oligonucleotides that are complementary to the 5' end of the message, *e.g.*, the 5' untranslated sequence up to and including the AUG initiation codon, should work most efficiently at inhibiting translation. However, sequences complementary to the 3' untranslated sequences of mRNAs have been shown to be effective at inhibiting translation of mRNAs as well. (Wagner, R. (1994) Nature
- 10 372:333). Antisense oligonucleotides complementary to mRNA coding regions are less efficient inhibitors of translation but could be used in accordance with the invention. Whether designed to hybridize to the 5', 3' or coding region of the protease polypeptide mRNA, antisense nucleic acids should be at least six nucleotides in length, and are preferably less than about 100 and more preferably
- 15 less than about 50 or 30 nucleotides in length. Typically they should be between 10 and 25 nucleotides in length. Such principles will inform the practitioner in selecting the appropriate oligonucleotides. In preferred embodiments, the antisense sequence is selected from an oligonucleotide sequence that comprises, consists of, or consists essentially of about 10-30, and more preferably 15-25, contiguous
- 20 nucleotide bases of a nucleic acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID
- 25 NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:44, SEQ ID NO:45, SEQ ID

NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:58, and SEQ ID NO:59 or domains thereof.

In another preferred embodiment, the invention includes an isolated,
5 enriched or purified nucleic acid molecule comprising, consisting of or consisting essentially of about 10-30, and more preferably 15-25 contiguous nucleotide bases of a nucleic acid sequence that encodes a polypeptide that selected from the group consisting of SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118.

Using the sequences of the present invention, antisense oligonucleotides can be designed. Such antisense oligonucleotides would be administered to cells expressing the target protease and the levels of the target RNA or protein with that of an internal control RNA or protein would be compared. Results obtained using
25 the antisense oligonucleotide would also be compared with those obtained using a suitable control oligonucleotide. A preferred control oligonucleotide is an oligonucleotide of approximately the same length as the test oligonucleotide. Those antisense oligonucleotides resulting in a reduction in levels of target RNA or protein would be selected.

The oligonucleotides can be DNA or RNA or chimeric mixtures or derivatives or modified versions thereof, single-stranded or double-stranded. The oligonucleotide can be modified at the base moiety, sugar moiety, or phosphate backbone, for example, to improve stability of the molecule, hybridization, etc. The oligonucleotide may include other appended groups such as peptides (*e.g.*, for targeting host cell receptors *in vivo*), or agents facilitating transport across the cell membrane (*see, e.g.*, Letsinger *et al.* (1989) *Proc. Natl. Acad. Sci. U.S.A.* 86:6553-6556; Lemaitre *et al.* (1987) *Proc. Natl. Acad. Sci. USA* 84:648-652; PCT Publication No. WO 88/09810, published Dec. 15, 1988) or the blood-brain barrier (*see, e.g.*, PCT Publication No. WO 89/10134, published Apr. 25, 1988), hybridization-triggered cleavage agents. (*See, e.g.*, Krol *et al.* (1988) *BioTechniques* 6:958-976) or intercalating agents. (*See, e.g.*, Zon (1988) *Pharm. Res.* 5:539-549). To this end, the oligonucleotide may be conjugated to another molecule, *e.g.*, a peptide, hybridization triggered cross-linking agent, transport agent, hybridization-triggered cleavage agent, etc.

The antisense oligonucleotide may comprise at least one modified base moiety which is selected from moieties such as 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acetylcytosine, and 5-(carboxyhydroxyethyl) uracil. The antisense oligonucleotide may also comprise at least one modified sugar moiety selected from the group including but not limited to arabinose, 2-fluoroarabinose, xylulose, and hexose.

In yet another embodiment, the antisense oligonucleotide comprises at least one modified phosphate backbone selected from the group consisting of a phosphorothioate, a phosphorodithioate, a phosphoramidothioate, a phosphoramidate, a phosphordiamidate, a methylphosphonate, an alkyl phosphotriester, and a formacetal or analog thereof. (*see also* U.S. Pat. Nos. 5,176,996; 5,264,564; and 5,256,775)

In yet a further embodiment, the antisense oligonucleotide is an α -anomeric oligonucleotide. An α -anomeric oligonucleotide forms specific double-stranded

hybrids with complementary RNA in which, contrary to the usual β -units, the strands run parallel to each other (Gautier *et al.* (1987) *Nucl. Acids Res.* 15:6625-6641). The oligonucleotide is a 2'-O-methylribonucleotide (Inoue *et al.* (1987) *Nucl. Acids Res.* 15:6131-6148), or a chimeric RNA-DNA analogue (Inoue *et al.* (1987) FEBS Lett. 215:327-330).

Also suitable are peptidyl nucleic acids, which are polypeptides such as polyserine, polythreonine, etc. including copolymers containing various amino acids, which are substituted at side-chain positions with nucleic acids (T,A,G,C,U). Chains of such polymers are able to hybridize through complementary bases in the same manner as natural DNA/RNA.. Alternatively, an antisense construct of the present invention can be delivered, for example, as an expression plasmid or vector that, when transcribed in the cell, produces RNA complementary to at least a unique portion of the cellular mRNA which encodes a protease polypeptide of the invention.

While antisense nucleotides complementary to the protease polypeptide coding region sequence can be used, those complementary to the transcribed untranslated region are most preferred.

In another preferred embodiment, a method of gene replacement is set forth. "Gene replacement" as used herein means supplying a nucleic acid sequence which is capable of being expressed *in vivo* in an animal and thereby providing or augmenting the function of an endogenous gene which is missing or defective in the animal.

Pharmaceutical Formulations And Routes Of Administration

The compounds described herein, including protease polypeptides of the invention, antisense molecules, ribozymes, and any other compound that modulates the activity of a protease polypeptide of the invention, can be administered to a human patient *per se*, or in pharmaceutical compositions where it is mixed with other active ingredients, as in combination therapy, or suitable carriers or

excipient(s). Techniques for formulation and administration of the compounds of the instant application may be found in "Remington's Pharmaceutical Sciences," Mack Publishing Co., Easton, PA, latest edition.

A. Routes Of Administration

5 Suitable routes of administration may, for example, include oral, rectal, transmucosal, or intestinal administration; parenteral delivery, including intramuscular, subcutaneous, intravenous, intramedullary injections, as well as intrathecal, direct intraventricular, intraperitoneal, intranasal, or intraocular injections.

10 Alternately, one may administer the compound in a local rather than systemic manner, for example, via injection of the compound directly into a solid tumor, often in a depot or sustained release formulation.

 Furthermore, one may administer the drug in a targeted drug delivery system, for example, in a liposome coated with tumor-specific antibody. The liposomes will
15 be targeted to and taken up selectively by the tumor.

B. Composition/Formulation

 The pharmaceutical compositions of the present invention may be manufactured in a manner that is itself known, *e.g.*, by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying,
20 encapsulating, entrapping or lyophilizing processes.

 Pharmaceutical compositions for use in accordance with the present invention thus may be formulated in conventional manner using one or more physiologically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used
25 pharmaceutically. Proper formulation is dependent upon the route of administration chosen.

 For injection, the agents of the invention may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks's solution, Ringer's solution, or physiological saline buffer. For transmucosal administration,

penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

For oral administration, the compounds can be formulated readily by combining the active compounds with pharmaceutically acceptable carriers well known in the art. Such carriers enable the compounds of the invention to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions and the like, for oral ingestion by a patient to be treated. Suitable carriers include excipients such as, fillers such as sugars, including lactose, sucrose, mannitol, or sorbitol; cellulose preparations such as, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, gum tragacanth, methyl cellulose, hydroxypropylmethyl cellulose, sodium carboxymethylcellulose, and/or polyvinylpyrrolidone (PVP). If desired, disintegrating agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof such as sodium alginate.

Dragee cores are provided with suitable coatings. For this purpose, concentrated sugar solutions may be used, which may optionally contain gum arabic, talc, polyvinyl pyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for identification or to characterize different combinations of active compound doses.

Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The push-fit capsules can contain the active ingredients in admixture with filler such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In addition, stabilizers may be added. All formulations for oral administration should be in dosages suitable for such administration.

For buccal administration, the compositions may take the form of tablets or lozenges formulated in conventional manner.

For administration by inhalation, the compounds for use according to the present invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebuliser, with the use of a suitable propellant, *e.g.*, dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of *e.g.* gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

The compounds may be formulated for parenteral administration by injection, *e.g.*, by bolus injection or continuous infusion. Formulations for injection may be presented in unit dosage form, *e.g.*, in ampoules or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents.

Pharmaceutical formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble form. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions.

Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, *e.g.*, sterile pyrogen-free water, before use.

The compounds may also be formulated in rectal compositions such as suppositories or retention enemas, *e.g.*, containing conventional suppository bases such as cocoa butter or other glycerides.

In addition to the formulations described previously, the compounds may also be formulated as a depot preparation. Such long acting formulations may be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds may be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

A pharmaceutical carrier for the hydrophobic compounds of the invention is a cosolvent system comprising benzyl alcohol, a nonpolar surfactant, a water-miscible organic polymer, and an aqueous phase. The cosolvent system may be the VPD co-solvent system. VPD is a solution of 3% w/v benzyl alcohol, 8% w/v of the nonpolar surfactant polysorbate 80, and 65% w/v polyethylene glycol 300, made up to volume in absolute ethanol. The VPD co-solvent system (VPD:D5W) consists of VPD diluted 1:1 with a 5% dextrose in water solution. This co-solvent system dissolves hydrophobic compounds well, and itself produces low toxicity upon systemic administration. Naturally, the proportions of a co-solvent system may be varied considerably without destroying its solubility and toxicity characteristics. Furthermore, the identity of the co-solvent components may be varied: for example, other low-toxicity nonpolar surfactants may be used instead of polysorbate 80; the fraction size of polyethylene glycol may be varied; other biocompatible polymers may replace polyethylene glycol, *e.g.* polyvinyl pyrrolidone; and other sugars or polysaccharides may substitute for dextrose.

Alternatively, other delivery systems for hydrophobic pharmaceutical compounds may be employed. Liposomes and emulsions are well known examples of delivery vehicles or carriers for hydrophobic drugs. Certain organic solvents such as dimethylsulfoxide also may be employed, although usually at the cost of greater

toxicity. Additionally, the compounds may be delivered using a sustained-release system, such as semipermeable matrices of solid hydrophobic polymers containing the therapeutic agent. Various sustained-release materials have been established and are well known by those skilled in the art. Sustained-release capsules may, 5 depending on their chemical nature, release the compounds for a few weeks up to over 100 days. Depending on the chemical nature and the biological stability of the therapeutic reagent, additional strategies for protein stabilization may be employed.

The pharmaceutical compositions also may comprise suitable solid or gel phase carriers or excipients. Examples of such carriers or excipients include but are 0 not limited to calcium carbonate, calcium phosphate, various sugars, starches, cellulose derivatives, gelatin, and polymers such as polyethylene glycols.

Many of the protease modulating compounds of the invention may be provided as salts with pharmaceutically compatible counterions. Pharmaceutically compatible salts may be formed with many acids, including but not limited to 5 hydrochloric, sulfuric, acetic, lactic, tartaric, malic, succinic, etc. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free base forms.

C. Effective Dosage

Pharmaceutical compositions suitable for use in the present invention include 10 compositions where the active ingredients are contained in an amount effective to achieve its intended purpose. More specifically, a therapeutically effective amount means an amount of compound effective to prevent, alleviate or ameliorate symptoms of disease or prolong the survival of the subject being treated. Determination of a therapeutically effective amount is well within the capability of 15 those skilled in the art, especially in light of the detailed disclosure provided herein.

For any compound used in the methods of the invention, the therapeutically effective dose can be estimated initially from cell culture assays. For example, a dose can be formulated in animal models to achieve a circulating concentration range that includes the IC_{50} as determined in cell culture (*i.e.*, the concentration of

the test compound which achieves a half-maximal inhibition of the protease activity). Such information can be used to more accurately determine useful doses in humans.

Toxicity and therapeutic efficacy of the compounds described herein can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, *e.g.*, for determining the LD₅₀ (the dose lethal to 50% of the population) and the ED₅₀ (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio between LD₅₀ and ED₅₀. Compounds which exhibit high therapeutic indices are preferred. The data obtained from these cell culture assays and animal studies can be used in formulating a range of dosage for use in human. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED₅₀ with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. The exact formulation, route of administration and dosage can be chosen by the individual physician in view of the patient's condition. (See *e.g.*, Fingl *et al.*, 1975, in The Pharmacological Basis of Therapeutics, Ch. 1 p.1).

Dosage amount and interval may be adjusted individually to provide plasma levels of the active moiety which are sufficient to maintain the protease modulating effects, or minimal effective concentration (MEC). The MEC will vary for each compound but can be estimated from *in vitro* data; *e.g.*, the concentration necessary to achieve 50-90% inhibition of the protease using the assays described herein.

Dosages necessary to achieve the MEC will depend on individual characteristics and route of administration. However, HPLC assays or bioassays can be used to determine plasma concentrations.

Dosage intervals can also be determined using MEC value. Compounds should be administered using a regimen which maintains plasma levels above the MEC for 10-90% of the time, preferably between 30-90% and most preferably between 50-90%.

In cases of local administration or selective uptake, the effective local concentration of the drug may not be related to plasma concentration.

The amount of composition administered will, of course, be dependent on the subject being treated, on the subject's weight, the severity of the affliction, the
5 manner of administration and the judgment of the prescribing physician.

D. Packaging

The compositions may, if desired, be presented in a pack or dispenser device which may contain one or more unit dosage forms containing the active ingredient. The pack may for example comprise metal or plastic foil, such as a blister pack. The
10 pack or dispenser device may be accompanied by instructions for administration. The pack or dispenser may also be accompanied with a notice associated with the container in form prescribed by a governmental agency regulating the manufacture, use, or sale of pharmaceuticals, which notice is reflective of approval by the agency of the form of the polynucleotide for human or veterinary administration. Such
15 notice, for example, may be the labeling approved by the U.S. Food and Drug Administration for prescription drugs, or the approved product insert. Compositions comprising a compound of the invention formulated in a compatible pharmaceutical carrier may also be prepared, placed in an appropriate container, and labeled for treatment of an indicated condition. Suitable conditions indicated on the label may
20 include treatment of a tumor, inhibition of angiogenesis, treatment of fibrosis, diabetes, and the like.

Functional Derivatives

Also provided herein are functional derivatives of a polypeptide or nucleic
25 acid of the invention. By "functional derivative" is meant a "chemical derivative," "fragment," or "variant," of the polypeptide or nucleic acid of the invention, which terms are defined below. A functional derivative retains at least a portion of the function of the protein, for example reactivity with an antibody specific for the protein, enzymatic activity or binding activity mediated through noncatalytic

domains, which permits its utility in accordance with the present invention. It is well known in the art that due to the degeneracy of the genetic code numerous different nucleic acid sequences can code for the same amino acid sequence. Equally, it is also well known in the art that conservative changes in amino acid can
5 be made to arrive at a protein or polypeptide that retains the functionality of the original. In both cases, all permutations are intended to be covered by this disclosure.

Included within the scope of this invention are the functional equivalents of the herein-described isolated nucleic acid molecules. The degeneracy of the genetic
10 code permits substitution of certain codons by other codons that specify the same amino acid and hence would give rise to the same protein. The nucleic acid sequence can vary substantially since, with the exception of methionine and tryptophan, the known amino acids can be coded for by more than one codon. Thus, portions or all of the genes of the invention could be synthesized to give a nucleic
15 acid sequence significantly different from one selected from the group consisting of those set forth in SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID
20 NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:44, SEQ ID
25 NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:58, and SEQ ID NO:59. The encoded amino acid sequence thereof would, however, be preserved.

In addition, the nucleic acid sequence may comprise a nucleotide sequence which results from the addition, deletion or substitution of at least one nucleotide to the 5'-end and/or the 3'-end of the nucleic acid formula selected from the group consisting of those set forth in SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:44, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:58, and SEQ ID NO:59, or a derivative thereof. Any nucleotide or polynucleotide may be used in this regard, provided that its addition, deletion or substitution does not alter the amino acid sequence selected from the group consisting of those set forth in SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113,

SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118 which is encoded by the nucleotide sequence. For example, the present invention is intended to include any nucleic acid sequence resulting from the addition of ATG as an initiation codon at the 5'-end of the inventive nucleic acid sequence or its derivative; or from the addition of TTA, TAG or TGA as a termination codon at the 3'-end of the inventive nucleotide sequence or its derivative. Moreover, the nucleic acid molecule of the present invention may, as necessary, have restriction endonuclease recognition sites added to its 5'-end and/or 3'-end.

Such functional alterations of a given nucleic acid sequence afford an opportunity to promote secretion and/or processing of heterologous proteins encoded by foreign nucleic acid sequences fused thereto. All variations of the nucleotide sequence of the protease genes of the invention and fragments thereof permitted by the genetic code are, therefore, included in this invention.

Further, it is possible to delete codons or to substitute one or more codons with codons other than degenerate codons to produce a structurally modified polypeptide, but one which has substantially the same utility or activity as the polypeptide produced by the unmodified nucleic acid molecule. As recognized in the art, the two polypeptides are functionally equivalent, as are the two nucleic acid molecules that give rise to their production, even though the differences between the nucleic acid molecules are not related to the degeneracy of the genetic code.

A "chemical derivative" of the complex contains additional chemical moieties not normally a part of the protein. Covalent modifications of the protein or peptides are included within the scope of this invention. Such modifications may be introduced into the molecule by reacting targeted amino acid residues of the peptide with an organic derivatizing agent that is capable of reacting with selected side chains or terminal residues, as described below.

Cysteinyl residues most commonly are reacted with α -haloacetates (and corresponding amines), such as chloroacetic acid or chloroacetamide, to give carboxymethyl or carboxyamidomethyl derivatives. Cysteinyl residues also are derivatized by reaction with bromotrifluoroacetone, chloroacetyl phosphate, N-alkylmaleimides, 3-nitro-2-pyridyl disulfide, methyl 2-pyridyl disulfide, p-chloromercuribenzoate, 2-chloromercuri-4-nitrophenol, or chloro-7-nitrobenzo-2-oxa-1,3-diazole.

Histidyl residues are derivatized by reaction with diethylprocarbonate at pH 5.5-7.0 because this agent is relatively specific for the histidyl side chain. Para-bromophenacyl bromide also is useful; the reaction is preferably performed in 0.1 M sodium cacodylate at pH 6.0.

Lysinyl and amino terminal residues are reacted with succinic or other carboxylic acid anhydrides. Derivatization with these agents has the effect of reversing the charge of the lysinyl residues. Other suitable reagents for derivatizing primary amine containing residues include imidoesters such as methyl picolinimide; pyridoxal phosphate; pyridoxal; chloroborohydride; trinitrobenzenesulfonic acid; O-methylisourea; 2,4 pentanedione; and transaminase-catalyzed reaction with glyoxylate.

Arginyl residues are modified by reaction with one or several conventional reagents, among them phenylglyoxal, 2,3-butanedione, 1,2-cyclohexanedione, and ninhydrin. Derivatization of arginine residues requires that the reaction be performed in alkaline conditions because of the high pKa of the guanidine functional group. Furthermore, these reagents may react with the groups of lysine as well as the arginine α -amino group.

Tyrosyl residues are well-known targets of modification for introduction of spectral labels by reaction with aromatic diazonium compounds or tetranitromethane. Most commonly, N-acetylimidazol and tetranitromethane are used to form O-acetyl tyrosyl species and 3-nitro derivatives, respectively.

Carboxyl side groups (aspartyl or glutamyl) are selectively modified by reaction with carbodiimide ($R'-N-C-N-R'$) such as 1-cyclohexyl-3-(2-morpholinyl(4-ethyl) carbodiimide or 1-ethyl-3-(4-azonia-4,4-dimethylpentyl) carbodiimide. Furthermore, aspartyl and glutamyl residues are converted to asparaginy and glutaminy residues by reaction with ammonium ions.

Glutaminy and asparaginy residues are frequently deamidated to the corresponding glutamyl and aspartyl residues. Alternatively, these residues are deamidated under mildly acidic conditions. Either form of these residues falls within the scope of this invention.

Derivatization with bifunctional agents is useful, for example, for cross-linking the component peptides of the protein to each other or to other proteins in a complex to a water-insoluble support matrix or to other macromolecular carriers. Commonly used cross-linking agents include, for example, 1,1-bis(diazoacetyl)-2-phenylethane, glutaraldehyde, N-hydroxysuccinimide esters, for example, esters with 4-azidosalicylic acid, homobifunctional imidoesters, including disuccinimidyl esters such as 3,3'-dithiobis(succinimidylpropionate), and bifunctional maleimides such as bis-N-maleimido-1,8-octane. Derivatizing agents such as methyl-3-[p-azidophenyl] dithiolpropionimidate yield photoactivatable intermediates that are capable of forming crosslinks in the presence of light. Alternatively, reactive water-insoluble matrices such as cyanogen bromide-activated carbohydrates and the reactive substrates described in U.S. Patent Nos. 3,969,287; 3,691,016; 4,195,128; 4,247,642; 4,229,537; and 4,330,440 are employed for protein immobilization.

Other modifications include hydroxylation of proline and lysine, phosphorylation of hydroxyl groups of seryl or threonyl residues, methylation of the α -amino groups of lysine, arginine, and histidine side chains (Creighton, T.E., Proteins: Structure and Molecular Properties, W.H. Freeman & Co., San Francisco, pp. 79-86 (1983)), acetylation of the N-terminal amine, and, in some instances, amidation of the C-terminal carboxyl groups.

Such derivatized moieties may improve the stability, solubility, absorption, biological half life, and the like. The moieties may alternatively eliminate or attenuate any undesirable side effect of the protein complex and the like. Moieties capable of mediating such effects are disclosed, for example, in Remington's
5 Pharmaceutical Sciences, 18th ed., Mack Publishing Co., Easton, PA (1990).

The term "fragment" is used to indicate a polypeptide derived from the amino acid sequence of the proteins, of the complexes having a length less than the full-length polypeptide from which it has been derived. Such a fragment may, for example, be produced by proteolytic cleavage of the full-length protein. Preferably,
10 the fragment is obtained recombinantly by appropriately modifying the DNA sequence encoding the proteins to delete one or more amino acids at one or more sites of the C-terminus, N-terminus, and/or within the native sequence. Fragments of a protein are useful for screening for substances that act to modulate signal transduction, as described herein. It is understood that such fragments may retain
15 one or more characterizing portions of the native complex. Examples of such retained characteristics include: catalytic activity; substrate specificity; interaction with other molecules in the intact cell; regulatory functions; or binding with an antibody specific for the native complex, or an epitope thereof.

Another functional derivative intended to be within the scope of the present
20 invention is a "variant" polypeptide which either lacks one or more amino acids or contains additional or substituted amino acids relative to the native polypeptide. The variant may be derived from a naturally occurring complex component by appropriately modifying the protein DNA coding sequence to add, remove, and/or to modify codons for one or more amino acids at one or more sites of the C-terminus,
25 N-terminus, and/or within the native sequence. It is understood that such variants having added, substituted and/or additional amino acids retain one or more characterizing portions of the native protein, as described above.

A functional derivative of a protein with deleted, inserted and/or substituted amino acid residues may be prepared using standard techniques well-known to those

of ordinary skill in the art. For example, the modified components of the functional derivatives may be produced using site-directed mutagenesis techniques (as exemplified by Adelman *et al.*, 1983, *DNA* 2:183) wherein nucleotides in the DNA coding the sequence are modified such that a modified coding sequence is modified, and thereafter expressing this recombinant DNA in a prokaryotic or eukaryotic host cell, using techniques such as those described above. Alternatively, proteins with amino acid deletions, insertions and/or substitutions may be conveniently prepared by direct chemical synthesis, using methods well-known in the art. The functional derivatives of the proteins typically exhibit the same qualitative biological activity as the native proteins.

TABLES AND DESCRIPTION THEREOF

This patent describes novel protease identified in databases of genomic sequence. The results are summarized in four tables, which are described below.

Table 1 documents the name of each gene, the classification of each gene, the positions of the open reading frames within the sequence, and the length of the corresponding peptide. From left to right the data presented is as follows: "Gene Name", "ID#na", "ID#aa", "FL/Cat", "Superfamily", "Group", "Family", "NA_length", "ORF Start", "ORF End", "ORF Length", and "AA_length". "Gene name" refers to name given the sequence encoding the protease enzyme. Each gene is represented by "SGPr" designation followed by an arbitrary number. The SGPr name usually represents multiple overlapping sequences built into a single contiguous sequence (a "contig"). The "ID#na" and "ID#aa" refer to the identification numbers given each nucleic acid and amino acid sequence in this patent application. "FL/Cat" refers to the length of the gene, with FL indicating full length, and "Cat" indicating that only the catalytic domain is presented. "Partial" in this column indicates that the sequence encodes a partial catalytic domain. "Superfamily" identifies whether the gene is a protease. "Group" and "Family"

refer to the protease classification defined by sequence homology. "NA_length" refers to the length in nucleotides of the corresponding nucleic acid sequence. "ORF start" refers to the beginning nucleotide of the open reading frame. "ORF end" refers to the last nucleotide of the open reading frame, including the stop codon.

- 5 "ORF length" refers to the length in nucleotides of the open reading frame (including the stop codon). "AA length" refers to the length in amino acids of the peptide encoded in the corresponding nucleic acid sequence.

Table 1 - Proteases

Gene Name	IDna	IDaa	FL/Cat	Superfamily	Group	Family	MA Length	ORF Start	ORF End	ORF Length	AA Length
SGP397	1	60	FL	Protease	Carboxypeptidase	Zn carboxypeptidase	848	1	948	948	316
SGP413	2	81	FL	Protease	Carboxypeptidase	Zn carboxypeptidase	1126	1	1125	1125	374
SGP404	3	62	FL	Protease	Carboxypeptidase	Zn carboxypeptidase	1590	1	1590	1590	528
SGP638	4	63	FL	Protease	Cysteine	papain	1404	1	1404	1404	467
SGP414	5	64	FL	Protease	Cysteine	UCH2b	10062	1	10062	10062	3353
SGP430	6	65	FL	Protease	Cysteine	UCH2b	2843	1	2843	2843	980
SGP498	7	66	FL	Protease	Cysteine	UCH2b	2862	1	2862	2862	953
SGP485	8	67	FL	Protease	Cysteine	UCH2b	2362	1	2362	2362	783
SGP407	9	68	FL	Protease	Cysteine	UCH2b	2269	1	2269	2269	762
SGP453	10	69	FL	Protease	Cysteine	UCH2b	2139	1	2139	2139	712
SGP445	11	70	FL	Protease	Cysteine	UCH2b	870	1	870	870	289
SGP401	12	71	FL	Protease	Cysteine	UCH2b	1101	1	1101	1101	366
SGP408	13	72	FL	Protease	Cysteine	UCH2b	3884	1	3884	3884	1287
SGP480	14	73	FL	Protease	Cysteine	UCH2b	4816	1	4816	4816	1604
SGP431	15	74	FL	Protease	Cysteine	UCH2b	3129	1	3129	3129	1042
SGP426	16	75	FL	Protease	Cysteine	UCH2b	3102	1	3102	3102	1033
SGP403	17	76	FL	Protease	Cysteine	UCH2b	1554	1	1554	1554	517
SGP427	18	77	FL	Protease	Cysteine	UCH2b	3372	1	3372	3372	1123
SGP402	19	78	FL	Protease	Metalloprotease	PepM10	786	1	786	786	261
SGP459	20	79	FL	Protease	Metalloprotease	PepM10	1462	1	1462	1462	483
SGP104	21	80	FL	Protease	Metalloprotease	PepM13	2288	1	2288	2288	765
SGP303	22	81	CAT	Protease	Metalloprotease	PepM2	1267	1	1267	1267	418
SGP402	23	82	FL	Protease	Serine	subtilase	2268	1	2268	2268	765
SGP434	24	83	FL	Protease	Serine	trypsin	1178	1	1178	1178	381
SGP446	25	84	CAT	Protease	Serine	trypsin	681	1	681	681	228
SGP447	26	85	FL	Protease	Serine	trypsin	888	1	888	888	285
SGP432	27	86	FL	Protease	Serine	trypsin	1887	1	1887	1887	628
SGP428	28	87	FL	Protease	Serine	trypsin	831	1	831	831	276
SGP425	29	88	CAT	Protease	Serine	trypsin	858	1	858	858	285
SGP448	30	89	FL	Protease	Serine	trypsin	1242	1	1242	1242	413
SGP398	31	90	FL	Protease	Serine	trypsin	983	1	983	983	320
SGP428	32	91	FL	Protease	Serine	trypsin	987	1	987	987	328
SGP426	33	92	FL	Protease	Serine	trypsin	1278	1	1278	1278	425
SGP462	34	93	CAT	Protease	Serine	trypsin	666	1	666	666	221
SGP405	35	94	FL	Protease	Serine	trypsin	2847	1	2847	2847	948
SGP485	36	95	FL	Protease	Serine	trypsin	1059	1	1059	1059	352
SGP634	37	96	FL	Protease	Serine	trypsin	762	1	762	762	263
SGP621	38	97	FL	Protease	Serine	trypsin	3387	1	3387	3387	1128
SGP630	39	98	FL	Protease	Serine	trypsin	762	1	762	762	253
SGP620	40	99	CAT	Protease	Serine	trypsin	816	1	816	816	271
SGP455	41	100	FL	Protease	Serine	trypsin	1737	1	1737	1737	578
SGP607	42	101	FL	Protease	Serine	trypsin	2913	1	2913	2913	970
SGP607	43	102	FL	Protease	Serine	trypsin	788	1	788	788	265
SGP658	44	103	FL	Protease	Serine	trypsin	1365	1	1365	1365	454
SGP687	45	104	FL	Protease	Serine	trypsin	1614	1	1614	1614	537
SGP478	46	105	FL	Protease	Serine	trypsin	981	1	981	981	326
SGP489	47	106	CAT	Protease	Serine	trypsin	1671	1	1671	1671	556
SGP405	48	107	CAT	Protease	Serine	trypsin	884	1	884	884	287
SGP624	49	108	FL	Protease	Serine	trypsin	2553	1	2553	2553	850
SGP422	50	109	FL	Protease	Serine	trypsin	1344	1	1344	1344	447
SGP638	51	110	FL	Protease	Serine	trypsin	1374	1	1374	1374	457
SGP627	52	111	FL	Protease	Serine	trypsin	2457	1	2457	2457	818
SGP642	53	112	FL	Protease	Serine	trypsin	855	1	855	855	284
SGP651	54	113	FL	Protease	Serine	trypsin	2409	1	2409	2409	802
SGP451	55	114	FL	Protease	Serine	trypsin	1080	1	1080	1080	359
SGP452	56	115	FL	Protease	Serine	trypsin	867	1	867	867	288
SGP604	57	116	Partial	Protease	Serine	trypsin	135	1	135	135	44
SGP488	58	117	Partial	Protease	Serine	trypsin	138	1	138	138	45
SGP400	59	118	Partial	Protease	Serine	trypsin	930	1	930	930	309

Table 2 lists the following features of the genes described in this patent application: chromosomal localization, single nucleotide polymorphisms (SNPs), representation in dbEST, and repeat regions. From left to right the data presented is as follows: "Gene Name", "ID#na", "ID#aa", "FL/Cat", "Superfamily", "Group", "Family", "Chromosome", "SNPs", "dbEST_hits", & "Repeats". The contents of the first 7 columns (i.e., "Gene Name", "ID#na", "ID#aa", "FL/Cat", "Superfamily", "Group", "Family") are as described above for Table 1. "Chromosome" refers to the cytogenetic localization of the gene. Information in the "SNPs" column describes the nucleic acid position and degenerate nature of candidate single nucleotide polymorphisms (SNPs; please see table of polymorphism below). These SNPs were identified by blastn of the DNA sequence against the database of single nucleotide polymorphisms maintained at NCBI (<http://www.ncbi.nlm.nih.gov/SNP/snpblastByChr.html>). "dbEST hits" lists accession numbers of entries in the public database of ESTs (dbEST, <http://www.ncbi.nlm.nih.gov/dbEST/index.html>) that contain at least 150 bp of 100% identity to the corresponding gene. These ESTs were identified by blastn of dbEST. "Repeats" contains information about the location of short sequences, approximately 20 bp in length, that are of low complexity and that are present in several distinct genes.

Gene Name	Ortho	Data	FL/CAT	Superfamily	Group	Family	Chromosome	SNP*	dbEST hits	Repeats
SGP430	6	65	FL	Protease	Cysteine	UCH2b	2q37			
SGP490	7	68	FL	Protease	Cysteine	UCH2b	X011.4			
SGP495	8	67	FL	Protease	Cysteine	UCH2b	6p16			
SGP407	9	68	FL	Protease	Cysteine	UCH2b	2q37			
SGP453	10	69	FL	Protease	Cysteine	UCH2b	12q23			
SGP448	11	70	FL	Protease	Cysteine	UCH2b	6p16			
SGP401	12	71	FL	Protease	Cysteine	UCH2b	4q11			
SGP400	13	72	FL	Protease	Cysteine	UCH2b	17p15			
SGP480	14	73	FL	Protease	Cysteine	UCH2b	17q24			
SGP431	15	74	FL	Protease	Cysteine	UCH2b	4q13.3			
SGP429	16	75	FL	Protease	Cysteine	UCH2b	13q32			
SGP603	17	76	FL	Protease	Cysteine	UCH2b	12q23.3			
SGP427	18	77	FL	Protease	Cysteine	UCH2b	17q13			
SGP402	19	78	FL	Protease	Metalloprotease	Pgm10	17p15			
SGP459	20	79	FL	Protease	Metalloprotease	Pgm10	17q23			
SGP103	21	80	FL	Protease	Metalloprotease	Pgm10	3q27			
SGP103	22	81	CAT	Protease	Serine	subbase	17q11.1			
SGP102	23	82	FL	Protease	Serine	subbase	19q11			
SGP414	24	83	FL	Protease	Serine	subbase	3q21			
SGP448	25	84	CAT	Protease	Serine	subbase	3q21			
SGP447	26	85	FL	Protease	Serine	subbase	18q13.3			
SGP432	27	86	FL	Protease	Serine	subbase	Unknown			
SGP429	28	87	FL	Protease	Serine	subbase	19q13.4			
SGP428	29	88	CAT	Protease	Serine	subbase	6p23			
SGP445	30	89	FL	Protease	Serine	subbase	6p14			
SGP428	31	90	FL	Protease	Serine	subbase	19q13.4			
SGP496	32	91	FL	Protease	Serine	subbase	4q32			
SGP428	33	92	FL	Protease	Serine	subbase	4q13			
SGP452	34	93	CAT	Protease	Serine	subbase	16p13.3			
SGP405	35	94	FL	Protease	Serine	subbase	Unknown			
SGP405	36	95	FL	Protease	Serine	subbase	9p23			
SGP634	37	96	FL	Protease	Serine	subbase	16q23			
SGP630	38	97	FL	Protease	Serine	subbase	19q11			
SGP621	39	98	FL	Protease	Serine	subbase	19q13.4			
SGP630	40	99	CAT	Protease	Serine	subbase	9q23			
SGP630	41	100	FL	Protease	Serine	subbase	2q37			
SGP455	42	101	FL	Protease	Serine	subbase	12p11.2			
SGP407	43	102	FL	Protease	Serine	subbase	17q26			
SGP659	44	103	FL	Protease	Serine	subbase	21q22			
SGP487	45	104	FL	Protease	Serine	subbase	11q23			
SGP478	46	105	FL	Protease	Serine	subbase	19q22			
SGP489	47	106	CAT	Protease	Serine	subbase	17p15			
SGP465	48	107	CAT	Protease	Serine	subbase	Unknown			
SGP652	49	108	FL	Protease	Serine	subbase	Unknown			
SGP422	50	109	FL	Protease	Serine	subbase	4q13			
SGP458	51	110	FL	Protease	Serine	subbase	11q23			
SGP657	52	111	FL	Protease	Serine	subbase	Unknown			
SGP642	53	112	FL	Protease	Serine	subbase	18q13.1			
SGP651	54	113	FL	Protease	Serine	subbase	22q13			
SGP451	55	114	FL	Protease	Serine	subbase	12q23			
SGP452	56	115	FL							

Table 3 lists the extent and the boundaries of the protease catalytic domains, and other protein domains. The column headings are: "Gene Name", "ID#na", "ID#aa", "FL/Cat", "Profile_start", "Profile_end", "Domain_start", "Domain_end", and "Profile".

5 The contents of the first 7 columns (i.e., "Gene Name", "ID#na", "ID#aa", "FL/Cat", "Superfamily", "Group", "Family") are as described above for Table 1. "Profile_start", "Profile_end", "Domain_start" and "Domain_end" refer to data obtained using a Hidden-Markov Model to define catalytic range boundaries. The boundaries of the catalytic domain(s) within the overall protein are noted in the

10 "Domain_start" and "Domain_end" columns. "Profile" indicates the identity of the Hidden-Markov Model used to identify catalytic and other types of domains within the protein sequence. Whether the HMMr search was done with a complete ("Global") or Smith Waterman ("Local") model, is described below. Starting from a multiple sequence alignment of catalytic domains, two hidden Markov models were

15 built. One of them allows for partial matches to the catalytic domain; this is a "local" HMM, similar to Smith-Waterman alignments in sequence matching. The other model allows matches only to the complete catalytic domain; this is a "global" HMM similar to Needleman-Wunsch alignments in sequence matching. The Smith Waterman local model is more specific, allowing for fragmentary matches to the

20 catalytic domain whereas the global "complete" model is more sensitive, allowing for remote homologue identification. These domains were identified using PFAM (<http://pfam.wustl.edu/hmmsearch.shtml>) models, a large collection of multiple sequence alignments and hidden Markov models covering many common protein domains. Version 5.5 of Pfam (Sept 2000) contains alignments and models for 2478

25 protein families (<http://pfam.wustl.edu/faq.shtml>). The PFAM alignments were downloaded from <http://pfam.wustl.edu/hmmsearch.shtml> and the HMMr searches were run locally on a Timelogic computer (TimeLogic Corporation, Incline Village, NV). A number of proteins have more than one domain recognized by the HMM searches. For these proteins, the domains have been listed in separate rows.

Table 3 - Protease Domains, Other Domains

Gene Name	ID#na	ID#aa	FL/Cat	Profile_start	Profile_end	Domain_start	Domain_end	Profile
SGP1397	1	60	FL	1	146	139	280	Zn carboxypeptidase (PF00246)
SGP1397	1	60	FL	1	82	41	120	Carboxypeptidase activation peptide
SGP1413	2	61	FL	1	248	50	291	Zn carboxypeptidase (PF00246)
SGP1404	3	62	FL	1	248	91	466	Zn carboxypeptidase (PF00246)
SGP1536	4	63	FL	1	337	203	456	papain (PF00112)
SGP1414	5	64	FL	1	72	1851	2045	Ubiquitin carboxyl-terminal hydrolase family 2b (PF00443)
SGP1414	5	64	FL	1	32	1701	1732	UCH2b (PF00442)
SGP1430	6	65	FL	1	72	886	951	Ubiquitin carboxyl-terminal hydrolase family 2b (PF00443)
SGP1430	6	65	FL	1	32	342	373	UCH2b (PF00442)
SGP1498	7	66	FL	1	72	875	935	Ubiquitin carboxyl-terminal hydrolase family 2b (PF00443)
SGP1498	7	66	FL	1	32	593	624	UCH2b (PF00442)
SGP1498	7	66	FL	1	82	465	534	Zn-finger in ubiquitin-hydrolases (PF02148)
SGP1498	8	67	FL	1	72	695	781	Ubiquitin carboxyl-terminal hydrolase family 2b (PF00443)
SGP1498	8	67	FL	1	32	190	221	UCH2b (PF00442)
SGP1498	8	67	FL	1	72	82	148	Zn-finger in ubiquitin-hydrolases (PF02148)
SGP1498	9	68	FL	80	90	481	491	Ubiquitin carboxyl-terminal hydrolase family 2b (PF00443)
SGP1498	10	69	FL	1	72	615	677	Ubiquitin carboxyl-terminal hydrolase family 2b (PF00443)
SGP1498	10	69	FL	1	32	273	304	UCH2b (PF00442)
SGP1498	10	69	FL	1	82	29	99	Zn-finger in ubiquitin-hydrolases (PF02148)
SGP1445	11	70	FL	1	32	190	221	Ubiquitin carboxyl-terminal hydrolase family 2b (PF00443)
SGP1445	11	70	FL	7	82	78	148	Zn-finger in ubiquitin-hydrolases (PF02148)
SGP1401	12	71	FL	1	72	292	364	Ubiquitin carboxyl-terminal hydrolase family 2b (PF00443)
SGP1401	12	71	FL	1	32	35	66	UCH2b (PF00442)
SGP1408	13	72	FL	1	72	395	475	Ubiquitin carboxyl-terminal hydrolase family 2b (PF00443)
SGP1408	13	72	FL	1	32	100	131	UCH2b (PF00442)
SGP1480	14	73	FL	1	72	1506	1586	Ubiquitin carboxyl-terminal hydrolase family 2b (PF00443)
SGP1480	14	73	FL	1	32	734	765	UCH2b (PF00442)
SGP1480	14	73	FL	1	29	268	296	EF hand (PF00036)
SGP1480	14	73	FL	1	29	232	260	EF hand (PF00036)
SGP1431	15	74	FL	1	72	838	948	Ubiquitin carboxyl-terminal hydrolase family 2b (PF00443)
SGP1431	15	74	FL	1	32	445	478	UCH2b (PF00442)
SGP1429	16	75	FL	1	72	332	419	Ubiquitin carboxyl-terminal hydrolase family 2b (PF00443)
SGP1429	16	75	FL	1	32	89	120	UCH2b (PF00442)
SGP1503	17	76	FL	1	72	432	501	Ubiquitin carboxyl-terminal hydrolase family 2b (PF00443)
SGP1503	17	76	FL	1	32	88	99	UCH2b (PF00442)
SGP1427	18	77	FL	1	72	648	709	Ubiquitin carboxyl-terminal hydrolase family 2b (PF00443)
SGP1427	18	77	FL	1	29	101	129	UCH2b (PF00442)
SGP1092	19	78	FL	49	168	75	194	Peptidase M10 (PF00413)
SGP1092	19	78	FL	168	179	207	218	ADAM (PF00413)
SGP1359	20	79	FL	1	168	44	212	Peptidase M10 (PF00413)
SGP1359	20	79	FL	1	50	302	443	3 x Hemaphysin (PF00045)
SGP1014	21	80	FL	1	222	561	764	Peptidase M13 (PF01431)
SGP1003	22	81	CAT	1	416	10	307	Peptidase M1 (PF01133)
SGP1402	23	82	FL	1	360	118	437	subtilase (PF00082)
SGP1434	24	83	FL	129	136	39	46	p20-ICE (PF00858)
SGP1448	25	84	CAT	1	242	13	227	Trypsin (PF00089)
SGP1447	26	85	FL	1	259	33	270	Trypsin (PF00089)
SGP1432	27	86	FL	6	259	117	343	Trypsin (PF00089)
SGP1529	28	87	FL	413	416	184	187	Trypsin (PF00089)
SGP1428	29	88	CAT	7	259	24	246	Trypsin (PF00089)
SGP1425	30	89	FL	387	408	287	308	Trypsin (PF00089)
SGP1548	31	90	FL	1	259	88	313	Trypsin (PF00089)
SGP1398	32	91	FL	1	259	28	262	Trypsin (PF00089)
SGP1426	33	92	FL	1	259	194	419	Trypsin (PF00089)
SGP1552	34	93	CAT	1	255	2	222	Trypsin (PF00089)
SGP1405	35	94	FL	60	259	218	406	Trypsin (PF00089)
SGP1405	35	94	FL	126	209	419	488	Trypsin (PF00089)
SGP1405	35	94	FL	122	251	639	781	Trypsin (PF00089)
SGP1485	36	95	FL	1	259	88	285	Trypsin (PF00089)
SGP1534	37	96	FL	1	259	34	258	Trypsin (PF00089)
SGP1390	38	97	FL	1	259	898	1122	Trypsin (PF00089)
SGP1390	38	97	FL	1	259	264	500	Trypsin (PF00089)
SGP1390	38	97	FL	1	259	573	800	Trypsin (PF00089)
SGP1521	39	98	FL	1	259	30	245	Trypsin (PF00089)
SGP1530	40	99	CAT	1	259	14	255	Trypsin (PF00089)
SGP1520	41	100	FL	1	259	73	306	Trypsin (PF00089)
SGP1455	42	101	FL	1	259	433	674	Trypsin (PF00089)
SGP1455	42	101	FL	109	259	4	156	Trypsin (PF00089)
SGP1455	42	101	FL	2	118	175	812	3 x CUB domains (PF00431)
SGP1507	43	102	FL	35	148	42	135	Trypsin (PF00089)
SGP1507	43	102	FL	247	259	248	258	Trypsin (PF00089)
SGP1559	44	103	FL	1	259	217	444	Trypsin (PF00089)
SGP1559	44	103	FL	1	43	71	109	Low-density lipoprotein receptor domain class A (PF00057)
SGP1587	45	104	FL	1	259	268	524	Trypsin (PF00089)
SGP1476	46	105	FL	1	259	60	288	Trypsin (PF00089)
SGP1489	47	106	CAT	1	227	58	257	Trypsin (PF00089)
SGP1489	47	106	CAT	1	118	304	533	2 x CUB domains (PF00431)
SGP1485	48	107	FL	12	259	2	240	Trypsin (PF00089)
SGP1524	49	108	FL	1	259	613	842	Trypsin (PF00089)
SGP1524	49	108	FL	1	43	489	603	3 x Low-density lipoprotein receptor domain class A (PF00057)
SGP1422	50	109	FL	1	259	218	441	Trypsin (PF00089)
SGP1538	51	110	FL	1	259	218	448	Trypsin (PF00089)
SGP1527	52	111	FL	1	259	47	266	Trypsin (PF00089)
SGP1527	52	111	FL	1	158	323	454	Trypsin (PF00089)
SGP1527	52	111	FL	12	149	564	678	Trypsin (PF00089)
SGP1542	53	112	FL	1	259	35	259	Trypsin (PF00089)
SGP1551	54	113	FL	1	259	568	797	Trypsin (PF00089)
SGP1551	54	113	FL	1	43	447	559	3 x Low-density lipoprotein receptor domain class A (PF00057)
SGP1451	55	114	FL	1	259	89	324	Trypsin (PF00089)
SGP1452	56	115	FL	1	259	73	280	Trypsin (PF00089)
SGP1504	57	116	Partial	1	52	1	45	Trypsin (PF00089)
SGP1469	58	117	Partial	210	259	1	48	Trypsin (PF00089)
SGP1400	59	118	Partial	1	198	133	281	Trypsin (PF00089)

Table 4 describes the results of Smith Waterman similarity searches (Matrix: Pam100; gap open/extension penalties 12/2) of the amino acid sequences against the NCBI database of non-redundant protein sequences

- 5 (<http://www.ncbi.nlm.nih.gov/Entrez/protein.html>). The column headings are: "Gene Name", "ID#na", "ID#aa", "FL/Cat", "Superfamily", "Group", "Family", "Pscore", "aa_length", "aa_ID_match", "%Identity", "%Similar", "ACC#_nraa_match", and "Description". The contents of the first 7 columns (i.e., "Gene Name", "ID#na", "ID#aa", "FL/Cat", "Superfamily", "Group", "Family") are
 10 as described above for Table 1. "Pscore" refers to the Smith Waterman probability score. This number approximates the chance that the alignment occurred by chance. Thus, a very low number, such as $2.10E-64$, indicates that there is a very significant match between the query and the database target. "aa_length" refers to the length of the protein in amino acids. "aa_ID_match" indicates the number of amino acids that
 15 were identical in the alignment. "% Identity" lists the percent of amino acids that were identical over the aligned region. "% Similarity" lists the percent of amino acids that were similar over the alignment. "ACC#nraa_match" lists the accession number of the most similar protein in the NCBI database of non-redundant proteins. "Description" contains the name of the most similar protein in the NCBI database of
 20 non-redundant proteins.

Table 4
Smith Waterman

Gene Name	Unig	Accession	FL/Cat	Superfamily	Group	Family	Protein	aa length	% ID match	% Identity	% Similar	ACSW area match	Description
SGP437	1	60	FL	Protease	Carboxypeptidase	Zn carboxypeptidase	3,10E-220	315	100	100	100	NP_055994.1	Carboxypeptidase B precursor (Homo sapiens)
SGP438	2	81	FL	Protease	Carboxypeptidase	Zn carboxypeptidase	5.00E-09	374	148	49	66	AA01344.1	Carboxypeptidase B precursor (Homo sapiens)
SGP439	3	62	FL	Protease	Carboxypeptidase	Zn carboxypeptidase	1.00E-276	407	502	84	96	NP_001355.1	Carboxypeptidase B precursor (Homo sapiens)
SGP440	4	63	FL	Protease	Cysteine	Uch2b	0	487	487	100	100	NP_071447.1	Protein (Homo sapiens)
SGP441	5	64	FL	Protease	Cysteine	Uch2b	0	353	1259	89	100	NP_055524.1	Protein (Homo sapiens)
SGP442	6	65	FL	Protease	Cysteine	Uch2b	0	180	630	89	96	BA012420.1	Protein (Homo sapiens)
SGP443	7	66	FL	Protease	Cysteine	Uch2b	2.00E-180	653	498	95	98	AA059533.1	Protein (Homo sapiens)
SGP444	8	67	FL	Protease	Cysteine	Uch2b	2.40E-178	763	282	100	100	AA059533.1	Protein (Homo sapiens)
SGP445	9	68	FL	Protease	Cysteine	Uch2b	2.00E-40	753	60	76	84	NP_036071.1	Protein (Homo sapiens)
SGP446	10	69	FL	Protease	Cysteine	Uch2b	3.60E-105	269	289	100	100	AA059533.1	Protein (Homo sapiens)
SGP447	11	70	FL	Protease	Cysteine	Uch2b	7.30E-234	369	369	100	100	NP_071443.1	Protein (Homo sapiens)
SGP448	12	71	FL	Protease	Cysteine	Uch2b	0	1267	1267	100	100	NP_115523.1	Protein (Homo sapiens)
SGP449	13	72	FL	Protease	Cysteine	Uch2b	0	1804	1272	99	99	NP_115971.1	Protein (Homo sapiens)
SGP450	14	73	FL	Protease	Cysteine	Uch2b	0	1042	397	100	100	NP_115946.1	Protein (Homo sapiens)
SGP451	15	74	FL	Protease	Cysteine	Uch2b	1.50E-250	1033	369	100	100	NP_115612.1	Protein (Homo sapiens)
SGP452	16	75	FL	Protease	Cysteine	Uch2b	0	517	509	100	100	AA013609.1	Protein (Homo sapiens)
SGP453	17	76	FL	Protease	Cysteine	Uch2b	1.50E-92	1123	269	36	53	AA013609.1	Protein (Homo sapiens)
SGP454	18	77	FL	Protease	Cysteine	Uch2b	4.70E-171	261	261	100	100	NP_011917.1	Protein (Homo sapiens)
SGP455	19	78	FL	Protease	Metalloprotease	Pap10	0	483	483	100	100	NP_004762.1	Protein (Homo sapiens)
SGP456	20	79	FL	Protease	Metalloprotease	Pap10	0	765	765	100	100	NP_055508.1	Protein (Homo sapiens)
SGP457	21	80	FL	Protease	Metalloprotease	Pap10	2.20E-284	419	407	97	98	CA010709.1	Protein (Homo sapiens)
SGP458	22	81	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP459	23	82	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP460	24	83	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP461	25	84	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP462	26	85	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP463	27	86	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP464	28	87	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP465	29	88	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP466	30	89	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP467	31	90	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP468	32	91	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP469	33	92	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP470	34	93	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP471	35	94	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP472	36	95	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP473	37	96	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP474	38	97	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP475	39	98	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP476	40	99	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP477	41	100	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP478	42	101	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP479	43	102	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP480	44	103	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP481	45	104	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP482	46	105	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP483	47	106	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP484	48	107	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP485	49	108	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP486	50	109	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP487	51	110	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP488	52	111	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP489	53	112	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP490	54	113	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP491	55	114	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP492	56	115	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP493	57	116	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP494	58	117	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP495	59	118	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP496	60	119	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP497	61	120	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP498	62	121	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP499	63	122	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP500	64	123	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP501	65	124	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP502	66	125	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP503	67	126	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP504	68	127	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP505	69	128	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP506	70	129	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP507	71	130	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP508	72	131	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP509	73	132	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP510	74	133	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP511	75	134	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP512	76	135	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP513	77	136	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP514	78	137	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP515	79	138	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP516	80	139	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP517	81	140	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP518	82	141	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP519	83	142	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP520	84	143	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP521	85	144	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP522	86	145	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP523	87	146	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP524	88	147	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP525	89	148	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP526	90	149	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP527	91	150	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP528	92	151	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP529	93	152	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP530	94	153	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP531	95	154	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP532	96	155	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP533	97	156	FL	Protease	Metalloprotease	Pap10	0	755	613	82	89	NP_055508.1	Protein (Homo sapiens)
SGP534	98												

EXAMPLES

The examples below are not limiting and are merely representative of various aspects and features of the present invention. The examples below demonstrate the isolation and characterization of the proteases of the invention.

5

EXAMPLE 1: Identification of Genomic Fragments Encoding Proteases

- Novel proteases were identified from the Celera human genomic sequence databases, and from the public Human Genome Sequencing project
- 10 (<http://www.ncbi.nlm.nih.gov/>) using hidden Markov models (HMMR). The genomic database entries were translated in six open reading frames and searched against the model using a Timelogic Decypher box with a Field programmable array (FPGA) accelerated version of HMMR2.1. The DNA sequences encoding the predicted protein sequences aligning to the HMMR profile were extracted from the
- 15 original genomic database. The nucleic acid sequences were then clustered using the Pangea Clustering tool to eliminate repetitive entries. The putative protease sequences were then sequentially run through a series of queries and filters to identify novel protease sequences. Specifically, the HMMR identified sequences were searched using BLASTN and BLASTX against a nucleotide and amino acid
- 20 repository containing known human proteases and all subsequent new protease sequences as they are identified. The output was parsed into a spreadsheet to facilitate elimination of known genes by manual inspection. Two models were used, a "complete" model and a "partial" or Smith Waterman model. The partial model was used to identify sub-catalytic domains, whereas the complete model was used to
- 25 identify complete catalytic domains. The selected hits were then queried using BLASTN against the public NRNA and EST databases to confirm they are indeed unique.

Extension of partial DNA sequences to encompass the longer sequences, including full-length open-reading frame, was carried out by several methods. Iterative blastn searching of the cDNA databases listed in Table 5 was used to find cDNAs that extended the genomic sequences. "LifeGold" databases are from Incyte
5 Genomics, Inc (<http://www.incyte.com/>). NCBI databases are from the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/>). All blastn searches were conducted using a penalty for a nucleotide mismatch of -3 and reward for a nucleotide match of 1. The gapped blast algorithm is described in: Altschul, Stephen F., Thomas L. Madden, Alejandro A. Schaffer, Jinghui Zhang, Zheng
10 Zhang, Webb Miller, and David J. Lipman (1997), "Gapped BLAST and PSI-BLAST: a new generation of protein database search-programs", *Nucleic Acids Res.* 25:3389-3402).

Extension of partial DNA sequences to encompass the full-length open-reading frame was also carried out by iterative searches of genomic databases. The first
15 method made use of the Smith-Waterman algorithm to carry out protein-protein searches of the closest homologue or orthologue to the partial. The target databases consisted of Genscan [Chris Burge and Sam Karlin "Prediction of Complete Gene Structures in Human Genomic DNA", *JMB* (1997) 268(1):78-94] and open-reading frame (ORF) predictions of all human genomic sequence derived from the human
20 genome project (HGP) as well as from Celera. The complete set of genomic databases searched is shown in Table 6 below. Genomic sequences encoding potential extensions were further assessed by blastp analysis against the NCBI nonredundant to confirm the novelty of the hit. The extending genomic sequences were incorporated into the cDNA sequence after removal of potential introns using
25 the Seqman program from DNASTar. The default parameters used for Smith-Waterman searches were Matrix: PAM100; gap-opening penalty: 12; gap extension penalty: 2. Genscan predictions were made using the Genscan program as detailed in Chris Burge and Sam Karlin "Prediction of Complete Gene Structures in Human

Genomic DNA", JMB (1997) 268(1):78-94). ORF predictions from genomic DNA were made using a standard 6-frame translation.

Another method for defining DNA extensions from genomic sequence used iterative searches of genomic databases through the Genscan program to predict exon splicing [Burge and Karlin, JMB (1997) 268(1):78-94)]. These predicted genes were then assessed to see if they represented "real" extensions of the partial genes based on homology to related proteases.

Another method involved using the Genewise program (<http://www.sanger.ac.uk/Software/Wise2/>) to predict potential ORFs based on homology to the closest orthologue/homologue. Genewise requires two inputs, the homologous protein, and genomic DNA containing the gene of interest. The genomic DNA was identified by blastn searches of Celera and Human Genome Project databases. The orthologs were identified by blastp searches of the NCBI non-redundant protein database (NR). Genewise compares the protein sequence to a genomic DNA sequence, allowing for introns and frameshifting errors.

Table 5. Databases used for cDNA-based sequence extensions

Database	Database Date
LifeGold templates	May 2001
LifeGold compseqs	May 2001
LifeGold compseqs	May 2001
LifeGold compseqs	May 2001
LifeGold fl	May 2001
LifeGold flt	May 2001
NCBI human Ests	May 2001
NCBI murine Ests	May 2001
NCBI nonredundant	May 2001

TABLE 6. DATABASES USED FOR GENOMIC-BASED SEQUENCE EXTENSIONS

Database	Number of entries	Database Date
Celera v. 1-5	5,306,158	Jan 2000
Celera v. 6-10	4,209,980	May 2000
Celera v. 11-14	7,222,425	April 2000
Celera v. 15	243,044	April 2000
Celera v. 16-17	25,885	April 2000
Celera Assembly 5 (release 25h)	479,986	May 2001
HGP Phase 0	3,189	Nov 1/00
HGP Phase 1	20,447	Jan 1/01
HGP Phase 2	1,619	Jan 1/01
HGP Phase 3	9,224	May 2001
HGP Chromosomal assemblies	2759	May 2001

5 Results:

The sources for the sequence information used to extend the genes in the provisional patents are listed below. For genes that were extended using Genewise, the accession numbers of the protein ortholog and the genomic DNA are given. (Genewise uses the ortholog to assemble the coding sequence of the target gene from the genomic sequence). The amino acid sequences for the orthologs were obtained from the NCBI non-redundant database of proteins (<http://www.ncbi.nlm.nih.gov/Entrez/protein.html>). The genomic DNA came from two sources: Celera and NCBI-NRNA, as indicated below. cDNA sources are also listed below. All of the genomic sequences were used as input for Genscan predictions to predict splice sites [Burge and Karlin, JMB (1997)

268(1):78-94)]. Abbreviations: HGP: Human Genome Project; NCBI, National Center for Biotechnology Information.

- SGPr397, SEQ ID NO:1; SEQ ID NO:60
- 5 Genewise orthologs: BAB25826.1, XP_005284.2, NP_065094.1.
Genomic DNA sources: Celera_asm5h 181000001172043
cDNA Sources: Public (gi|9966830|ref|NM_020361.1)
- SGPr413, SEQ ID NO:2; SEQ ID NO:61
- 10 Genewise orthologs: gi|6013463, XP_003009.1, P15086.
Genomic DNA sources: Celera_asm5h 300475633
SGPr404, SEQ ID NO:3; SEQ ID NO:62
Genewise orthologs: BAB31768.1, NP_061355.1, AAH03713.
- 15 Genomic DNA sources: Celera_asm5h 90000641768196
SGPr536_1, SEQ ID NO:4; SEQ ID NO:63
Genewise orthologs: BAB18637
Genomic DNA sources: 90000642234172
- 20 SGPr414, SEQ ID NO:5; SEQ ID NO:64
Genewise orthologs: AAF50752.1
Genomic DNA sources: 90000628114448
cDNA Sources: AK023845.1|AK023845 Homo sapiens cDNA FLJ13783 fis; Incyte
25 399773.5
SGPr430, SEQ ID NO:6; SEQ ID NO:65
Genewise orthologs: NP_065954.
Genomic DNA sources: 301015601

cDNA Sources:AB046814 Homo sapiens mRNA for KIAA1594.

SGPr496_1, SEQ ID NO:7, SEQ ID NO:66

Genewise orthologs: AAH07196, AAF66953.

5 Genomic DNA sources:90000627702299

SGPr495, SEQ ID NO:8, SEQ ID NO:67

Genewise orthologs: NP_006438.

Genomic DNA sources:90000627041101

10

SGPr407, SEQ ID NO:9, SEQ ID NO:68

Genewise orthologs: BAB27431, AAH03130, NP_057656.

Genomic DNA sources:92000003986525

15 SGPr453, SEQ ID NO:10, SEQ ID NO:69

Genewise orthologs: NP_006528.

Genomic DNA sources:90000640175777

cDNA Sources:AL136825.1|HSM801793 Homo sapiens mRNA; Incyte 428428.1

20 SGPr445, SEQ ID NO:11, SEQ ID NO:70

Genewise orthologs: NP_006438.

Genomic DNA sources:90000627041101

cDNA Sources:9863487 328 bp ubiquitin carboxyl-terminal hydrolase; Incyte
4802789CA2

25

SGPr401_1, SEQ ID NO:12, SEQ ID NO:71

Genewise orthologs: BAB14881, NP_073743, BAB24720.

Genomic DNA sources:92000004473288

cDNA Sources:NM_022832.1| Homo sapiens hypothetical protein FLJ12552
(FLJ12552)

SGPr408, SEQ ID NO:13, SEQ ID NO:72

5 Genewise orthologs: Q24574, AAF50752.

Genomic DNA sources:90000628565543.

cDNA Sources:AK027362.

SGPr480, SEQ ID NO:14, SEQ ID NO:73

10 Genewise orthologs: AAF49100, T29010.

Genomic DNA sources:90000640697688.

cDNA Sources:EF_hand; CAAX: NP_115971.

SGPr431, SEQ ID NO:15, SEQ ID NO:74

15 Genewise orthologs: AAK26248, BAA92610, Q92353.

Genomic DNA sources:90000642340202.

SGPr429, SEQ ID NO:16, SEQ ID NO:75

Genewise orthologs: BAB15591, AAG42764, gi_11358453.

20 Genomic DNA sources:90000642540891

cDNA Sources:AK026930.1|AK026930 Homo sapiens cDNA: FLJ23277.

SGPr503, SEQ ID NO:17, SEQ ID NO:76

Genewise orthologs: AAF40451, AAF46096, AAH04868.

25 Genomic DNA sources:90000642658172

cDNA Sources:BC004868.1|BC004868 Homo sapiens, clone MGC:10702; Incyte
5432879CB1.

SGPr427, SEQ ID NO:18, SEQ ID NO:77

Genewise orthologs: XP_003288, AAC27356, BAA86517.

Genomic DNA sources:181000001646773

cDNA Sources:Incyte 7485896CB1

5 SGPr092, SEQ ID NO:19, SEQ ID NO:78

Genewise orthologs: XP_011971.1, NP_068573.1, AAF80180.1.

Genomic DNA sources:Celera_asm5h 300261795

cDNA Sources:gi|12736016.

0 SGPr359, SEQ ID NO:20, SEQ ID NO:79

Genewise orthologs: 1.) gi|11545845|ref 2.) gi|12006364|gb 3.) gi|3511149|gb|A.

Genomic DNA sources:Celera_asm5h 90000642045264

cDNA Sources:gi|13639688.

5 SGPr104_1, SEQ ID NO:21, SEQ ID NO:80

Genewise orthologs: 1.) gi|7662200|ref.

Genomic DNA sources:HGP_s gi|12039078|4

cDNA Sources:NP_055508.1.

0 SGPr303, SEQ ID NO:22, SEQ ID NO:81

Genewise orthologs: CAA10709.1.

Genomic DNA sources:HGP_s gi|8082389_31

SGPr402_1, SEQ ID NO:23, SEQ ID NO:82

5 Genewise orthologs: A54306, , I77530, A45357

Genomic DNA sources:Celera_asm5h 92000004018126

SGPr434, SEQ ID NO:24, SEQ ID NO:83

Genewise orthologs: gi|6755819, gi|6912728, gi|8570164.

cDNA Sources:gi|6141221, gi|3754092, Incyte 1856589CB1.

- Genomic DNA sources:90000628646080

- Genewise orthologs: [gi_11181573](#), [gi_12832944](#), [gi_13124769](#), [gi_13277969](#).

- cDNA Sources:** Incyte EST 474674.1 (1975 to 1985 nucleotides) (GenBank)

- cDNA Sources:gi|4506157.

- Genomic DNA sources:90000627342893

- Genewise orthologs:** gi_12844896, gi_6005882.

Genomic DNA sources:181000004221955

cDNA Sources: Incyte Sequence 400833.1.

SGPr548, SEQ ID NO:31, SEQ ID NO:90

- 5 Genewise orthologs: gi|9957760, gi|5803199, gi|6681654.

Genomic DNA sources:92000003497776, gi|11178143

cDNA Sources:gi|9957759.

SGPr396, SEQ ID NO:32, SEQ ID NO:91

- 10 Genewise orthologs: gi_11055972, gi_12839280, gi_6680267, gi_8393560,
gi_9757698.

Genomic DNA sources:90000632590917

cDNA Sources: Incyte Sequence 7480224CB1.

- 15 SGPr426, SEQ ID NO:33, SEQ ID NO:92

Genewise orthologs: gi_13640890, gi_13646365, gi_7661558.

Genomic DNA sources:90000641479138

cDNA Sources:Incyte Sequence 7481056CB1.

- 20 SGPr552, SEQ ID NO:34, SEQ ID NO:93

Genewise orthologs: gi|7661558, gi|4758508.

Genomic DNA sources:90000641479138

SGPr405, SEQ ID NO:35, SEQ ID NO:94

- 25 Genewise orthologs: gi_7415931, gi_126839, gi_136423, gi_13183572.

Genomic DNA sources: gi|13509126

cDNA Sources:Incyte seqs 7474351CB1 and 134360.1.

SGPr485_1, SEQ ID NO:36, SEQ ID NO:95

Genomic DNA sources:90000627342893

cDNA Sources: Incyte Sequence 6026494CA2.

Genewise orthologs: [gi|4503135](#).

Genomic DNA sources:92000004436076, 165000101932709, 92000004433469

cDNA Sources: Incyte ESTs: 1383391.20 , 1383391.10 , 1383391.13 , 7691434H1, 2070278CB1, 741522CA2; NCBI ESTs: gi|7260671, gi|7260006, gi|7260642.

0 gi|7259962, gi|2018619, gi|7260655, gi|2019751. <http://www.ncbi.nlm.nih.gov/blast/>

SGPr390, SEO ID NO:38, SEO ID NO:97

Genewise orthologs: BAB23684

Genomic DNA sources: hCG22693 (100% identity) (GenBank accession number: U00180.1)

515 - Irving TST, PG 12190.

SGPr521, SEQ ID NO:39, SEQ ID NO:98

Genewise orthologs: BAB55604, AAF01139, AAF01139

Genomic DNA sources: HGP: sgi|11178143:10

cDNA Sources: gi|4826949; accession number U75090.1; GenBank

107. *Journal of the American Medical Association*, 1990; 263: 1033-1037.

SGPr530_1, SEQ ID NO:40, SEQ ID NO:99

Genewise orthologs: gi_12314133, NP_033381.1 3, NP_033382.1

Genomic DNA sources:Celera asm5h:181000001848433

5 SGPt520, SEQ ID NO:41, SEQ ID NO:100

Genewise orthologs: gi|12839535 gi|1352368; gi|4506151.

Genomic DNA sources:90000640807190

cDNA Sources: ESTs gi|13745759, 7472044CB1, 7474338CB1, gi|13703426, gi|5392427, gi|2142177, gi|2103202, LIB4218-103-R1-K1-H5, LIB4218-085-Q1-K1-C6, LIB4752-019-R1-K1-H4

5 SGPr455, SEQ ID NO:42, SEQ ID NO:101

Genewise orthologs: gi|7512178, gi|7512176.

Genomic DNA sources: 90000641321557

cDNA Sources: Incyte template 987279.1.

10 SGPr507_2, SEQ ID NO:43, SEQ ID NO:102

Genewise orthologs: gi|13385812, gi|12854692, gi|2499862.

Genomic DNA sources: 90000642611957

SGPr559, SEQ ID NO:44, SEQ ID NO:103

15 Genewise orthologs: XP_016993, BAB20079

Genomic DNA sources: Celera_asm5h 335001064013332

cDNA Sources: gi|13173471.

SGPr567_1, SEQ ID NO:45, SEQ ID NO:104

20 Genewise orthologs: NP_114435, Q9JIQ8

Genomic DNA sources: Celera_asm5h 90000642045213

cDNA Sources: gi|14042983|ref|NM_032046.1.

SGPr479_1, SEQ ID NO:46, SEQ ID NO:105

25 Genewise orthologs: NP_114154, NP_038949, NP_033382

Genomic DNA sources: 90000624931837

cDNA Sources: EST gi|13997890 and Incyte EST 7480124CB1,.

SGPr489_1, SEQ ID NO:47, SEQ ID NO:106

Genewise orthologs: gi|7512176, gi|7512178, gi|9757698.

Genomic DNA sources:90000628565500

SGPr465_1, SEQ ID NO:48, SEQ ID NO:107

5 Genewise orthologs: gi|6678293, gi|6678295.

Genomic DNA sources:gi|13431162

SGPr524_1, SEQ ID NO:49, SEQ ID NO:108

Genewise orthologs: gi|12836503, gi|10257390, gi|11415040.

10 Genomic DNA sources:90000626428259

SGPr422, SEQ ID NO:50, SEQ ID NO:109

Genewise orthologs: gi|7661558, gi|4758508.

Genomic DNA sources:90000641479138

15

SGPr538, SEQ ID NO:51, SEQ ID NO:110

Genewise orthologs: NP_110397, Q9ER04, NP_109634

Genomic DNA sources:Celera_asm5h 90000642044035 and 90000642045412

cDNA Sources:gi|13540535.

20

SGPr527_1, SEQ ID NO:52, SEQ ID NO:111

Genewise orthologs: gi|11181573, gi|13277969, gi|10441463.

Genomic DNA sources:90000631961624

cDNA Sources:Incye 2751509CB1.

25

SGPr542, SEQ ID NO:53, SEQ ID NO:112

Genewise orthologs: gi|1705760, gi|4885369.

Genomic DNA sources:92000004018116, gi|2896799, 92000004013323,

92000004013330, 165000100427031

SGPr551, SEQ ID NO:54, SEQ ID NO:113

Genewise orthologs: BAB23684.1, NP_035306.2, BAB03502.1

Genomic DNA sources: Celera_asm5h 90000643090998

5

SGPr451, SEQ ID NO:55, SEQ ID NO:114

Genewise orthologs: gi_5002340, gi|12018322, gi|1480413.

Genomic DNA sources: 181000000828193

0 SGPr452_1, SEQ ID NO:56, SEQ ID NO:115

Genewise orthologs: gi|13183572, gi|339983, gi|7415931.

Genomic DNA sources: 92000004034678

SGPr504, SEQ ID NO:57, SEQ ID NO:116

5 Genewise orthologs: gi|1633237

Genomic DNA sources: celera_asm5h 92000004018137

SGPr469, SEQ ID NO:58, SEQ ID NO:117

Genewise orthologs: BAB30277, CAB41988, XP_016204

0 Genomic DNA sources: GA_x2HTBKPYW7D

SGPr400, SEQ ID NO:59, SEQ ID NO:118

Genewise orthologs: gi|6755819, gi|6912728.

Genomic DNA sources: 90000632590917

5

DESCRIPTION OF NOVEL PROTEASE POLYNUCLEOTIDES

SGPr397, SEQ ID NO:1, SEQ ID NO:60 is 948 nucleotides long. The open reading frame starts at position 1 and ends at position 948, giving an ORF length of 948

nucleotides. The predicted protein is 315 amino acids long. This sequence codes for a full length protein. It is classified as (superfamily/group/family): Protease, Carboxypeptidase, Zn carboxypeptidase. The cytogenetic position of this gene is 8q12. This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: AV763490.

SGPr413, SEQ ID NO:2, SEQ ID NO:61 is 1125 nucleotides long. The open reading frame starts at position 1 and ends at position 1125, giving an ORF length of 1125 nucleotides. The predicted protein is 374 amino acids long. This sequence codes for a full length protein. It is classified as (superfamily/group/family): Protease, Carboxypeptidase, Zn carboxypeptidase. The cytogenetic position of this gene is 2q35. This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: none.

SGPr404, SEQ ID NO:3, SEQ ID NO:62 is 1590 nucleotides long. The open reading frame starts at position 1 and ends at position 1590, giving an ORF length of 1590 nucleotides. The predicted protein is 529 amino acids long. This sequence codes for a full length protein. It is classified as (superfamily/group/family): Protease, Carboxypeptidase, Zn carboxypeptidase. The cytogenetic position of this gene is 10q26. This nucleotide sequence contains the following single nucleotide polymorphisms (the accession number of SNP is given, with the allele position, followed by the sequence surrounding the SNP within the gene): ss1782198_allelePos=201; agaaggcctaygaagggg. SNP ss1782198 occurs at nucleotide 612 (aa 58) of the ORF (C or T = silent; AA 204 = tyrosine with either nucleotide). This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: AA045748, AA148684, AA047483. The nucleic acid contains short repetitive sequence (the position and sequence of the repeat): 477 ggagctgctgctgctgctggtg 498.

SGPr536_1, SEQ ID NO:4, SEQ ID NO:63 is 1404 nucleotides long. The open reading frame starts at position 1 and ends at position 1404, giving an ORF length of 1404 nucleotides. The predicted protein is 467 amino acids long. This sequence codes for a full length protein. It is classified as (superfamily/group/family):

- 5 Protease, Cysteine, papain. The cytogenetic position of this gene is 1p35. This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: AL542213, AL547246, AL552037. The nucleic acid contains short repetitive sequence (the position and sequence of the repeat): 480 gctgctgctgctgctgggcag
501.

10

SGPr414, SEQ ID NO:5, SEQ ID NO:64 is 10062 nucleotides long. The open reading frame starts at position 1 and ends at position 10062, giving an ORF length of 10062 nucleotides. The predicted protein is 3353 amino acids long. This sequence codes for a full length protein. It is classified as

- 15 (superfamily/group/family): Protease, Cysteine, UCH2b. The cytogenetic position of this gene is 2p14. This nucleotide sequence contains the following single nucleotide polymorphisms (the accession number of SNP is given, followed by the sequence surrounding the SNP within the gene):ss16542_allelePos=101, ctaccctagcygaggaaga. SNP ss16542 occurs at nucleotide 9807 (aa 3269, alanine) of
20 the ORF. The SNP is silent. This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: AU118237, AU131420, AU125083. The nucleic acid contains short repetitive sequence (the position and sequence of the repeat): 2249 accaccaccaccaccaccatcaccaccaccac 2280.

- 25 SGPr430, SEQ ID NO:6, SEQ ID NO:65 is 2943 nucleotides long. The open reading frame starts at position 1 and ends at position 2943, giving an ORF length of 2943 nucleotides. The predicted protein is 980 amino acids long. This sequence codes for a full length protein. It is classified as (superfamily/group/family):
Protease, Cysteine, UCH2b. The cytogenetic position of this gene is 2q37. This

nucleotide sequence contains the following single nucleotide polymorphisms (the accession number of SNP is given, with the allele position, followed by the sequence surrounding the SNP within the gene): ss1534585_allelePos=51, tggaatarctcggac; rs1055687_allelePos=51, tggtaatccgkgtagagg. SNP ss1534585 occurs at nucleotide
 5 538 (aa 180) of the ORF (A or G). The SNP ss1534585 changes amino acid 180. If nucleotide 538 is an adenine, amino acid 180 is a threonine; if it is a guanine, amino acid 180 is an alanine. A second SNP, rs1055687, codes for a G or T at nucleotide 499. rs1055687 changes the amino acid sequence of the gene. Amino acid 167 is a cysteine if nucleotide 499 is a thymidine; amino acid 167 is a glycine if nucleotide
 10 499 is a guanine. This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: W87666, AI076108, BG612864.

SGPr496_1, SEQ ID NO:7, SEQ ID NO:66 is 2862 nucleotides long. The open reading frame starts at position 1 and ends at position 2862, giving an ORF length of
 15 2862 nucleotides. The predicted protein is 953 amino acids long. This sequence codes for a full length protein. It is classified as (superfamily/group/family): Protease, Cysteine, UCH2b. The cytogenetic position of this gene is Xp11.4. This nucleotide sequence contains the following single nucleotide polymorphisms (the accession number of SNP is given, with the allele position, followed by the sequence surrounding the SNP within the gene): ss1029756_allelePos=101,
 20 agagaaataygagggtatt. SNP ss1029756 codes for a C or T at nucleotide 351. Amino acid 117 is a tyrosine with either nucleotide, so the SNP is silent. This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: AW851066, AW851065, AW851076.

25 SGPr495, SEQ ID NO:8, SEQ ID NO:67 is 2352 nucleotides long. The open reading frame starts at position 1 and ends at position 2352, giving an ORF length of 2352 nucleotides. The predicted protein is 783 amino acids long. This sequence codes for a full length protein. It is classified as (superfamily/group/family):

Protease, Cysteine, UCH2b. The cytogenetic position of this gene is 6q16. This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: AL559960, AL530470, AL516184.

- 5 SGPr407, SEQ ID NO:9, SEQ ID NO:68 is 2259 nucleotides long. The open reading frame starts at position 1 and ends at position 2259, giving an ORF length of 2259 nucleotides. The predicted protein is 752 amino acids long. This sequence codes for a full length protein. It is classified as (superfamily/group/family):
Protease, Cysteine, UCH2b. The cytogenetic position of this gene is 2q37. This
0 sequence is represented in the database of public ESTs (dbEST) by the following ESTs: none.

- SGPr453, SEQ ID NO:10, SEQ ID NO:69 is 2139 nucleotides long. The open reading frame starts at position 1 and ends at position 2139, giving an ORF length of
5 2139 nucleotides. The predicted protein is 712 amino acids long. This sequence codes for a full length protein. It is classified as (superfamily/group/family):
Protease, Cysteine, UCH2b. The cytogenetic position of this gene is 12q23. This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: BG722436, AI927881, BG771888. The nucleic acid contains short repetitive
0 sequence (the position and sequence of the repeat): 553 gtagtaaaaagagaagtaaa 572.

- SGPr445, SEQ ID NO:11, SEQ ID NO:70 is 870 nucleotides long. The open reading frame starts at position 1 and ends at position 870, giving an ORF length of 870 nucleotides. The predicted protein is 289 amino acids long. This sequence
5 codes for a full length protein. It is classified as (superfamily/group/family):
Protease, Cysteine, UCH2b. The cytogenetic position of this gene is 6q16. This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: AL559960, AL530470, AL516184.

SGPr401_1, SEQ ID NO:12, SEQ ID NO:71 is 1101 nucleotides long. The open reading frame starts at position 1 and ends at position 1101, giving an ORF length of 1101 nucleotides. The predicted protein is 366 amino acids long. This sequence codes for a full length protein. It is classified as (superfamily/group/family):

- 5 Protease, Cysteine, UCH2b. The cytogenetic position of this gene is 4q11. This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: AU124898, AU134553, AI269069.

10 SGPr408, SEQ ID NO:13, SEQ ID NO:72 is 3864 nucleotides long. The open reading frame starts at position 1 and ends at position 3864, giving an ORF length of 3864 nucleotides. The predicted protein is 1287 amino acids long. This sequence codes for a full length protein. It is classified as (superfamily/group/family):

- 15 Protease, Cysteine, UCH2b. The cytogenetic position of this gene is 11p15. This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: BG741190, BF575498, BG170829.

20 SGPr480, SEQ ID NO:14, SEQ ID NO:73 is 4815 nucleotides long. The open reading frame starts at position 1 and ends at position 4815, giving an ORF length of 4815 nucleotides. The predicted protein is 1604 amino acids long. This sequence codes for a partial protein. It is classified as (superfamily/group/family): Protease, Cysteine, UCH2b. The cytogenetic position of this gene is 17q24. This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: AU131748, AU120381, BG420766.

- 25 SGPr431, SEQ ID NO:15, SEQ ID NO:74 is 3129 nucleotides long. The open reading frame starts at position 1 and ends at position 3129, giving an ORF length of 3129 nucleotides. The predicted protein is 1042 amino acids long. This sequence codes for a full length protein. It is classified as (superfamily/group/family): Protease, Cysteine, UCH2b. The cytogenetic position of this gene is 4q31.3. This

sequence is represented in the database of public ESTs (dbEST) by the following ESTs: BG575871, BG113469, BG112979.

5 SGPr429, SEQ ID NO:16, SEQ ID NO:75 is 3102 nucleotides long. The open reading frame starts at position 1 and ends at position 3102, giving an ORF length of 3102 nucleotides. The predicted protein is 1033 amino acids long. This sequence codes for a full length protein. It is classified as (superfamily/group/family):
Protease, Cysteine, UCH2b. The cytogenetic position of this gene is 1p36.2. This sequence is represented in the database of public ESTs (dbEST) by the following
0 ESTs: AL518266, BG681225, BG217186.

SGPr503, SEQ ID NO:17, SEQ ID NO:76 is 1554 nucleotides long. The open reading frame starts at position 1 and ends at position 1554, giving an ORF length of 1554 nucleotides. The predicted protein is 517 amino acids long. This sequence
5 codes for a full length protein. It is classified as (superfamily/group/family):
Protease, Cysteine, UCH2b. The cytogenetic position of this gene is 12q24.3. This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: BG678894, BG476418, BE264732. The nucleic acid contains short repetitive sequence (the position and sequence of the repeat): 1534 gagtgcaagtctgaagaatg
10 1553.

SGPr427, SEQ ID NO:18, SEQ ID NO:77 is 3372 nucleotides long. The open reading frame starts at position 1 and ends at position 3372, giving an ORF length of 3372 nucleotides. The predicted protein is 1123 amino acids long. This sequence
15 codes for a full length protein. It is classified as (superfamily/group/family):
Protease, Cysteine, UCH2b. The cytogenetic position of this gene is 17p13. This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: BG831111, AW996553, BE614914.

SGPr092, SEQ ID NO:19, SEQ ID NO:78 is 786 nucleotides long. The open reading frame starts at position 1 and ends at position 786, giving an ORF length of 786 nucleotides. The predicted protein is 261 amino acids long. This sequence codes for a full length protein. It is classified as (superfamily/group/family):
5 Protease, Metalloprotease, PepM10. The cytogenetic position of this gene is 11p15. This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: BG189720, AW966183, BG198356.

SGPr359, SEQ ID NO:20, SEQ ID NO:79 is 1452 nucleotides long. The open
10 reading frame starts at position 1 and ends at position 1452, giving an ORF length of 1452 nucleotides. The predicted protein is 483 amino acids long. This sequence codes for a full length protein. It is classified as (superfamily/group/family):
Protease, Metalloprotease, PepM10. The cytogenetic position of this gene is 11q22. This sequence is represented in the database of public ESTs (dbEST) by the
15 following ESTs: BG187290.

SGPr104_1, SEQ ID NO:21, SEQ ID NO:80 is 2298 nucleotides long. The open reading frame starts at position 1 and ends at position 2298, giving an ORF length of 2298 nucleotides. The predicted protein is 765 amino acids long. This sequence
20 codes for a full length protein. It is classified as (superfamily/group/family):
Protease, Metalloprotease, PepM13. The cytogenetic position of this gene is 3q27. This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: BF511209, AW341249, AL119270.

SGPr303, SEQ ID NO:22, SEQ ID NO:81 is 1257 nucleotides long. The open reading frame starts at position 1 and ends at position 1257, giving an ORF length of 1257 nucleotides. The predicted protein is 418 amino acids long. This sequence codes for a full length catalytic domain. It is classified as
25 (superfamily/group/family): Protease, Metalloprotease, PepM2. The cytogenetic

position of this gene is 17q11.1. This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: AU138954, BG251083, AW161660.

5 SGPr402_1, SEQ ID NO:23, SEQ ID NO:82 is 2268 nucleotides long. The open reading frame starts at position 1 and ends at position 2268, giving an ORF length of 2268 nucleotides. The predicted protein is 755 amino acids long. This sequence codes for a full length protein. It is classified as (superfamily/group/family): Protease, Serine, subtilase. The cytogenetic position of this gene is 19q11. This sequence is represented in the database of public ESTs (dbEST) by the following
10 ESTs: AL041695, AA454137, BG719638.

SGPr434, SEQ ID NO:24, SEQ ID NO:83 is 1176 nucleotides long. The open reading frame starts at position 1 and ends at position 1176, giving an ORF length of 1176 nucleotides. The predicted protein is 391 amino acids long. This sequence
15 codes for a full length protein. It is classified as (superfamily/group/family): Protease, Serine, Trypsin. The cytogenetic position of this gene is 3p21. This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: AW137088, BF593342.

20 SGPr446_1, SEQ ID NO:25, SEQ ID NO:84 is 681 nucleotides long. The open reading frame starts at position 1 and ends at position 681, giving an ORF length of 681 nucleotides. The predicted protein is 226 amino acids long. This sequence codes for a full length catalytic domain. It is classified as (superfamily/group/family): Protease, Serine, Trypsin. The cytogenetic position of this gene is 3p21. This
25 sequence is represented in the database of public ESTs (dbEST) by the following ESTs: AW243584. The nucleic acid contains short repetitive sequence (the position and sequence of the repeat): 798 ggtgggcatcatcagctgggg 818.

SGPr447, SEQ ID NO:26, SEQ ID NO:85 is 888 nucleotides long. The open reading frame starts at position 1 and ends at position 888, giving an ORF length of 888 nucleotides. The predicted protein is 295 amino acids long. This sequence codes for a partial protein. It is classified as (superfamily/group/family): Protease, Serine, Trypsin. The cytogenetic position of this gene is 16p13.3. This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: none.

SGPr432_1, SEQ ID NO:27, SEQ ID NO:86 is 1887 nucleotides long. The open reading frame starts at position 1 and ends at position 1887, giving an ORF length of 1887 nucleotides. The predicted protein is 628 amino acids long. This sequence codes for a full length protein. It is classified as (superfamily/group/family): Protease, Serine, Trypsin. The cytogenetic position of this gene is unknown. This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: BE264142, BG474605, BF304202.

SGPr529, SEQ ID NO:28, SEQ ID NO:87 is 831 nucleotides long. The open reading frame starts at position 1 and ends at position 831, giving an ORF length of 831 nucleotides. The predicted protein is 276 amino acids long. This sequence codes for a full length protein. It is classified as (superfamily/group/family): Protease, Serine, Trypsin. The cytogenetic position of this gene is 19q13.4. This nucleotide sequence contains the following single nucleotide polymorphisms (the accession number of SNP is given, with the allele position, followed by the sequence surrounding the SNP within the gene): ss1550333_allelePos=51, taggggatgaycacctgct; ss1546197_allelePos=51, gccggacsactcgc. SNP ss1550333 codes for a C or T at nucleotide 297. Amino acid 99 is an aspartic acid with either nucleotide, so the SNP is silent. There is another SNP, ss1546197, that codes for a C or G at position 336; amino acid 112 is a threonine when either nucleotide is present, and so this SNP is silent. This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: BE898352, BG469321.

SGPr428_1, SEQ ID NO:29, SEQ ID NO:88 is 858 nucleotides long. The open reading frame starts at position 1 and ends at position 858, giving an ORF length of 858 nucleotides. The predicted protein is 285 amino acids long. This sequence
5 codes for a full length catalytic domain. It is classified as (superfamily/group/family): Protease, Serine, Trypsin. The cytogenetic position of this gene is 8p23. This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: none. The nucleic acid contains short repetitive sequence (the position and sequence of the repeat): 473 catgcacctggaaaagctg 491.

10

SGPr425, SEQ ID NO:30, SEQ ID NO:89 is 1242 nucleotides long. The open reading frame starts at position 1 and ends at position 1242, giving an ORF length of 1242 nucleotides. The predicted protein is 413 amino acids long. This sequence codes for a full length protein. It is classified as (superfamily/group/family):
15 Protease, Serine, Trypsin. The cytogenetic position of this gene is 6q14. This nucleotide sequence contains the following single nucleotide polymorphisms (the accession number of SNP is given, with the allele position, followed by the sequence surrounding the SNP within the gene): ss674620_allelePos=201, gagcatctgcVggagagag,. SNP ss674620 codes for a G or A or C at nucleotide 671. If
20 the nucleotide is a guanine, amino acid 224 is an arginine; if it is an adenine, amino acid 224 is a glutamine; if the nucleotide is a cytosine, the amino acid at 224 is a proline. This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: AL551286, AA445948, AA424073. The nucleic acid contains short repetitive sequence (the position and sequence of the repeat): 1111
25 tcagggcaccagtgggtgga 1130.

SGPr548, SEQ ID NO:31, SEQ ID NO:90 is 963 nucleotides long. The open reading frame starts at position 1 and ends at position 963, giving an ORF length of 963 nucleotides. The predicted protein is 320 amino acids long. This sequence

codes for a full length protein. It is classified as (superfamily/group/family):
Protease, Serine, Trypsin. The cytogenetic position of this gene is 19q13.4. This
sequence is represented in the database of public ESTs (dbEST) by the following
ESTs: none.

5

SGPr396, SEQ ID NO:32, SEQ ID NO:91 is 987 nucleotides long. The open
reading frame starts at position 1 and ends at position 987, giving an ORF length of
987 nucleotides. The predicted protein is 328 amino acids long. This sequence
codes for a full length protein. It is classified as (superfamily/group/family):

10 Protease, Serine, Trypsin. The cytogenetic position of this gene is 4q32. This
sequence is represented in the database of public ESTs (dbEST) by the following
ESTs: none.

SGPr426, SEQ ID NO:33, SEQ ID NO:92 is 1278 nucleotides long. The open
15 reading frame starts at position 1 and ends at position 1278, giving an ORF length of
1278 nucleotides. The predicted protein is 425 amino acids long. This sequence
codes for a full length protein. It is classified as (superfamily/group/family):
Protease, Serine, Trypsin. The cytogenetic position of this gene is 4q13. This
sequence is represented in the database of public ESTs (dbEST) by the following
20 ESTs: none.

SGPr552, SEQ ID NO:34, SEQ ID NO:93 is 666 nucleotides long. The open
reading frame starts at position 1 and ends at position 666, giving an ORF length of
666 nucleotides. The predicted protein is 221 amino acids long. This sequence
25 codes for a full length catalytic domain. It is classified as
(superfamily/group/family): Protease, Serine, Trypsin. The cytogenetic position of
this gene is 4q13. This sequence is represented in the database of public ESTs
(dbEST) by the following ESTs: none.

SGPr405, SEQ ID NO:35, SEQ ID NO:94 is 2847 nucleotides long. The open reading frame starts at position 1 and ends at position 2847, giving an ORF length of 2847 nucleotides. The predicted protein is 948 amino acids long. This sequence codes for a full length protein. It is classified as (superfamily/group/family):

- 5 Protease, Serine, Trypsin. The cytogenetic position of this gene is 16p13.3. This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: none.

- 10 SGPr485_1, SEQ ID NO:36, SEQ ID NO:95 is 1059 nucleotides long. The open reading frame starts at position 1 and ends at position 1059, giving an ORF length of 1059 nucleotides. The predicted protein is 352 amino acids long. This sequence codes for a full length protein. It is classified as (superfamily/group/family): Protease, Serine, Trypsin. The cytogenetic position of this gene is 8p23. This nucleotide sequence contains the following single nucleotide polymorphisms (the
- 15 accession number of SNP is given, with the allele position, followed by the sequence surrounding the SNP within the gene):ss1532791_allelePos=51 , tggagakaagaacac. ss1532791 codes for a G or a T at position 834. This polymorphism changes amino acid 278. If the nucleotide at 834 is a guanine, amino acid 278 is a glutamic acid (E); if the nucleotide is a thymine, amino acid 278 is an aspartic acid (D). This
- 20 sequence is represented in the database of public ESTs (dbEST) by the following ESTs: AA781356.

- 25 SGPr534, SEQ ID NO:37, SEQ ID NO:96 is 792 nucleotides long. The open reading frame starts at position 1 and ends at position 792, giving an ORF length of 792 nucleotides. The predicted protein is 263 amino acids long. This sequence codes for a full length protein. It is classified as (superfamily/group/family): Protease, Serine, Trypsin. The cytogenetic position of this gene is 16q23. This nucleotide sequence contains the following six single nucleotide polymorphisms (the

accession number of SNP is given, with the allele position, followed by the sequence surrounding the SNP within the gene):

ss1522946_allelePos=51, gctctaccwccacgccc;

ss1522943_allelePos=51, cgcacctgctcyaccaccac;

5 ss1522933_allelePos=51, ctgccagaaggayggagcctgg;

ss1522931_allelePos=51 total len = 101, gtctgccaraaggacg;

ss1522930_allelePos=51, gggtgactctggmggcccct;

ss1522928_allelePos=51, tgcattgggygactctgg;

10 SNP ss1522946 codes for A or T at position 721. If 721 is adenine, amino acid 241 is threonine (T); if 721 is Thymine, amino acid 241 is serine (S).

SNP ss1522943 codes for C or T at position 717; this SNP is silent (239 = serine).

SNP ss1522933 codes for C or T at 666; this SNP is silent (222 = aspartic acid).

15 SNP ss1522931 codes for A or G at position 660; this SNP is silent (220 = glutamine).

SNP ss1522930 codes for A or C at position 642; this SNP is silent (214 = glycine).

SNP ss1522928 codes for a C or T at position 633; this SNP is silent (211 = glycine).

20 This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: AW583018, AW582942, AW960025. The nucleic acid contains short repetitive sequence (the position and sequence of the repeat): 172 cacttctgcgggggctccctcatc 195.

25 SGPr390, SEQ ID NO:38, SEQ ID NO:97 is 3387 nucleotides long. The open reading frame starts at position 1 and ends at position 3387, giving an ORF length of 3387 nucleotides. The predicted protein is 1128 amino acids long. This sequence codes for a full length protein. It is classified as (superfamily/group/family): Protease, Serine, Trypsin. The cytogenetic position of this gene is 19q11. This

nucleotide sequence contains the following single nucleotide polymorphisms (the accession number of SNP is given, with the allele position, followed by the sequence surrounding the SNP within the gene): ss82431_allelePos=99 , gccgtgarcaccactg; ss1320361_allelePos=225, agcggccascattggcgt. ss82431 codes for an A or G at position 2585. If this nucleotide is an adenine, amino acid 862 is an asparagine (N); if this nucleotide is a guanine, amino acid 862 is a serine. The SNP ss1320361 codes for C or G at position 89. If position 89 is a cytosine, amino acid 30 is a threonine (T). If position 89 is a guanine, amino acid 30 is a serine. This sequence is represented in the database of public ESTs (dbEST) by the following ESTs:

10 C16607.

SGPr521, SEQ ID NO:39, SEQ ID NO:98 is 762 nucleotides long. The open reading frame starts at position 1 and ends at position 762, giving an ORF length of 762 nucleotides. The predicted protein is 253 amino acids long. This sequence codes for a full length protein. It is classified as (superfamily/group/family): Protease, Serine, Trypsin. The cytogenetic position of this gene is 19q13.4. This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: AA542994, BE713379, W58737. The nucleic acid contains short repetitive sequence (the position and sequence of the repeat): 646 caaggtctggtgtcctgggg 665.

15

20 SGPr530_1, SEQ ID NO:40, SEQ ID NO:99 is 816 nucleotides long. The open reading frame starts at position 1 and ends at position 816, giving an ORF length of 816 nucleotides. The predicted protein is 271 amino acids long. This sequence codes for a full length catalytic domain. It is classified as (superfamily/group/family): Protease, Serine, Trypsin. The cytogenetic position of this gene is 9q22. This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: none.

25

SGPr520, SEQ ID NO:41, SEQ ID NO:100 is 1737 nucleotides long. The open reading frame starts at position 1 and ends at position 1737, giving an ORF length of 1737 nucleotides. The predicted protein is 578 amino acids long. This sequence codes for a full length protein. It is classified as (superfamily/group/family):

- 5 Protease, Serine, Trypsin. The cytogenetic position of this gene is 2q37. This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: none.

SGPr455, SEQ ID NO:42, SEQ ID NO:101 is 2913 nucleotides long. The open reading frame starts at position 1 and ends at position 2913, giving an ORF length of 2913 nucleotides. The predicted protein is 970 amino acids long. This sequence codes for a full length protein. It is classified as (superfamily/group/family):

- 10 Protease, Serine, Trypsin. The cytogenetic position of this gene is 12p11.2. This sequence is represented in the database of public ESTs (dbEST) by the following
15 ESTs: AW450155, AW995496.

SGPr507_2, SEQ ID NO:43, SEQ ID NO:102 is 798 nucleotides long. The open reading frame starts at position 1 and ends at position 798, giving an ORF length of 798 nucleotides. The predicted protein is 265 amino acids long. This sequence codes for a full length protein. It is classified as (superfamily/group/family):

- 20 Protease, Serine, Trypsin. The cytogenetic position of this gene is 7q36. This sequence is represented in the database of public ESTs (dbEST) by the following
ESTs: BG217724, BG219738, BG192709.

SGPr559, SEQ ID NO:44, SEQ ID NO:103 is 1365 nucleotides long. The open reading frame starts at position 1 and ends at position 1365, giving an ORF length of 1365 nucleotides. The predicted protein is 454 amino acids long. This sequence codes for a full length protein. It is classified as (superfamily/group/family):

- 25 Protease, Serine, Trypsin. The cytogenetic position of this gene is 21q22. This

sequence is represented in the database of public ESTs (dbEST) by the following ESTs: AI978874, AI469095, BF435670

5 SGPr567_1, SEQ ID NO:45, SEQ ID NO:104 is 1614 nucleotides long. The open reading frame starts at position 1 and ends at position 1614, giving an ORF length of 1614 nucleotides. The predicted protein is 537 amino acids long. This sequence codes for a full length protein. It is classified as (superfamily/group/family): Protease, Serine, Trypsin. The cytogenetic position of this gene is 11q23. This sequence is represented in the database of public ESTs (dbEST) by the following
10 ESTs: BE732381, R78581, AW845106.

SGPr479_1, SEQ ID NO:46, SEQ ID NO:105 is 981 nucleotides long. The open reading frame starts at position 1 and ends at position 981, giving an ORF length of 981 nucleotides. The predicted protein is 326 amino acids long. This sequence
15 codes for a full length protein. It is classified as (superfamily/group/family): Protease, Serine, Trypsin. The cytogenetic position of this gene is 1q42. This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: BG718703, AA401705, AA398170. The nucleic acid contains short repetitive sequence (the position and sequence of the repeat): 780
20 tggaattgtgagctggggccg 800.

SGPr489_1, SEQ ID NO:47, SEQ ID NO:106 is 1671 nucleotides long. The open reading frame starts at position 1 and ends at position 1671, giving an ORF length of 1671 nucleotides. The predicted protein is 556 amino acids long. This sequence
25 codes for a full length catalytic domain. It is classified as (superfamily/group/family): Protease, Serine, Trypsin. The cytogenetic position of this gene is 11p15. This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: AW271430, AW237893.

SGPr465_1, SEQ ID NO:48, SEQ ID NO:107 is 894 nucleotides long. The open reading frame starts at position 1 and ends at position 894, giving an ORF length of 894 nucleotides. The predicted protein is 297 amino acids long. This sequence codes for a full length catalytic domain. It is classified as
5 (superfamily/group/family): Protease, Serine, Trypsin. The cytogenetic position of this gene is unknown. This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: none.

SGPr524_1, SEQ ID NO:49, SEQ ID NO:108 is 2553 nucleotides long. The open
10 reading frame starts at position 1 and ends at position 2553, giving an ORF length of 2553 nucleotides. The predicted protein is 850 amino acids long. This sequence codes for a full length protein. It is classified as (superfamily/group/family): Protease, Serine, Trypsin. The cytogenetic position of this gene is unknown. This nucleotide sequence contains the following single nucleotide polymorphisms (the
15 accession number of SNP is given, with the allele position, followed by the sequence surrounding the SNP within the gene): ss2013558_allelePos=201, gacatggawgtggacgac; ss2014128_allelePos=358, acaatttttygagtgccca.ss2013558 codes for a T of C at position 675; this is a silent SNP. Ss2014128 codes for a C or T at 1369; if the nucleotide is a cytosine, amino acid 457 is an arginine; if the
20 nucleotide 1369 is a thymine, a stop codon is introduced, truncating the protein to 456 amino acids. This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: none. The nucleic acid contains short repetitive sequence (the position and sequence of the repeat): 711
aaaaaaaaagaaaagaaagaaaa 734.

25 SGPr422, SEQ ID NO:50, SEQ ID NO:109 is 1344 nucleotides long. The open reading frame starts at position 1 and ends at position 1344, giving an ORF length of 1344 nucleotides. The predicted protein is 447 amino acids long. This sequence codes for a full length protein. It is classified as (superfamily/group/family):

Protease, Serine, Trypsin. The cytogenetic position of this gene is 4q13. This nucleotide sequence contains the following single nucleotide polymorphisms (the accession number of SNP is given, with the allele position, followed by the sequence surrounding the SNP within the gene): ss1091793_allelePos=101,

5 acatacgcrcrgattgtttg; ss448607_allelePos=101, tgggagcrggtcctgcct. SNP ss1091793 codes for an adenine or guanine at position 956. If 956 is guanine, amino acid 319 is arginine (R); if nucleotide 956 is adenine, amino acid 319 is glutamine (Q). The SNP ss448607 codes for an A or G at position 552. This is silent (amino acid 184 = alanine). This sequence is represented in the database of public ESTs (dbEST) by

10 the following ESTs: none.

SGPr538, SEQ ID NO:51, SEQ ID NO:110 is 1374 nucleotides long. The open reading frame starts at position 1 and ends at position 1374, giving an ORF length of 1374 nucleotides. The predicted protein is 457 amino acids long. This sequence

15 codes for a full length protein. It is classified as (superfamily/group/family): Protease, Serine, Trypsin. The cytogenetic position of this gene is 11q23. This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: AL538140, BF934870. The nucleic acid contains short repetitive sequence (the position and sequence of the repeat): 545 tgggaggcttctctggaggag 564.

20 SGPr527_1, SEQ ID NO:52, SEQ ID NO:111 is 2457 nucleotides long. The open reading frame starts at position 1 and ends at position 2457, giving an ORF length of 2457 nucleotides. The predicted protein is 818 amino acids long. This sequence codes for a full length protein. It is classified as (superfamily/group/family):

25 Protease, Serine, Trypsin. The cytogenetic position of this gene is unknown. This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: AW450407, AI190509, AI864473.

SGPr542, SEQ ID NO:53, SEQ ID NO:112 is 855 nucleotides long. The open reading frame starts at position 1 and ends at position 855, giving an ORF length of 855 nucleotides. The predicted protein is 284 amino acids long. This sequence codes for a full length protein. It is classified as (superfamily/group/family):

- 5 Protease, Serine, Trypsin. The cytogenetic position of this gene is 19q13.1. This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: none.

SGPr551, SEQ ID NO:54, SEQ ID NO:113 is 2409 nucleotides long. The open reading frame starts at position 1 and ends at position 2409, giving an ORF length of 2409 nucleotides. The predicted protein is 802 amino acids long. This sequence codes for a full length protein. It is classified as (superfamily/group/family):

- 15 Protease, Serine, Trypsin. The cytogenetic position of this gene is 22q13. This nucleotide sequence contains the following single nucleotide polymorphisms (the accession number of SNP is given, with the allele position, followed by the sequence surrounding the SNP within the gene): rs881144_allelePos=200, ctgcagccctaygccgagagg; rs855791_allelePos=101, agcgaggycatcgcta. SNP rs881144 codes for a C or T at position 1227; this is a silent SNP (409 = tyrosine). SNP rs855791 codes for C or T at position 2180. If the nucleotide at 2180 is
20 cytosine, the amino acid at 727 is alanine; if the nucleotide is thymine, amino acid 727 is valine. This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: AV693114, N70418, AA609066

- SGPr451, SEQ ID NO:55, SEQ ID NO:114 is 1080 nucleotides long. The open
25 reading frame starts at position 1 and ends at position 1080, giving an ORF length of 1080 nucleotides. The predicted protein is 359 amino acids long. This sequence codes for a full length protein. It is classified as (superfamily/group/family):
Protease, Serine, Trypsin. The cytogenetic position of this gene is 12q23. This nucleotide sequence contains the following single nucleotide polymorphisms (the

accession number of SNP is given, with the allele position, followed by the sequence surrounding the SNP within the gene): ss1881349_allelePos=201, gggcgcgatgcaragg; ss1266911_allelePos=101, ccactgcactaaagacrctag. SNP ss1881349 codes for an A or G at position 217. If the nucleotide at 217 is adenine, amino acid 73 is lysine (K);
5 if the nucleotide is guanine, amino acid 73 is glutamic acid (E). The SNP ss1266911 codes for an A or G at position 412. If 412 is guanine, amino acid 138 is alanine (A); if 412 is adenine, amino acid 138 is threonine (T). This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: BG722131, BG722203,

10 SGPr452_1, SEQ ID NO:56, SEQ ID NO:115 is 867 nucleotides long. The open reading frame starts at position 1 and ends at position 867, giving an ORF length of 867 nucleotides. The predicted protein is 288 amino acids long. This sequence codes for a full length protein. It is classified as (superfamily/group/family):
15 Protease, Serine, Trypsin. The cytogenetic position of this gene is 16p13.3. This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: none

20 SGPr504, SEQ ID NO:57, SEQ ID NO:116 is 135 nucleotides long. The open reading frame starts at position 1 and ends at position 135, giving an ORF length of 135 nucleotides. The predicted protein is 44 amino acids long. This sequence codes for a partial length protein. It is classified as (superfamily/group/family): Protease, Serine, Trypsin. The cytogenetic position of this gene is unknown. This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: none

25 SGPr469, SEQ ID NO:58, SEQ ID NO:117 is 138 nucleotides long. The open reading frame starts at position 1 and ends at position 138, giving an ORF length of 138 nucleotides. The predicted protein is 45 amino acids long. This sequence codes for a partial length protein. It is classified as (superfamily/group/family): Protease,

Serine, Trypsin. The cytogenetic position of this gene is unknown. This sequence is represented in the database of public ESTs (dbEST) by the following ESTs:

AW753029, Z19070. The nucleic acid contains short repetitive sequence (the position and sequence of the repeat): 55 gggattgtgagctggggc 72.

5

SGPr400, SEQ ID NO:59, SEQ ID NO:118 is 930 nucleotides long. The open reading frame starts at position 1 and ends at position 930, giving an ORF length of 930 nucleotides. The predicted protein is 309 amino acids long. This sequence codes for a partial length protein. It is classified as (superfamily/group/family):

10 Protease, Serine, Trypsin. The cytogenetic position of this gene is 4q32. This sequence is represented in the database of public ESTs (dbEST) by the following ESTs: none

15 **DESCRIPTION OF NOVEL PROTEASE POLYPEPTIDES**

SGPr397, SEQ ID NO:1, SEQ ID NO:60 encodes a protein that is 315 amino acids long. It is classified as an Carboxypeptidase protease, of the Zn carboxypeptidase family. The protease domain(s) in this protein match the hidden Markov profile for
20 a Zn carboxypeptidase (PF00246) domain, from amino acid 139 to amino acid 280. The positions within the HMMR profile that match the protein sequence are from profile position 1 to profile position 146. Other domains identified within this protein are: Carboxypeptidase activation peptide (PF02244) from amino acid 41 to 120. The pro-segment moiety (activation peptide) is responsible for modulation of
25 folding and activity of the pro-enzyme (see http://pfam.wustl.edu/cgi-bin/getdesc?name=Propep_M14). The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 3.10E-220; number of identical amino acids = 315; percent identity

= 100%; percent similarity = 100%; the accession number of the most similar entry in NRAA is NP_065094.1; the name or description, and species, of the most similar protein in NRAA is: carboxypeptidase B precursor [Homo sapiens].

- 5 SGPr413, SEQ ID NO:2, SEQ ID NO:61 encodes a protein that is 374 amino acids long. It is classified as an Carboxypeptidase protease, of the Zn carboxypeptidase family. The protease domain(s) in this protein match the hidden Markov profile for a Zn carboxypeptidase (PF00246), from amino acid 50 to amino acid 291. The positions within the HMMR profile that match the protein sequence are from profile position 1 to profile position 248. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 5.90E-93; number of identical amino acids = 146; percent identity = 49%; percent similarity = 68%; the accession number of the most similar entry in NRAA is AAF01344.1; the name or description, and species, of the most similar protein in NRAA is: (AF190274) carboxypeptidase homolog [Bothrops jararaca].

- 20 SGPr404, SEQ ID NO:3, SEQ ID NO:62 encodes a protein that is 529 amino acids long. It is classified as an Carboxypeptidase protease, of the Zn carboxypeptidase family. The protease domain(s) in this protein match the hidden Markov profile for a Zn carboxypeptidase (PF00246), from amino acid 91 to amino acid 466. The positions within the HMMR profile that match the protein sequence are from profile position 1 to profile position 248. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 0; number of identical amino acids = 502; percent identity = 94%; percent similarity = 98%; the accession number of the most similar entry in NRAA is NP_061355.1; the name or description, and species, of the most similar protein in

NRAA is: carboxypeptidase X2 [Mus musculus].

SGPr536_1, SEQ ID NO:4, SEQ ID NO:63 encodes a protein that is 467 amino
5 acids long. It is classified as an Cysteine protease, of the papain family. The
protease domain(s) in this protein match the hidden Markov profile for a papain
(PF00112), from amino acid 203 to amino acid 456. The positions within the
HMMR profile that match the protein sequence are from profile position 1 to profile
position 337. The results of a Smith Waterman search (PAM100, gap open and
10 extend penalties of 12 and 2) of the public database of amino acid sequences
(NRAA) with this protein sequence yielded the following results: Pscore = 1.10E-
276; number of identical amino acids = 467; percent identity = 100%; percent
similarity = 100%; the accession number of the most similar entry in NRAA is
NP_071447.1; the name or description, and species, of the most similar protein in
15 NRAA is: P3ECSL [Homo sapiens].

SGPr414, SEQ ID NO:5, SEQ ID NO:64 encodes a protein that is 3353 amino acids
long. It is classified as a Cysteine protease, of the UCH2b family. The protease
domain(s) in this protein match the hidden Markov profile for a Ubiquitin carboxyl-
20 terminal hydrolase family 2b (PF00443), from amino acid 1951 to amino acid 2045.
The positions within the HMMR profile that match the protein sequence are from
profile position 1 to profile position 72. Other domains identified within this protein
are: Ubiquitin carboxyl-terminal hydrolases family 2 (UCH2b, PF00442) from
amino acid 1701 to 1731. Ubiquitin carboxyl-terminal hydrolases (EC 3.1.2.15)
25 (UCH) (deubiquitinating enzymes) are thiol proteases that recognize and hydrolyze
the peptide bond at the C-terminal glycine of ubiquitin. These enzymes are involved
in the processing of poly-ubiquitin precursors as well as that of ubiquitinated proteins.
The results of a Smith Waterman search (PAM100, gap open and extend penalties of
12 and 2) of the public database of amino acid sequences (NRAA) with this protein

sequence yielded the following results: Pscore = 0; number of identical amino acids = 1259; percent identity = 99%; percent similarity = 100%; the accession number of the most similar entry in NRAA is NP_055524.1; the name or description, and species, of the most similar protein in NRAA is: KIAA0570 gene product [Homo sapiens].

SGPr430, SEQ ID NO:6, SEQ ID NO:65 encodes a protein that is 980 amino acids long. It is classified as a Cysteine protease, of the UCH2b family. The protease domain(s) in this protein match the hidden Markov profile for a Ubiquitin carboxyl-terminal hydrolase family 2b (PF00443), from amino acid 886 to amino acid 951. The positions within the HMMR profile that match the protein sequence are from profile position 1 to profile position 72. Other domains identified within this protein are: UCH2b (PF00442) from amino acids 342 to 373. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 0; number of identical amino acids = 930; percent identity = 99%; percent similarity = 99%; the accession number of the most similar entry in NRAA is BAB13420.1; the name or description, and species, of the most similar protein in NRAA is: (AB046814) KIAA1594 protein [Homo sapiens].

SGPr496_1, SEQ ID NO:7, SEQ ID NO:66 encodes a protein that is 953 amino acids long. It is classified as a Cysteine protease, of the UCH2b family. The protease domain(s) in this protein match the hidden Markov profile for a Ubiquitin carboxyl-terminal hydrolase family 2b (PF00443), from amino acid 875 to amino acid 935. The positions within the HMMR profile that match the protein sequence are from profile position 1 to profile position 72. Other domains identified within this protein are: UCH2b (PF00442) from 593 to 694; and a Zn-finger domain (PF02148), found in ubiquitin-hydrolases, from 465 to 534. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the

public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 2.00E-190; number of identical amino acids = 496; percent identity = 95%; percent similarity = 98%; the accession number of the most similar entry in NRAA is AAF66953.1; the name or description, and species, of the most similar protein in NRAA is: (AF229643) ubiquitin specific protease [Mus musculus].

SGPr495, SEQ ID NO:8, SEQ ID NO:67 encodes a protein that is 783 amino acids long. It is classified as a Cysteine protease, of the UCH2b family. The protease domain(s) in this protein match the hidden Markov profile for a Ubiquitin carboxyl-terminal hydrolase family 2b (PF00443), from amino acid 695 to amino acid 781. The positions within the HMMR profile that match the protein sequence are from profile position 1 to profile position 72. Other domains identified within this protein are: UCH2b (PF00442) from 190 to 221; and Zn-finger in ubiquitin-hydrolases (PF02148) from 465 to 534. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 2.40E-176; number of identical amino acids = 282; percent identity = 100%; percent similarity = 100%; the accession number of the most similar entry in NRAA is AAH05991.1; the name or description, and species, of the most similar protein in NRAA is: (BC005991) Unknown (protein for MGC:14793) [Homo sapiens].

SGPr407, SEQ ID NO:9, SEQ ID NO:68 encodes a protein that is 752 amino acids long. It is classified as a Cysteine protease, of the UCH2b family. The protease domain(s) in this protein match the hidden Markov profile for a Ubiquitin carboxyl-terminal hydrolase family 2b (PF00443), from amino acid 481 to amino acid 491. The positions within the HMMR profile that match the protein sequence are from profile position 80 to profile position 90. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of

amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 2.60E-40; number of identical amino acids = 80; percent identity = 76%; percent similarity = 84%; the accession number of the most similar entry in NRAA is NP_036607.1; the name or description, and species, of the most similar protein in NRAA is: ubiquitin specific protease 23; NEDD8-specific protease [Homo sapiens].

SGPr453, SEQ ID NO:10, SEQ ID NO:69 encodes a protein that is 712 amino acids long. It is classified as a Cysteine protease, of the UCH2b family. The protease domain(s) in this protein match the hidden Markov profile for a Ubiquitin carboxyl-terminal hydrolase family 2b (PF00443), from amino acid 615 to amino acid 677. The positions within the HMMR profile that match the protein sequence are from profile position 1 to profile position 72. Other domains identified within this protein are: UCH2b (PF00442) from 273 to 304; and Zn-finger in ubiquitin-hydrolases (PF02148) from amino acids 29 to 99. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 0; number of identical amino acids = 712; percent identity = 100%; percent similarity = 100%; the accession number of the most similar entry in NRAA is NP_115523.1; the name or description, and species, of the most similar protein in NRAA is: hypothetical protein DKFZp434D0127 [Homo sapiens].

SGPr445, SEQ ID NO:11, SEQ ID NO:70 encodes a protein that is 289 amino acids long. It is classified as a Cysteine protease, of the UCH2b family. The protease domain(s) in this protein match the hidden Markov profile for a Ubiquitin carboxyl-terminal hydrolase family 2b (PF00443), from amino acid 190 to amino acid 221. The positions within the HMMR profile that match the protein sequence are from profile position 1 to profile position 32. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of

amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 3.60E-185; number of identical amino acids = 289; percent identity = 100%; percent similarity = 100%; the accession number of the most similar entry in NRAA is AAH05991.1; the name or description, and species, of the most similar protein in NRAA is: (BC005991) Unknown (protein for MGC:14793) [Homo sapiens].

SGPr401_1, SEQ ID NO:12, SEQ ID NO:71 encodes a protein that is 366 amino acids long. It is classified as a Cysteine protease, of the UCH2b family. The protease domain(s) in this protein match the hidden Markov profile for a Ubiquitin carboxyl-terminal hydrolase family 2b (PF00443), from amino acid 292 to amino acid 364. The positions within the HMMR profile that match the protein sequence are from profile position 1 to profile position 72. Other domains identified within this protein are: UCH2b (PF00442) from amino acids 35 to 66. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 7.30E-254; number of identical amino acids = 366; percent identity = 100%; percent similarity = 100%; the accession number of the most similar entry in NRAA is NP_073743.1; the name or description, and species, of the most similar protein in NRAA is: hypothetical protein FLJ12552 [Homo sapiens].

SGPr408, SEQ ID NO:13, SEQ ID NO:72 encodes a protein that is 1287 amino acids long. It is classified as a Cysteine protease, of the UCH2b family. The protease domain(s) in this protein match the hidden Markov profile for a Ubiquitin carboxyl-terminal hydrolase family 2b (PF00443), from amino acid 395 to amino acid 475. The positions within the HMMR profile that match the protein sequence are from profile position 1 to profile position 72. Other domains identified within this protein are: UCH2b (PF00442) from amino acids 100 to 131. The results of a

Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 0; number of identical amino acids = 1287; percent identity = 100%; percent similarity = 100%; the accession number of the most similar entry in NRAA is BAB55063.1; the name or description, and species, of the most similar protein in NRAA is: (AK027362) unnamed protein product [Homo sapiens].

SGPr480, SEQ ID NO:14, SEQ ID NO:73 encodes a protein that is 1604 amino acids long. It is classified as a Cysteine protease, of the UCH2b family. The protease domain(s) in this protein match the hidden Markov profile for a Ubiquitin carboxyl-terminal hydrolase family 2b (PF00443), from amino acid 1506 to amino acid 1566. The positions within the HMMR profile that match the protein sequence are from profile position 1 to profile position 72. Other domains identified within this protein are: UCH2b (PF00442) from 734 to 765; and two EF hands (PF00036) from 232 to 260, and from 268 to 296. Many calcium-binding proteins belong to the same evolutionary family and share a type of calcium-binding domain known as the EF-hand (see <http://www.expasy.ch/cgi-bin/prosite-search-ac?PDOC00018>). This type of domain consists of a twelve residue loop flanked on both side by a twelve residue alpha-helical domain. In an EF-hand loop the calcium ion is coordinated in a pentagonal bipyramidal configuration. This protein has a putative CAAX motif (CVLQ) which may direct it to the membrane fraction. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 0; number of identical amino acids = 1272; percent identity = 99%; percent similarity = 99%; the accession number of the most similar entry in NRAA is NP_115971.1; the name or description, and species, of the most similar protein in NRAA is: ubiquitin specific protease [Homo sapiens].

- SGPr431, SEQ ID NO:15, SEQ ID NO:74 encodes a protein that is 1042 amino acids long. It is classified as a Cysteine protease, of the UCH2b family. The protease domain(s) in this protein match the hidden Markov profile for a Ubiquitin carboxyl-terminal hydrolase family 2b (PF00443), from amino acid 836 to amino acid 948. The positions within the HMMR profile that match the protein sequence are from profile position 1 to profile position 72. Other domains identified within this protein are: UCH2b (PF00442) from 445 to 476. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 2.40E-251; number of identical amino acids = 397; percent identity = 100%; percent similarity = 100%; the accession number of the most similar entry in NRAA is NP_115946.1; the name or description, and species, of the most similar protein in NRAA is: HP43.8KD protein [Homo sapiens].
- SGPr429, SEQ ID NO:16, SEQ ID NO:75 encodes a protein that is 1033 amino acids long. It is classified as a Cysteine protease, of the UCH2b family. The protease domain(s) in this protein match the hidden Markov profile for a Ubiquitin carboxyl-terminal hydrolase family 2b (PF00443), from amino acid 332 to amino acid 419. The positions within the HMMR profile that match the protein sequence are from profile position 1 to profile position 72. Other domains identified within this protein are: UCH2b (PF00442) from 89 to 120. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 1.50E-250; number of identical amino acids = 368; percent identity = 100%; percent similarity = 100%; the accession number of the most similar entry in NRAA is NP_115612.1; the name or description, and species, of the most similar protein in NRAA is: hypothetical protein FLJ23277 [Homo sapiens]. This protein has a transmembrane domain from amino acid 87 to amino acid 109.

SGPr503, SEQ ID NO:17, SEQ ID NO:76 encodes a protein that is 517 amino acids long. It is classified as a Cysteine protease, of the UCH2b family. The protease domain(s) in this protein match the hidden Markov profile for a Ubiquitin carboxyl-terminal hydrolase family 2b (PF00443), from amino acid 432 to amino acid 501. The positions within the HMMR profile that match the protein sequence are from profile position 1 to profile position 72. Other domains identified within this protein are: UCH2b (PF00442) from 68 to 99. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 0; number of identical amino acids = 508; percent identity = 100%; percent similarity = 100%; the accession number of the most similar entry in NRAA is AAH04868.1; the name or description, and species, of the most similar protein in NRAA is: (BC004868) Unknown (protein for MGC:10702) [Homo sapiens]. This protein has a transmembrane domain from amino acid 35 to amino acid 57.

SGPr427, SEQ ID NO:18, SEQ ID NO:77 encodes a protein that is 1123 amino acids long. It is classified as a Cysteine protease, of the UCH2b family. The protease domain(s) in this protein match the hidden Markov profile for a Ubiquitin carboxyl-terminal hydrolase family 2b (PF00443), from amino acid 648 to amino acid 709. The positions within the HMMR profile that match the protein sequence are from profile position 1 to profile position 72. Other domains identified within this protein are: UCH2b (PF00442) from 101 to 129. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 1.80E-92; number of identical amino acids = 269; percent identity = 36%; percent similarity = 53%; the accession number of the most similar entry in NRAA is AAF47260.1; the name or description, and species, of the

most similar protein in NRAA is: (AE003465) CG3872 gene product [*Drosophila melanogaster*].

SGPr092, SEQ ID NO:19, SEQ ID NO:78 encodes a protein that is 261 amino acids
5 long. It is classified as a Metalloprotease protease, of the PepM10 family. The
protease domain(s) in this protein match the hidden Markov profile for a
Peptidase_M10 (PF00413), from amino acid 75 to amino acid 194. The positions
within the HMMR profile that match the protein sequence are from profile position
49 to profile position 168. Other domains identified within this protein are: ADAM
10 domain, amino acid 207 to 218. The results of a Smith Waterman search (PAM100,
gap open and extend penalties of 12 and 2) of the public database of amino acid
sequences (NRAA) with this protein sequence yielded the following results: Pscore
= 4.70E-171; number of identical amino acids = 261; percent identity = 100%;
percent similarity = 100%; the accession number of the most similar entry in NRAA
15 is XP_011971.1; the name or description, and species, of the most similar protein in
NRAA is: matrix metalloproteinase 26 [*Homo sapiens*].

SGPr359, SEQ ID NO:20, SEQ ID NO:79 encodes a protein that is 483 amino acids
long. It is classified as a Metalloprotease protease, of the PepM10 family. The
20 protease domain(s) in this protein match the hidden Markov profile for a
Peptidase_M10 (PF00413), from amino acid 44 to amino acid 212. The positions
within the HMMR profile that match the protein sequence are from profile position
1 to profile position 168. Other domains identified within this protein are: 3 x
Hemopexin (PF00045) domains from 302 to 403. Hemopexin is a serum
25 glycoprotein that binds heme and transports it to the liver for breakdown and iron
recovery, after which the free hemopexin returns to the circulation. Hemopexin-like
domains have been found in two types of proteins: - in vitronectin, a cell adhesion
and spreading factor found in plasma and tissues and in most members of the matrix
metalloproteinases family (matrixins), including MMP-1, MMP-2, MMP-3, MMP-8,

MMP-9, MMP-10, MMP-11, MMP-12, MMP-13, MMP-14, MMP-15, MMP-16, MMP-17, MMP-18, MMP-19, MMP-20, MMP-24, and MMP-25 (see <http://www.expasy.ch/cgi-bin/prosite-search-ac?PDOC00023>). The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 0; number of identical amino acids = 483; percent identity = 100%; percent similarity = 100%; the accession number of the most similar entry in NRAA is NP_004762.1; the name or description, and species, of the most similar protein in NRAA is: matrix metalloproteinase 20 preproprotein; enamelysin [Homo sapiens]. This protein has a transmembrane domain from amino acid 7 to amino acid 29. This may function as a signal peptide.

SGPr104_1, SEQ ID NO:21, SEQ ID NO:80 encodes a protein that is 765 amino acids long. It is classified as a Metalloprotease protease, of the PepM13 family. The protease domain(s) in this protein match the hidden Markov profile for a Peptidase_M13 (PF01431), from amino acid 561 to amino acid 764. The positions within the HMMR profile that match the protein sequence are from profile position 1 to profile position 222. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 0; number of identical amino acids = 765; percent identity = 100%; percent similarity = 100%; the accession number of the most similar entry in NRAA is NP_055508.1; the name or description, and species, of the most similar protein in NRAA is: KIAA0604 gene product [Homo sapiens]. This protein has a transmembrane domain from amino acid 61 to amino acid 83.

SGPr303, SEQ ID NO:22, SEQ ID NO:81 encodes a protein that is 418 amino acids long. It is classified as a Metalloprotease protease, of the PepM2 family. The protease domain(s) in this protein match the hidden Markov profile for a

- Peptidase_M1 (PF01433), from amino acid 10 to amino acid 397. The positions within the HMMR profile that match the protein sequence are from profile position 1 to profile position 416. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 2.20E-284; number of identical amino acids = 407; percent identity = 97%; percent similarity = 98%; the accession number of the most similar entry in NRAA is CAA10709.1; the name or description, and species, of the most similar protein in NRAA is: (AJ132583) puromycin sensitive aminopeptidase [Homo sapiens].
- SGPr402_1, SEQ ID NO:23, SEQ ID NO:82 encodes a protein that is 755 amino acids long. It is classified as a Serine protease, of the subtilase family. The protease domain(s) in this protein match the hidden Markov profile for a subtilase (PF00082), from amino acid 118 to amino acid 437. The positions within the HMMR profile that match the protein sequence are from profile position 1 to profile position 360. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 0; number of identical amino acids = 513; percent identity = 82%; percent similarity = 89%; the accession number of the most similar entry in NRAA is P29121; the name or description, and species, of the most similar protein in NRAA is: NEUROENDOCRINE CONVERTASE 3 PRECURSOR [Mus musculus].
- SGPr434, SEQ ID NO:24, SEQ ID NO:83 encodes a protein that is 391 amino acids long. It is classified as a Serine protease, of the trypsin family. The protease domain(s) in this protein match the hidden Markov profile for a p20-ICE (PF00656), from amino acid 39 to amino acid 46. The positions within the HMMR profile that match the protein sequence are from profile position 129 to profile position 136. The results of a Smith Waterman search (PAM100, gap open and extend penalties of

12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 6.20E-43; number of identical amino acids = 104; percent identity = 42%; percent similarity = 59%; the accession number of the most similar entry in NRAA is NP_036164.1; the name or
5 description, and species, of the most similar protein in NRAA is: transmembrane tryptase [Mus musculus].

SGPr446_1, SEQ ID NO:25, SEQ ID NO:84 encodes a protein that is 226 amino acids long. It is classified as a Serine protease, of the trypsin family. The protease
10 domain(s) in this protein match the hidden Markov profile for a trypsin (PF00089), from amino acid 13 to amino acid 227. The positions within the HMMR profile that match the protein sequence are from profile position 1 to profile position 242. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein
15 sequence yielded the following results: Pscore = 2.50E-40; number of identical amino acids = 107; percent identity = 45%; percent similarity = 57%; the accession number of the most similar entry in NRAA is NP_038949.1; the name or description, and species, of the most similar protein in NRAA is: distal intestinal serine protease [Mus musculus].

20 SGPr447, SEQ ID NO:26, SEQ ID NO:85 encodes a protein that is 295 amino acids long. It is classified as a Serine protease, of the trypsin family. The protease domain(s) in this protein match the hidden Markov profile for a trypsin (PF00089), from amino acid 33 to amino acid 270. The positions within the HMMR profile that
25 match the protein sequence are from profile position 1 to profile position 259. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 1.00E-97; number of identical amino acids = 167; percent identity = 60%; percent similarity = 77%; the accession

number of the most similar entry in NRAA is BAB30277.1; the name or description, and species, of the most similar protein in NRAA is: (AK016509) putative [Mus musculus].

- 5 SGPr432_1, SEQ ID NO:27, SEQ ID NO:86 encodes a protein that is 628 amino acids long. It is classified as a Serine protease, of the trypsin family. The protease domain(s) in this protein match the hidden Markov profile for a trypsin (PF00089), from amino acid 117 to amino acid 343. The positions within the HMMR profile that match the protein sequence are from profile position 6 to profile position 259.
- 10 The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 3.70E-56; number of identical amino acids = 95; percent identity = 100%; percent similarity = 100%; the accession number of the most similar entry in NRAA is NP_076869.1; the name or
- 15 description, and species, of the most similar protein in NRAA is: hypothetical protein IMAGE3455200 [Homo sapiens]. This protein has two transmembrane domains from amino acid 10 to amino acid 29, and from 82 to 99. The region from amino acid 10 to 29 may function as a signal peptide.
- 20 SGPr529, SEQ ID NO:28, SEQ ID NO:87 encodes a protein that is 276 amino acids long. It is classified as a Serine protease, of the trypsin family. The protease domain(s) in this protein match the hidden Markov profile for a trypsin (PF00089), from amino acid 184 to amino acid 187. The positions within the HMMR profile that match the protein sequence are from profile position 413 to profile position 416.
- 25 The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 1.70E-184; number of identical amino acids = 276; percent identity = 100%; percent similarity = 100%; the accession number of the most similar entry in NRAA is NP_002767.1; the name or

description, and species, of the most similar protein in NRAA is: kallikrein 10; protease, serine-like, 1 [Homo sapiens].

SGPr428_1, SEQ ID NO:29, SEQ ID NO:88 encodes a protein that is 285 amino acids long. It is classified as a Serine protease, of the trypsin family. The protease domain(s) in this protein match the hidden Markov profile for a trypsin (PF00089), from amino acid 24 to amino acid 246. The positions within the HMMR profile that match the protein sequence are from profile position 7 to profile position 259. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 1.90E-58; number of identical amino acids = 92; percent identity = 53%; percent similarity = 73%; the accession number of the most similar entry in NRAA is: BAB24215.1; the name or description, and species, of the most similar protein in NRAA is: (AK005740) putative [Mus musculus]. This protein has a transmembrane domain from amino acid 262 to amino acid 284.

SGPr425, SEQ ID NO:30, SEQ ID NO:89 encodes a protein that is 413 amino acids long. It is classified as a Serine protease, of the trypsin family. The protease domain(s) in this protein match the hidden Markov profile for a trypsin (PF00089), from amino acid 287 to amino acid 306. The positions within the HMMR profile that match the protein sequence are from profile position 387 to profile position 406. This protein has a putative CAAX motif (CAYG) which may direct it to the plasma membrane. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 5.80E-268; number of identical amino acids = 412; percent identity = 99%; percent similarity = 99%; the accession number of the most similar entry in NRAA is: CAC35071.1; the name or

description, and species, of the most similar protein in NRAA is: (AL121939)
dJ223E3.1 (putative secreted protein ZSIG13) [Homo sapiens].

5 SGPr548, SEQ ID NO:31, SEQ ID NO:90 encodes a protein that is 320 amino acids
long. It is classified as a Serine protease, of the trypsin family. The protease
domain(s) in this protein match the hidden Markov profile for a trypsin (PF00089),
from amino acid 86 to amino acid 313. The positions within the HMMR profile that
match the protein sequence are from profile position 1 to profile position 259. The
results of a Smith Waterman search (PAM100, gap open and extend penalties of 12
10 and 2) of the public database of amino acid sequences (NRAA) with this protein
sequence yielded the following results: Pscore = 2.60E-168; number of identical
amino acids = 256; percent identity = 100%; percent similarity = 100%; the
accession number of the most similar entry in NRAA is AAG09469.1; the name or
description, and species, of the most similar protein in NRAA is: (AF242195)
15 KLK15 [Homo sapiens].

SGPr396, SEQ ID NO:32, SEQ ID NO:91 encodes a protein that is 328 amino acids
long. It is classified as a Serine protease, of the trypsin family. The protease
domain(s) in this protein match the hidden Markov profile for a trypsin (PF00089),
20 from amino acid 28 to amino acid 262. The positions within the HMMR profile that
match the protein sequence are from profile position 1 to profile position 259. The
results of a Smith Waterman search (PAM100, gap open and extend penalties of 12
and 2) of the public database of amino acid sequences (NRAA) with this protein
sequence yielded the following results: Pscore = 1.60E-56; number of identical
25 amino acids = 111; percent identity = 44%; percent similarity = 61%; the accession
number of the most similar entry in NRAA is BAA84941.1; the name or
description, and species, of the most similar protein in NRAA is: (AB018694)
epidermis specific serine protease [Xenopus laevis].

SGPr426, SEQ ID NO:33, SEQ ID NO:92 encodes a protein that is 425 amino acids long. It is classified as a Serine protease, of the trypsin family. The protease domain(s) in this protein match the hidden Markov profile for a trypsin (PF00089),
5 from amino acid 194 to amino acid 419. The positions within the HMMR profile that match the protein sequence are from profile position 1 to profile position 259. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 7.70E-93; number of identical
10 amino acids = 181; percent identity = 43%; percent similarity = 61%; the accession number of the most similar entry in NRAA is NP_054777.1; the name or description, and species, of the most similar protein in NRAA is: DESC1 protein [Homo sapiens]. This protein has a transmembrane domain from amino acid 30 to amino acid 52. This region could function as a signal peptide.

15 SGPr552, SEQ ID NO:34, SEQ ID NO:93 encodes a protein that is 221 amino acids long. It is classified as a Serine protease, of the trypsin family. The protease domain(s) in this protein match the hidden Markov profile for a trypsin (PF00089), from amino acid 2 to amino acid 222. The positions within the HMMR profile that
20 match the protein sequence are from profile position 1 to profile position 255. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 1.20E-45; number of identical amino acids = 96; percent identity = 42%; percent similarity = 59%; the accession
25 number of the most similar entry in NRAA is NP_054777.1; the name or description, and species, of the most similar protein in NRAA is: DESC1 protein [Homo sapiens].

SGPr405, SEQ ID NO:35, SEQ ID NO:94 encodes a protein that is 948 amino acids long. It is classified as a Serine protease, of the trypsin family. The protease domain(s) in this protein match the hidden Markov profile for a trypsin (PF00089), from amino acid 218 to amino acid 406. The positions within the HMMR profile that match the protein sequence are from profile position 60 to profile position 259. Other domains identified within this protein are: two additional trypsin domains, from amino acids 419 to 496, and from amino acids 636 to 761. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = $1.10E-30$; number of identical amino acids = 111; percent identity = 54%; percent similarity = 65%; the accession number of the most similar entry in NRAA is P19236; the name or description, and species, of the most similar protein in NRAA is: MASTOCYTOMA PROTEASE PRECURSOR [Canis familiaris].

SGPr485_1, SEQ ID NO:36, SEQ ID NO:95 encodes a protein that is 352 amino acids long. It is classified as a Serine protease, of the trypsin family. The protease domain(s) in this protein match the hidden Markov profile for a trypsin (PF00089), from amino acid 68 to amino acid 295. The positions within the HMMR profile that match the protein sequence are from profile position 1 to profile position 259. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = $7.20E-133$; number of identical amino acids = 223; percent identity = 94%; percent similarity = 96%; the accession number of the most similar entry in NRAA is BAB03569.1; the name or description, and species, of the most similar protein in NRAA is: (AB046651) hypothetical protein [Macaca fascicularis].

- SGPr534, SEQ ID NO:37, SEQ ID NO:96 encodes a protein that is 263 amino acids long. It is classified as a Serine protease, of the trypsin family. The protease domain(s) in this protein match the hidden Markov profile for a trypsin (PF00089), from amino acid 34 to amino acid 256. The positions within the HMMR profile that match the protein sequence are from profile position 1 to profile position 259. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 3.60E-165; number of identical amino acids = 253; percent identity = 96%; percent similarity = 98%; the accession number of the most similar entry in NRAA is NP_001897.1; the name or description, and species, of the most similar protein in NRAA is: chymotrypsinogen B1 [Homo sapiens]. This protein has a transmembrane domain from amino acid 2 to amino acid 24. This region could function as a signal peptide.
- SGPr390, SEQ ID NO:38, SEQ ID NO:97 encodes a protein that is 1128 amino acids long. It is classified as a Serine protease, of the trypsin family. The protease domain(s) in this protein match the hidden Markov profile for a trypsin (PF00089), from amino acid 896 to amino acid 1122. The positions within the HMMR profile that match the protein sequence are from profile position 1 to profile position 259. Other domains identified within this protein are: two trypsin domains, from amino acids 264 to 500, and from amino acids 573 to 800. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 2.60E-53; number of identical amino acids = 135; percent identity = 46%; percent similarity = 59%; the accession number of the most similar entry in NRAA is BAB23684.1; the name or description, and species, of the most similar protein in NRAA is: (AK004939) putative [Mus musculus]. This protein has a transmembrane domain from amino acid 28 to amino acid 50. This region could function as a signal peptide.

SGPr521, SEQ ID NO:39, SEQ ID NO:98 encodes a protein that is 253 amino acids long. It is classified as a Serine protease, of the trypsin family. The protease domain(s) in this protein match the hidden Markov profile for a trypsin (PF00089),
5 from amino acid 30 to amino acid 245. The positions within the HMMR profile that match the protein sequence are from profile position 1 to profile position 259. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 2.30E-155; number of identical
10 amino acids = 253; percent identity = 100%; percent similarity = 100%; the accession number of the most similar entry in NRAA is NP_005037.1; the name or description, and species, of the most similar protein in NRAA is: kallikrein 7 (chymotryptic, stratum corneum); protease, serine, 6 (chymotryptic, stratum corneum) [Homo sapiens].

15 SGPr530_1, SEQ ID NO:40, SEQ ID NO:99 encodes a protein that is 271 amino acids long. It is classified as a Serine protease, of the trypsin family. The protease domain(s) in this protein match the hidden Markov profile for a trypsin (PF00089), from amino acid 14 to amino acid 255. The positions within the HMMR profile that
20 match the protein sequence are from profile position 1 to profile position 259. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 1.10E-95; number of identical amino acids = 142; percent identity = 100%; percent similarity = 100%; the
25 accession number of the most similar entry in NRAA is CAC12709.1; the name or description, and species, of the most similar protein in NRAA is: (AL136097) bA62C3.1 (similar to testicular serine protease) [Homo sapiens].

SGPr520, SEQ ID NO:41, SEQ ID NO:100 encodes a protein that is 578 amino acids long. It is classified as a Serine protease, of the trypsin family. The protease domain(s) in this protein match the hidden Markov profile for a trypsin (PF00089), from amino acid 73 to amino acid 306. The positions within the HMMR profile that match the protein sequence are from profile position 1 to profile position 259. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 1.50E-83; number of identical amino acids = 158; percent identity = 73%; percent similarity = 83%; the accession number of the most similar entry in NRAA is BAB24587.1; the name or description, and species, of the most similar protein in NRAA is: (AK006434) putative [Mus musculus].

SGPr455, SEQ ID NO:42, SEQ ID NO:101 encodes a protein that is 970 amino acids long. It is classified as a Serine protease, of the trypsin family. The protease domain(s) in this protein match the hidden Markov profile for a trypsin (PF00089), from amino acid 433 to amino acid 674. The positions within the HMMR profile that match the protein sequence are from profile position 1 to profile position 259. Other domains identified within this protein are: Trypsin, from amino acid 4 to 156; and three 3 x CUB domains (PF00431) from amino acid 175 to 812. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 5.90E-179; number of identical amino acids = 386; percent identity = 41%; percent similarity = 58%; the accession number of the most similar entry in NRAA is T30337; the name or description, and species, of the most similar protein in NRAA is: polyprotein - African clawed frog.

SGPr507_2, SEQ ID NO:43, SEQ ID NO:102 encodes a protein that is 265 amino acids long. It is classified as a Serine protease, of the trypsin family. The protease

domain(s) in this protein match the hidden Markov profile for a trypsin (PF00089), from amino acid 42 to amino acid 135. The positions within the HMMR profile that match the protein sequence are from profile position 35 to profile position 148. Other domains identified within this protein are: Trypsin domain from amino acid

5 247 to 258. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 2.40E-121; number of identical amino acids = 195; percent identity = 73%; percent similarity = 81%; the accession number of the most similar entry in NRAA is

10 NP_080593.1; the name or description, and species, of the most similar protein in NRAA is: RIKEN cDNA 1700016G05 gene [Mus musculus].

SGPr559, SEQ ID NO:44, SEQ ID NO:103 encodes a protein that is 454 amino acids long. It is classified as a Serine protease, of the trypsin family. The protease

15 domain(s) in this protein match the hidden Markov profile for a trypsin (PF00089), from amino acid 217 to amino acid 444. The positions within the HMMR profile that match the protein sequence are from profile position 1 to profile position 259. Other domains identified within this protein are: Low-density lipoprotein receptor domain class A (PF00057), from amino acid 71 to 109. LDL-receptors the class A

20 domains form the binding site for LDL and calcium. The acidic residues between the fourth and sixth cysteines are important for high-affinity binding of positively charged sequences in LDLR's ligands. The repeat has been shown to consist of a beta-hairpin structure followed by a series of beta turns (see <http://www.expasy.ch/cgi-bin/get-prodoc-entry?PDOC00929>). The results of a

25 Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 1.40E-288; number of identical amino acids = 454; percent identity = 100%; percent similarity = 100%; the accession number of the most similar entry in NRAA is NP_076927.1; the name or description, and

species, of the most similar protein in NRAA is: transmembrane protease, serine 3 [Homo sapiens]. This protein has a transmembrane domain from amino acid 49 to amino acid 71.

- 5 SGPr567_1, SEQ ID NO:45, SEQ ID NO:104 encodes a protein that is 537 amino acids long. It is classified as a Serine protease, of the trypsin family. The protease domain(s) in this protein match the hidden Markov profile for a trypsin (PF00089), from amino acid 296 to amino acid 524. The positions within the HMMR profile that match the protein sequence are from profile position 1 to profile position 259.
- 10 The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 1.70E-135; number of identical amino acids = 534; percent identity = 99%; percent similarity = 99%; the accession number of the most similar entry in NRAA is NP_114435.1; the name or
- 15 description, and species, of the most similar protein in NRAA is: mosaic serine protease [Homo sapiens].

- SGPr479_1, SEQ ID NO:46, SEQ ID NO:105 encodes a protein that is 326 amino acids long. It is classified as a Serine protease, of the trypsin family. The protease
- 20 domain(s) in this protein match the hidden Markov profile for a trypsin (PF00089), from amino acid 60 to amino acid 288. The positions within the HMMR profile that match the protein sequence are from profile position 1 to profile position. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein
- 25 sequence yielded the following results: Pscore = 1.70E-39; number of identical amino acids = 107; percent identity = 42%; percent similarity = 57%; the accession number of the most similar entry in NRAA is NP_114154.1; the name or description, and species, of the most similar protein in NRAA is: marapsin [Homo sapiens].

SGPr489_1, SEQ ID NO:47, SEQ ID NO:106 encodes a protein that is 556 amino acids long. It is classified as a Serine protease, of the trypsin family. The protease domain(s) in this protein match the hidden Markov profile for a trypsin (PF00089),
5 from amino acid 56 to amino acid 257. The positions within the HMMR profile that match the protein sequence are from profile position 1 to profile position 227. Other domains identified within this protein are: 2 x CUB domains (PF00431) from amino acids 304 to 503. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences
10 (NRAA) with this protein sequence yielded the following results: Pscore = 2.70E-90; number of identical amino acids = 194; percent identity = 37%; percent similarity = 54%; the accession number of the most similar entry in NRAA is T30338; the name or description, and species, of the most similar protein in NRAA is: oviductin - [Xenopus laevis].

15

SGPr465_1, SEQ ID NO:48, SEQ ID NO:107 encodes a protein that is 297 amino acids long. It is classified as a Serine protease, of the trypsin family. The protease domain(s) in this protein match the hidden Markov profile for a trypsin (PF00089),
20 from amino acid 2 to amino acid 240. The positions within the HMMR profile that match the protein sequence are from profile position 12 to profile position 259. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 2.70E-76; number of identical amino acids = 144; percent identity = 48%; percent similarity = 66%; the accession
25 number of the most similar entry in NRAA is NP_033381.1; the name or description, and species, of the most similar protein in NRAA is: testicular serine protease 1 [Mus musculus].

SGPr524_1, SEQ ID NO:49, SEQ ID NO:108 encodes a protein that is 850 amino acids long. It is classified as a Serine protease, of the trypsin family. The protease domain(s) in this protein match the hidden Markov profile for a trypsin (PF00089), from amino acid 613 to amino acid 842. The positions within the HMMR profile that match the protein sequence are from profile position 1 to profile position 259. Other domains identified within this protein are: three Low-density lipoprotein receptor domain class A domains (PF00057) from 489 to 603. LDL-receptors the class A domains form the binding site for LDL and calcium. The acidic residues between the fourth and sixth cysteines are important for high-affinity binding of positively charged sequences in LDLR's ligands. The repeat has been shown to consist of a beta-hairpin structure followed by a series of beta turns (see <http://www.expasy.ch/cgi-bin/get-prodoc-entry?PDOC00929>). The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 1.30E-79; number of identical amino acids = 193; percent identity = 41%; percent similarity = 55%; the accession number of the most similar entry in NRAA is BAB23684.1; the name or description, and species, of the most similar protein in NRAA is: (AK004939) putative [Mus musculus]. This protein has a transmembrane domain from amino acid 77 to amino acid 99.

SGPr422, SEQ ID NO:50, SEQ ID NO:109 encodes a protein that is 447 amino acids long. It is classified as a Serine protease, of the trypsin family. The protease domain(s) in this protein match the hidden Markov profile for a trypsin (PF00089), from amino acid 216 to amino acid 441. The positions within the HMMR profile that match the protein sequence are from profile position 1 to profile position 259. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 4.90E-80; number of identical amino acids = 173; percent identity = 39%; percent similarity = 59%; the accession

number of the most similar entry in NRAA is NP_054777.1; the name or description, and species, of the most similar protein in NRAA is: DESC1 protein [Homo sapiens]. This protein has a transmembrane domain from amino acid 32 to amino acid 54. This region could function as a signal peptide.

5

SGPr538, SEQ ID NO:51, SEQ ID NO:110 encodes a protein that is 457 amino acids long. It is classified as a Serine protease, of the trypsin family. The protease domain(s) in this protein match the hidden Markov profile for a trypsin (PF00089), from amino acid 218 to amino acid 448. The positions within the HMMR profile that match the protein sequence are from profile position 1 to profile position 259. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = $9.1e-315$; number of identical amino acids = 457; percent identity = 100%; percent similarity = 100%; the
15 accession number of the most similar entry in NRAA is NP_110397.1; the name or description, and species, of the most similar protein in NRAA is: spinesin [Homo sapiens]. This protein has a transmembrane domain from amino acid 48 to amino acid 70. This region could function as a signal peptide.

20 SGPr527_1, SEQ ID NO:52, SEQ ID NO:111 encodes a protein that is 818 amino acids long. It is classified as a Serine protease, of the trypsin family. The protease domain(s) in this protein match the hidden Markov profile for a trypsin (PF00089), from amino acid 47 to amino acid 286. The positions within the HMMR profile that match the protein sequence are from profile position 1 to profile position 259. Other
25 domains identified within this protein are: two additional trypsin domains, from 323 to 454, and from 564 to 679. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = $1.30E-52$; number of identical amino acids = 114; percent identity = 42%; percent

similarity = 59%; the accession number of the most similar entry in NRAA is AAH03851.1; the name or description, and species, of the most similar protein in NRAA is: (BC003851) Similar to protease, serine, 8 (prostasin) [Mus musculus].

- 5 SGPr542, SEQ ID NO:53, SEQ ID NO:112 encodes a protein that is 284 amino acids long. It is classified as a Serine protease, of the trypsin family. The protease domain(s) in this protein match the hidden Markov profile for a trypsin (PF00089), from amino acid 35 to amino acid 259. The positions within the HMMR profile that match the protein sequence are from profile position 1 to profile position 259. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 2.70E-41; number of identical amino acids = 110; percent identity = 43%; percent similarity = 58%; the accession number of the most similar entry in NRAA is NP_005308.1; the name or description, and species, of the most similar protein in NRAA is: granzyme M precursor; lymphocyte met-ase 1 [Homo sapiens].

- SGPr551, SEQ ID NO:54, SEQ ID NO:113 encodes a protein that is 802 amino acids long. It is classified as a Serine protease, of the trypsin family. The protease domain(s) in this protein match the hidden Markov profile for a trypsin (PF00089), from amino acid 568 to amino acid 797. The positions within the HMMR profile that match the protein sequence are from profile position 1 to profile position 259. Other domains identified within this protein are: three low-density lipoprotein receptor domain class A domains (PF00057) from 447 to 559. LDL-receptors the class A domains form the binding site for LDL and calcium. The acidic residues between the fourth and sixth cysteines are important for high-affinity binding of positively charged sequences in LDLR's ligands. The repeat has been shown to consist of a beta-hairpin structure followed by a series of beta turns (see <http://www.expasy.ch/cgi-bin/get-prodoc-entry?PDOC00929>). The results of a

Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 0; number of identical amino acids = 675; percent identity = 84%; percent similarity = 90%; the accession number of the most similar entry in NRAA is BAB23684.1; the name or description, and species, of the most similar protein in NRAA is: (AK004939) putative [Mus musculus]. This protein has a transmembrane domain from amino acid 44 to amino acid 66. This region could function as a signal peptide.

- 10 SGPr451, SEQ ID NO:55, SEQ ID NO:114 encodes a protein that is 359 amino acids long. It is classified as a Serine protease, of the trypsin family. The protease domain(s) in this protein match the hidden Markov profile for a trypsin (PF00089), from amino acid 89 to amino acid 324. The positions within the HMMR profile that match the protein sequence are from profile position 1 to profile position 259. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 9.90E-41; number of identical amino acids = 101; percent identity = 39%; percent similarity = 59%; the accession number of the most similar entry in NRAA is NP_072152.1; the name or description, and species, of the most similar protein in NRAA is: adrenal secretory serine protease precursor [Rattus norvegicus].

- 25 SGPr452_1, SEQ ID NO:56, SEQ ID NO:115 encodes a protein that is 288 amino acids long. It is classified as a Serine protease, of the trypsin family. The protease domain(s) in this protein match the hidden Markov profile for a trypsin (PF00089), from amino acid 73 to amino acid 280. The positions within the HMMR profile that match the protein sequence are from profile position 1 to profile position 259. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein

sequence yielded the following results: Pscore = 1.40E-81; number of identical amino acids = 142; percent identity = 57%; percent similarity = 72%; the accession number of the most similar entry in NRAA is AAK15264.1; the name or description, and species, of the most similar protein in NRAA is: (AF305425)

5 implantation serine proteinase 2 [Mus musculus].

SGPr504, SEQ ID NO:57, SEQ ID NO:116 encodes a protein that is 44 amino acids long. It is classified as a Serine protease, of the trypsin family. The protease domain(s) in this protein match the hidden Markov profile for a trypsin (PF00089),
10 from amino acid 1 to amino acid 45. The positions within the HMMR profile that match the protein sequence are from profile position 1 to profile position 52. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 2.40E-13; number of identical
15 amino acids = 26; percent identity = 61%; percent similarity = 88%; the accession number of the most similar entry in NRAA is NP_002095.1; the name or description, and species, of the most similar protein in NRAA is: granzyme K precursor; granzyme 3; granzyme K (serine protease, granzyme 3); tryptase II [Homo sapiens].

20 SGPr469, SEQ ID NO:58, SEQ ID NO:117 encodes a protein that is 45 amino acids long. It is classified as a Serine protease, of the trypsin family. The protease domain(s) in this protein match the hidden Markov profile for a trypsin (PF00089), from amino acid 1 to amino acid 46. The positions within the HMMR profile that
25 match the protein sequence are from profile position 210 to profile position 259. The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 2.20E-17; number of identical amino acids = 32; percent identity = 69%; percent similarity = 84%; the accession

number of the most similar entry in NRAA is BAB30277.1; the name or description, and species, of the most similar protein in NRAA is: (AK016509) putative [Mus musculus].

- 5 SGPr400, SEQ ID NO:59, SEQ ID NO:118 encodes a protein that is 309 amino acids long. It is classified as a Serine protease, of the trypsin family. The protease domain(s) in this protein match the hidden Markov profile for a trypsin (PF00089), from amino acid 133 to amino acid 281. The positions within the HMMR profile that match the protein sequence are from profile position 1 to profile position 198.
- 10 The results of a Smith Waterman search (PAM100, gap open and extend penalties of 12 and 2) of the public database of amino acid sequences (NRAA) with this protein sequence yielded the following results: Pscore = 2.30E-16; number of identical amino acids = 72; percent identity = 38%; percent similarity = 48%; the accession number of the most similar entry in NRAA is NP_036164.1; the name or
- 15 description, and species, of the most similar protein in NRAA is: transmembrane tryptase [Mus musculus].

EXAMPLE 2: Expression Analysis of Mammalian Proteases

Materials and Methods

20 Quantitative PCR Analysis

- RNA is isolated from a variety of normal human tissues and cell lines. Single stranded cDNA is synthesized from 10 µg of each RNA as described above using the Superscript Preamplification System (GibcoBRL). These single strand templates are then linearly amplified with a pair of specific primers in a real time
- 25 PCR reaction on a Light Cycler (Roche Molecular Biochemical). Graphical readout can provide quantitative analysis of the relative abundance of the targeted gene in the total RNA preparation.

DNA Array Based Expression Analysis

DNA-free RNA is isolated from a variety of normal human tissues, cryostat sections, and cell lines. Single stranded cDNA is synthesized from 10ug RNA or 1ug mRNA using a modification of the SMART PCR cDNA synthesis technique (Clontech). The procedure can be modified to allow asymmetric labeling of the 5' and 3' ends of each transcript with a unique oligonucleotide sequence. The resulting sscDNAs are then linearly amplified using Advantage long-range PCR (Clontech) on a Light Cycler PCR machine. Reactions are halted when the graphical real-time display demonstrates the products have begun to plateau. The double stranded cDNA products are purified using Millipore DNA purification matrix, dried, resuspended, quantified, and analyzed on an agarose gel. The resulting elements are referred to as "tissue cDNAs".

Tissue cDNAs are spotted onto GAPS coated glass slides (Corning) using a Genetic Microsystems (GMS) arrayer at 500 ng/ul.

Fluorescent labeled oligonucleotides are synthesized to each novel exon, ensuring they contained internal mismatches with the closest known homologue. Typically oligos are 45 nucleotides long, labeled on the 5' end with Cy5.

Exon-specific Cy5-labeled oligos are hybridized to the tissue cDNAs arrayed onto glass slides, and washed using standard buffers and conditions. Hybridizing signals are then quantified using a GMS Scanner.

Alternatively, tissue cDNAs are manually spotted onto Nylon membranes using a 384 pin replicator, and hybridized to ³²P-end labeled oligo probes.

Tissue cDNAs are generated from multiple RNA templates selected to provide information of relevance to the disease areas of interest and to reflect the biological mechanism of action for each protease. These templates include: human tumor cell lines, cryostat sections of primary human tumors and 32 normal human tissues to identify cancer-related genes; sections of normal, Alzheimer's, Parkinson's, and Schizophrenia brain regions for CNS-related genes; normal and diabetic or obese skeletal muscle, adipose, or liver for metabolic-related genes; and

purified hematopoietic cells, and lymphoid tissues for immune-related genes. To characterize gene mechanism of action, tissue cDNAs are generated to reflect angiogenesis (cultured endothelial cells treated with VEGF ligand, anti-angiogenic drugs, or hypoxia), motility (A549 cells stimulated with HGF ligand, orthotopic metastases, primary tumors with matched metastatic tumors), cell cycle (Hela, H1299, and other cell lines synchronized by drug block and harvested at various times in the cell cycle), checkpoint integrity and DNA repair (p53 normal or defective cells treated with γ -radiation, UV, cis-platinum, or oxidative stress), and cell survival (cells induced to differentiate or at various stages of apoptosis).

10

EXAMPLE 3: Isolation of cDNAs Encoding Mammalian Proteases

Materials and Methods

Identification of novel clones

Total RNAs are isolated using the Guanidine Salts/Phenol extraction protocol of Chomczynski and Sacchi (P. Chomczynski and N. Sacchi, *Anal. Biochem.* 162:156 (1987)) from primary human tumors, normal and tumor cell lines, normal human tissues, and sorted human hematopoietic cells. These RNAs are used to generate single-stranded cDNA using the Superscript Preamplification System (GIBCO BRL, Gaithersburg, MD; Gerard, GF *et al.* (1989), FOCUS 11, 66) under conditions recommended by the manufacturer. A typical reaction uses 10 μ g total RNA with 1.5 μ g oligo(dT)₁₂₋₁₈ in a reaction volume of 60 μ L. The product is treated with RNaseH and diluted to 100 μ L with H₂O. For subsequent PCR amplification, 1-4 μ L of this sscDNA is used in each reaction.

Degenerate oligonucleotides are synthesized on an Applied Biosystems 3948 DNA synthesizer using established phosphoramidite chemistry, precipitated with ethanol and used unpurified for PCR. These primers are derived from the sense and antisense strands of conserved motifs within the catalytic domain of several proteases. Degenerate nucleotide residue designations are: N = A, C, G, or T; R =

A or G; Y = C or T; H = A, C or T not G; D = A, G or T not C; S = C or G; and W = A or T.

PCR reactions are performed using degenerate primers applied to multiple single-stranded cDNAs. The primers are added at a final concentration of 5 μ M each to a mixture containing 10 mM TrisHCl, pH 8.3, 50 mM KCl, 1.5 mM MgCl₂, 200 μ M each deoxynucleoside triphosphate, 0.001% gelatin, 1.5 U AmpliTaq DNA Polymerase (Perkin-Elmer/Cetus), and 1-4 μ L cDNA. Following 3 min denaturation at 95 °C, the cycling conditions are 94 °C for 30 s, 50 °C for 1 min, and 72 °C for 1 min 45 s for 35 cycles. PCR fragments migrating between 300-350 bp are isolated from 2% agarose gels using the GeneClean Kit (Bio101), and T-A cloned into the pCRII vector (Invitrogen Corp. U.S.A.) according to the manufacturer's protocol.

Colonies are selected for mini plasmid DNA-preparations using Qiagen columns and the plasmid DNA is sequenced using a cycle sequencing dye-terminator kit with AmpliTaq DNA Polymerase, FS (ABI, Foster City, CA). Sequencing reaction products are run on an ABI Prism 377 DNA Sequencer, and analyzed using the BLAST alignment algorithm (Altschul, S.F. *et al.*, *J.Mol.Biol.* 215: 403-10).

Additional PCR strategies are employed to connect various PCR fragments or ESTs using exact or near exact oligonucleotide primers. PCR conditions are as described above except the annealing temperatures are calculated for each oligo pair using the formula: $T_m = 4(G+C)+2(A+T)$.

Isolation of cDNA clones:

Human cDNA libraries are probed with PCR or EST fragments corresponding to protease-related genes. Probes are ³²P-labeled by random priming and used at 2x10⁶ cpm/mL following standard techniques for library screening. Pre-hybridization (3 h) and hybridization (overnight) are conducted at 42 °C in 5X SSC, 5X Denhart's solution, 2.5% dextran sulfate, 50 mM Na₂PO₄/NaHPO₄, pH 7.0, 50% formamide with 100 mg/mL denatured salmon sperm DNA. Stringent washes are

performed at 65 °C in 0.1X SSC and 0.1% SDS. DNA sequencing is carried out on both strands using a cycle sequencing dye-terminator kit with AmpliTaq DNA Polymerase, FS (ABI, Foster City, CA). Sequencing reaction products are run on an ABI Prism 377 DNA Sequencer.

5

EXAMPLE 4: Expression Analysis of Mammalian Proteases

Materials and Methods

Northern blot analysis

Northern blots are prepared by running 10 µg total RNA isolated from 60 human tumor cell lines (such as HOP-92, EKVX, NCI-H23, NCI-H226, NCI-H322M, NCI-H460, NCI-H522, A549, HOP-62, OVCAR-3, OVCAR-4, OVCAR-5, OVCAR-8, IGROV1, SK-OV-3, SNB-19, SNB-75, U251, SF-268, SF-295, SF-539, CCRF-CEM, K-562, MOLT-4, HL-60, RPMI 8226, SR, DU-145, PC-3, HT-29, HCC-2998, HCT-116, SW620, Colo 205, HTC15, KM-12, UO-31, SN12C, A498, CaKi1, RXF-393, ACHN, 786-0, TK-10, LOX IMVI, Malme-3M, SK-MEL-2, SK-MEL-5, SK-MEL-28, UACC-62, UACC-257, M14, MCF-7, MCF-7/ADR RES, Hs578T, MDA-MB-231, MDA-MB-435, MDA-N, BT-549, T47D), from human adult tissues (such as thymus, lung, duodenum, colon, testis, brain, cerebellum, cortex, salivary gland, liver, pancreas, kidney, spleen, stomach, uterus, prostate, skeletal muscle, placenta, mammary gland, bladder, lymph node, adipose tissue), and 2 human fetal normal tissues (fetal liver, fetal brain), on a denaturing formaldehyde 1.2% agarose gel and transferring to nylon membranes.

Filters are hybridized with random primed [α - 32 P]dCTP-labeled probes synthesized from the inserts of several of the protease genes. Hybridization is performed at 42 °C overnight in 6X SSC, 0.1% SDS, 1X Denhardt's solution, 100 µg/mL denatured herring sperm DNA with $1-2 \times 10^6$ cpm/mL of 32 P-labeled DNA probes. The filters are washed in 0.1X SSC/0.1% SDS, 65 °C, and exposed on a Molecular Dynamics phosphorimager.

Quantitative PCR analysis

RNA is isolated from a variety of normal human tissues and cell lines.

Single stranded cDNA is synthesized from 10 µg of each RNA as described above

- 5 using the Superscript Preamplification System (GibcoBRL). These single strand templates are then used in a 25 cycle PCR reaction with primers specific to each clone. Reaction products are electrophoresed on 2% agarose gels, stained with ethidium bromide and photographed on a UV light box. The relative intensity of the STK-specific bands were estimated for each sample.

10

DNA Array Based Expression Analysis

Plasmid DNA array blots are prepared by loading 0.5 µg denatured plasmid

for each protease on a nylon membrane. The [$\gamma^{32}\text{P}$]dCTP labeled single stranded

DNA probes are synthesized from the total RNA isolated from several human

- 15 immune tissue sources or tumor cells (such as thymus, dendrocytes, mast cells, monocytes, B cells (primary, Jurkat, RPMI8226, SR), T cells (CD8/CD4+, TH1, TH2, CEM, MOLT4), K562 (megakaryocytes). Hybridization is performed at 42 °C for 16 hours in 6X SSC, 0.1% SDS, 1X Denhardt's solution, 100 µg/mL denatured herring sperm DNA with 10^6 cpm/mL of [$\gamma^{32}\text{P}$]dCTP labeled single stranded probe.

- 20 The filters are washed in 0.1X SSC/0.1% SDS, 65 °C, and exposed for quantitative analysis on a Molecular Dynamics phosphorimager.

EXAMPLE 5: Protease Gene Expression

Vector Construction

- 25 Materials and Methods

Expression Vector Construction

Expression constructs are generated for some of the human cDNAs including: a) full-length clones in a pCDNA expression vector; and b) a GST-fusion

construct containing the catalytic domain of the novel protease fused to the C-terminal end of a GST expression cassette; and c) a full-length clone containing a mutation within the predicted polypeptide cleaving site within the protease domain, inserted in the pCDNA vector.

- 5 These mutants of the protease might function as dominant negative constructs, and will be used to elucidate the function of these novel proteases.

EXAMPLE 6: Generation of Specific Immunoreagents to Proteases

Materials and Methods

- 10 Specific immunoreagents are raised in rabbits against KLH- or MAP-conjugated synthetic peptides corresponding to isolated protease polypeptides. C-terminal peptides were conjugated to KLH with glutaraldehyde, leaving a free C-terminus. Internal peptides were MAP-conjugated with a blocked N-terminus. Additional immunoreagents can also be generated by immunizing rabbits with the
- 15 bacterially expressed GST-fusion proteins containing the cytoplasmic domains of each novel PTK or STK.

 The various immune sera are first tested for reactivity and selectivity to recombinant protein, prior to testing for endogenous sources.

Western blots

- 20 Proteins in SDS PAGE are transferred to immobilon membrane. The washing buffer is PBST (standard phosphate-buffered saline pH 7.4 + 0.1% Triton X-100). Blocking and antibody incubation buffer is PBST +5% milk. Antibody dilutions are varied from 1:1000 to 1:2000.

25

EXAMPLE 7: Recombinant Expression and Biological Assays for Proteases

Materials and Methods

Transient Expression of Proteases in Mammalian Cells

The pcDNA expression plasmids (10 µg DNA/100 mm plate) containing the protease constructs are introduced into 293 cells with lipofectamine (Gibco BRL). After 72 hours, the cells are harvested in 0.5 mL solubilization buffer (20 mM HEPES, pH 7.35, 150 mM NaCl, 10% glycerol, 1% Triton X-100, 1.5 mM MgCl₂, 1 mM EGTA, 2 mM phenylmethylsulfonyl fluoride, 1 µg/mL aprotinin). Sample aliquots are resolved by SDS polyacrylamide gel electrophoresis (PAGE) on 6% acrylamide/0.5% bis-acrylamide gels and electrophoretically transferred to nitrocellulose. Non-specific binding is blocked by preincubating blots in Blotto (phosphate buffered saline containing 5% w/v non-fat dried milk and 0.2% v/v nonidet P-40 (Sigma)), and recombinant protein is detected using the various anti-peptide or anti-GST-fusion specific antisera.

In Vitro Protease Assays

In vitro Protease Assay Using Fluorogenic Peptides

Assays are carried out using a spectrofluorometer, such as Perkin-Elmer 204S. The standard reaction mixtures (100 µl) contains 200 mM Tris-HCl, pH8.5, and 200 µM fluorogenic peptide substrate. After enzyme addition, reaction mixtures are incubated at 37 °C for 30 min and terminated by addition of 1.9 ml of 125 mM ZnSO₄ (Brenner, C., and Fuller, R. S., 1992, *Proc. Natl. Acad. Sci. U. S. A.* 89:922-926). The precipitate is removed by centrifugation for 1 min in a microcentrifuge (15,000 × g), and the rate of product (7-amino-4-methyl-coumarin) released into the supernatant solution is determined fluorometrically [(excitation) = 385 nm, (emission) = 465 nm]. Examples of substrates used in the literature include: Boc-Gly-Arg-Arg-4-methylcoumaryl-7-amide (MCA), Boc-Gln-Arg-Arg-MCA, Z-Arg-Arg-MCA, and pGlu-Arg-Thr-Lys-Arg-MCA. Stock solutions (100 mM) are prepared by dissolving peptides in dimethyl sulfoxide that are then diluted in water to 1 mM working stock before use. (Details of this assay can be found in: R. Yosuf, *et al. J. Biol. Chem.*, Vol. 275, Issue 14, 9963-9969, April 7, 2000 which is

incorporated herein by reference in its entirety including any figures, tables, or drawings.)

Protease assay in intact cells using fluorogenic peptides-

5 Calpain activity is measured by the rate of generation of the fluorescent product, AMC, from intracellular thiol-conjugated Boc-Leu-Met-CMAC (Rosser, B. G., Powers, S. P., and Gores, G. J. (1993) *J. Biol. Chem.* 268, 23593-23600). Cells are dispersed, grown on glass coverslips, continuously superfused with physiologic saline solution at 37 °C, and sequentially imaged with a quantitative fluorescence
10 imaging system. At $t = 0$, Boc-Leu-Met-CMAC (10 μ M, Molecular Probes) is introduced into the superfusion solution, and mean fluorescence intensity (excitation 350 nm, emission 470 nm) of individual cells is measured at 60-s intervals. At 10 min, TNF-alpha (30 ng/ml) is added to the superfusion solution with 10 μ M Boc-Leu-Met-CMAC. The slope of the fluorescence change with respect to time
15 represents the intracellular calpain activity (Rosser, *et al.*, 1993, *J. Biol. Chem.* 268:23593-23600). For calpain assays in whole cell populations, suspension cultures of cells are loaded with 10 μ M Boc-Leu-Met-CMAC, and changes in intracellular fluorescence are measured prior to and after TNF-alpha addition at 37 °C using a FACS Vantage system. Cellular fluorescence of AMC is measured using
20 a 360-nm excitation filter and a 405-nm long-pass emission filter. (Details of this assay can be found in: Han, *et al.*, 1999, *J Biol Chem*, 274:787-794 which is incorporated herein by reference in its entirety including any figures, tables, or drawings)

25 Protease assay using chromogenic substrates

The proteolytic activity of enzymes is measured using a commercially available assay system (Athena Environmental Sciences, Inc.). The assay employs a universal substrate of a dye-protein conjugate cross linked to a matrix. Protease activity is determined spectrophotometrically by measuring the absorbance of the

dye released from the matrix to the supernatant. Reaction vials containing the enzyme and substrate are incubated for 3 h at 37 °C. The activity is measured at different incubation times, and reactions are terminated by adding 500 µl of 0.2 N NaOH to each vial. The absorbance of the supernatant in each reaction vial is measured at 450 nm. The proteolytic activity is monitored using 10 µl (approximately 10 µg) of purified protein incubated with 5 µg of -casein (Sigma) in 50 mM Tris-HCl (pH 7.5) for 30 min, 1 h or 2 h at 37 °C. The reaction products are resolved by SDS-polyacrylamide gel electrophoresis and proteins visualized by staining with Coomassie Blue (Details of this assay can be found in: Faccio, *et al.*, 2000, *J Biol. Chem.*, 275:2581-2588 which is incorporated herein by reference in its entirety including any figures, tables, or drawings).

Protease assay using radiolabeled substrate bound to membranes-

Unlabeled protease is mixed with radiolabeled substrate-containing membranes in buffer (100 mM HEPES, 100 mM NaCl, 125 µM magnesium acetate, 125 µM zinc acetate, pH 7.5) and incubated at 30 °C. Typically, each reaction had a final volume of 80-100 µl. Each reaction is normalized to the same final concentration of lysis buffer components (25 mM Tris, 0.1 M sorbitol, 0.5 mM EDTA, 0.01% NaN₃, pH 7.5) because the amount of membranes added to each reaction is varied. To examine metal ion specificity, reactions are assembled without substrate and pretreated with 1.125 mM 1,10-orthophenanthroline for 20 min on ice. Subsequently, metal ions and substrate-containing membranes are added, and reactions are initiated by incubation at 30 °C; the additions result in dilution of the 1,10-orthophenanthroline to a final concentration of 1 mM. The metal ions are added in the form of acetate salts from 25-100 mM stock solutions (Zn²⁺, Mg²⁺, Cu²⁺, Co²⁺, or Ca²⁺) that are first acidified with 2 mM concentrated HCl and then neutralized with 1 mM HEPES, pH 7.5; this step is necessary to achieve full solubilization of zinc acetate. For analysis by immunoprecipitation, samples are diluted 10-20× with immunoprecipitation buffer (Berkower, C., and Michaelis, S.

(1991) *EMBO J.* 10:3777-3785) containing 0.1% SDS, cleared of insoluble material (13,000 × g for 5-10 min at 4 °C), and immunoprecipitated with substrate-specific antibody. Alternatively, samples are solubilized by SDS (final concentration, 0.5%), boiled for 3 min, and directly immunoprecipitated after dilution with

5 immunoprecipitation buffer. Immunoprecipitates are subjected to SDS-polyacrylamide gel electrophoresis as described, fixed for 7 min with 20% trichloroacetic acid, dried, and exposed to a PhosphorImager screen for detection and quantitation (Molecular Dynamics, Sunnyvale, CA). All of the above reagents can be purchased from Sigma. (Details of this assay can be found in: Schmidt, *et*

10 *al.*, 2000, *J Biol Chem*, 275:6227-6233 which is incorporated herein by reference in its entirety including any figures, tables, or drawings). Variation of this assay to apply to substrate not bound to membrane is straightforward.

A comprehensive discussion of various protease assays can be found in: The Handbook of Proteolytic Enzymes by Alan J. Barrett (Editor), Neil D. Rawlings (Editor), J. Fred Woessner (Editor) (February 1998) Academic Press, San Diego;

15 ISBN: 0-12-079370-9 (which is incorporated herein by reference in its entirety including any figures, tables, or drawings).

Similar assays are performed on bacterially expressed GST-fusion constructs

20 of the proteases.

EXAMPLE 8a: Chromosomal Localization of Proteases

Materials And Methods

Several sources were used to find information about the chromosomal

25 localization of each of the genes described in this patent. First, the Celera Browser was used to map the genes. Alternatively, the accession number of a genomic contig (identified by BLAST against NRNA) was used to query the Entrez Genome Browser (<http://www.ncbi.nlm.nih.gov/PMGifs/Genomes/MapViewHelp.html>), and the cytogenetic localization was read from the NCBI data. References for

association of the mapped sites with chromosomal amplifications found in human cancer can be found in: Knuutila, et al., Am J Pathol, 1998, 152:1107-1123.

Information on mapped positions was also obtained by searching published literature (at NCBI, <http://www.ncbi.nlm.nih.gov/entrez/query.fcgi>) for documented

5 association of the mapped position with human disease.

Results

The chromosomal regions for mapped genes are listed in Table 2.

10 The following section describes various diseases that map to chromosomal locations established for proteases included in this patent application. The protease polynucleotides of the present invention can be used to identify individuals who have, or are at risk for developing, relevant diseases. As discussed elsewhere in this application, the polypeptides and polynucleotides of the present invention are useful in identifying compounds that modulate protease activity, and in turn ameliorate
15 various diseases.

SGPr397, SEQID_1, maps to human chromosomal position 8q12. Chromosomal aberrations in this region are associated with breast cancer: Rummukainen J, et al., *Cancer Genet Cytogenet.* 2001 Apr 1;126(1):1-7.

20

SGPr413, SEQ ID NO:2, maps to human chromosomal position 2q35. This region is highly implicated in osteoarthritis (Loughlin J, et al., Linkage analysis of chromosome 2q in osteoarthritis. *Rheumatology.* 2000 Apr;39(4): 377-81).

25 SGPr404, SEQ ID NO:3, maps to human chromosomal position 10q26. Genomic amplification of this region has been associated with the following cancers (Knuutila): Malignant fibrous histiocytoma of soft tissue.

SGPr536_1, SEQ ID NO:4, maps to human chromosomal position 1p35.

SGPr414, SEQ ID NO:5, maps to human chromosomal position 2p14.

5 SGPr430, SEQ ID NO:6, maps to human chromosomal position 2q37. This region is highly implicated in osteoarthritis (Loughlin J, et al. ,Linkage analysis of chromosome 2q in osteoarthritis. *Rheumatology*. 2000 Apr;39(4): 377-81).

SGPr496_1, SEQ ID NO:7, maps to human chromosomal position Xp11. 4. Genomic amplification of this region has been associated with the following cancers
10 (Knuutila): small cell lung cancer and prostate cancer.

SGPr495, SEQ ID NO:8, maps to human chromosomal position 6q16.

SGPr407, SEQ ID NO:9, maps to human chromosomal position 2q37. This region is
15 highly implicated in osteoarthritis (Loughlin J, et al. ,Linkage analysis of chromosome 2q in osteoarthritis. *Rheumatology*. 2000 Apr;39(4): 377-81).

SGPr453, SEQ ID NO:10, maps to human chromosomal position 12q23.

20 SGPr445, SEQ ID NO:11, maps to human chromosomal position 6q16.

SGPr401_1, SEQ ID NO:12, maps to human chromosomal position 4q11. Genomic amplification of this region has been associated with the following cancers (Knuutila): Follicular carcinoma.

25

SGPr408, SEQ ID NO:13, maps to human chromosomal position 11p15.

- SGPr480, SEQ ID NO:14, maps to human chromosomal position 17q24. Genomic amplification of this region has been associated with the following cancers (Knuutila): Non-small cell lung cancer, and testicular cancer.
- 5 SGPr431, SEQ ID NO:15, maps to human chromosomal position 4q31. 3. Genomic amplification of this region has been associated with the following cancers (Knuutila): Osteosarcoma.
- 10 SGPr429, SEQ ID NO:16, maps to human chromosomal position 1p36. 2. Genomic amplification of this region has been associated with the following cancers (Knuutila): alveolar cancer.
- 15 SGPr503, SEQ ID NO:17, maps to human chromosomal position 12q24. 3. Genomic amplification of this region has been associated with the following cancers (Knuutila): Non-small cell lung cancer.
- 20 SGPr427, SEQ ID NO:18, maps to human chromosomal position 17p13.
- SGPr092, SEQ ID NO:19, maps to human chromosomal position 11p15.
- 20 SGPr359, SEQ ID NO:20, maps to human chromosomal position 11q22. Genomic amplification of this region has been associated with the following cancers (Knuutila): Uterine cervix cancer.
- 25 SGPr104_1, SEQ ID NO:21, maps to human chromosomal position 3q27. Genomic amplification of this region has been associated with the following cancers (Knuutila): Squamous cell carcinomas of the head and neck; Malignant fibrous histiocytoma of soft tissue.

SGPr303, SEQ ID NO:22, maps to human chromosomal position 17q11. 1.

Genomic amplification of this region has been associated with the following cancers (Knuutila): Breast carcinoma and Hepatocellular carcinoma.

5 SGPr402_1, SEQ ID NO:23, maps to human chromosomal position 19q11.

Genomic amplification of this region has been associated with the following cancers (Knuutila): Leiomyosarcoma.

SGPr434, SEQ ID NO:24, maps to human chromosomal position 3p21. Genomic
10 amplification of this region has been associated with the following cancers (Knuutila): Bladder carcinoma.

SGPr446_1, SEQ ID NO:25, maps to human chromosomal position 3p21. Genomic
amplification of this region has been associated with the following cancers
15 (Knuutila): Bladder carcinoma.

SGPr447, SEQ ID NO:26, maps to human chromosomal position 16p13. 3.

SGPr432_1, SEQ ID NO:27, has not been assigned a chromosomal location.
20

SGPr529, SEQ ID NO:28, maps to human chromosomal position 19q13. 4.
Genomic amplification of this region has been associated with the following cancers (Knuutila): Breast carcinoma.

25 SGPr428_1, SEQ ID NO:29, maps to human chromosomal position 8p23.

SGPr425, SEQ ID NO:30, maps to human chromosomal position 6q14.

SGPr548, SEQ ID NO:31, maps to human chromosomal position 19q13. 4.

Genomic amplification of this region has been associated with the following cancers (Knuutila): Breast carcinoma.

- 5 SGPr396, SEQ ID NO:32, maps to human chromosomal position 4q32. Genomic amplification of this region has been associated with the following cancers (Knuutila): Non-small cell lung cancer.

- 10 SGPr426, SEQ ID NO:33, maps to human chromosomal position 4q13. Genomic amplification of this region has been associated with the following cancers (Knuutila): Non-small cell lung cancer.

- 15 SGPr552, SEQ ID NO:34, maps to human chromosomal position 4q13. Genomic amplification of this region has been associated with the following cancers (Knuutila): Non-small cell lung cancer.

- SGPr405, SEQ ID NO:35, maps to human chromosomal position 16p13. 3.

- 20 SGPr485_1, SEQ ID NO:36, maps to human chromosomal position 8p23.

- SGPr534, SEQ ID NO:37, maps to human chromosomal position 16q23. Genomic amplification of this region has been associated with the following cancers (Knuutila): Diffuse large cell lymphoma of stomach.

- 25 SGPr390, SEQ ID NO:38, maps to human chromosomal position 19q11. Genomic amplification of this region has been associated with the following cancers (Knuutila): Leiomyosarcoma.

SGPr521, SEQ ID NO:39, maps to human chromosomal position 19q13. 4.

Genomic amplification of this region has been associated with the following cancers (Knuutila): Breast carcinoma.

- 5 SGPr530_1, SEQ ID NO:40, maps to human chromosomal position 9q22. Genomic amplification of this region has been associated with the following cancers (Knuutila): Non-small cell lung cancer.

- 10 SGPr520, SEQ ID NO:41, maps to human chromosomal position 2q37. This region is highly implicated in osteoarthritis (Loughlin J, et al. ,Linkage analysis of chromosome 2q in osteoarthritis. *Rheumatology*. 2000 Apr;39(4): 377-81).

- 15 SGPr455, SEQ ID NO:42, maps to human chromosomal position 12p11. 2. Genomic amplification of this region has been associated with the following cancers (Knuutila): ovarian germ cell tumor, testicular cancer and non-small cell lung cancer.

- 20 SGPr507_2, SEQ ID NO:43, maps to human chromosomal position 7q36. Genomic amplification of this region has been associated with the following cancers (Knuutila): Ovarian cancer.

SGPr559, SEQ ID NO:44, maps to human chromosomal position 21q22.

- 25 SGPr567_1, SEQ ID NO:45, maps to human chromosomal position 11q23. Genomic amplification of this region has been associated with the following cancers (Knuutila): Pleural mesothelioma.

SGPr479_1, SEQ ID NO:46, maps to human chromosomal position 1q42.

SGPr489_1, SEQ ID NO:47, maps to human chromosomal position 11p15.

SGPr465_1, SEQ ID NO:48, has not been assigned a chromosomal location.

- 5 SGPr524_1, SEQ ID NO:49, has not been assigned a chromosomal location.

SGPr422, SEQ ID NO:50, maps to human chromosomal position 4q13. Genomic amplification of this region has been associated with the following cancers (Knuutila): Non-small cell lung cancer.

- 10 SGPr538, SEQ ID NO:51, maps to human chromosomal position 11q23. Genomic

amplification of this region has been associated with the following cancers (Knuutila): Pleural mesothelioma.

- 15 SGPr527_1, SEQ ID NO:52, has not been assigned a chromosomal location.

SGPr542, SEQ ID NO:53, maps to human chromosomal position 19q13. 1. Genomic amplification of this region has been associated with the following cancers (Knuutila): Small cell lung cancer (highly associated, with 10 of 35 patients tested showing amplification).

20

SGPr551, SEQ ID NO:54, maps to human chromosomal position 22q13. Genomic amplification of this region has been associated with the following cancers (Knuutila): Osteosarcoma.

25

SGPr451, SEQ ID NO:55, maps to human chromosomal position 12q23.

SGPr452_1, SEQ ID NO:56, maps to human chromosomal position 16p13. 3.

SGPr504, SEQ ID NO:57, has not been assigned a chromosomal location.

SGPr469, SEQ ID NO:58, has not been assigned a chromosomal location.

- 5 SGPr400, SEQ ID NO:59, maps to human chromosomal position 4q32. Genomic amplification of this region has been associated with the following cancers (Knuutila): Non-small cell lung cancer.

EXAMPLE 8b: Candidate Single Nucleotide Polymorphisms (SNPs)

10 Materials And Methods

The most common variations in human DNA are single nucleotide polymorphisms (SNPs), which occur approximately once every 100 to 300 bases. Because SNPs are expected to facilitate large-scale association genetics studies, there has recently been great interest in SNP discovery and detection. Candidate
15 SNPs for the genes in this patent were identified by blastn searching the nucleic acid sequences against the public database of sequences containing documented SNPs (dbSNP, at NCBI, <http://www.ncbi.nlm.nih.gov/SNP/snpblastpretty.html>). dbSNP accession numbers for the SNP-containing sequences are given. SNPs were also identified by comparing several databases of expressed genes (dbEST, NRNA) and
20 genomic sequence (i.e., NRNA) for single basepair mismatches. The results are shown in Table 1, in the column labeled "SNPs". These are candidate SNPs – their actual frequency in the human population was not determined. The code below is standard for representing DNA sequence:

G = Guanosine
25 A = Adenosine
T = Thymidine
C = Cytidine
R = G or A, puRine
Y = C or T, pYrimidine

K = G or T, Keto
W = A or T, Weak (2 H-bonds)
S = C or G, Strong (3 H-bonds)
M = A or C, aMino

5 B = C, G or T (i.e., not A)
D = A, G or T (i.e., not C)
H = A, C or T (i.e., not G)
V = A, C or G (i.e., not T)
N = A, C, G or T, aNy

10 X = A, C, G or T

complementary DNA strands

G	A	T	C	R	Y	W	S	K	M	B	V	D	H	N	X
+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
C	T	A	G	Y	R	S	W	M	K	V	B	H	D	N	X

For example, if two versions of a gene exist, one with a "C" at a given position, and a second one with a "T" at the same position, then that position is represented as a Y, which means C or T. SNPs may be important in identifying heritable traits associated with a gene.

20 THE UNIVERSITY OF CHICAGO PRESS 1987

Results

The results of SNP identification are contained in Table 2 above, and in Example 1, under the section entitled DESCRIPTION OF NOVEL PROTEASE POLYNUCLEOTIDES. As discussed above, a variety of SNPs were identified in the protease polynucleotides of the present invention.

EXAMPLE 9: Demonstration Of Gene Amplification By Southern Blotting

Materials and Methods

Nylon membranes are purchased from Boehringer Mannheim. Denaturing solution contains 0.4 M NaOH and 0.6 M NaCl. Neutralization solution contains 0.5 M Tris-HCL, pH 7.5 and 1.5 M NaCl. Hybridization solution contains 50% formamide, 6X SSPE, 2.5X Denhardt's solution, 0.2 mg/mL denatured salmon DNA, 0.1 mg/mL yeast tRNA, and 0.2 % sodium dodecyl sulfate. Restriction enzymes are purchased from Boehringer Mannheim. Radiolabeled probes are prepared using the Prime-it II kit by Stratagene. The β -actin DNA fragment used for a probe template is purchased from Clontech.

Genomic DNA is isolated from a variety of tumor cell lines (such as MCF-7, MDA-MB-231, Calu-6, A549, HCT-15, HT-29, Colo 205, LS-180, DLD-1, HCT-116, PC3, CAPAN-2, MIA-PaCa-2, PANC-1, AsPc-1, BxPC-3, OVCAR-3, SKOV3, SW 626 and PA-1, and from two normal cell lines.

A 10 μ g aliquot of each genomic DNA sample is digested with EcoR I restriction enzyme and a separate 10 μ g sample is digested with Hind III restriction enzyme. The restriction-digested DNA samples are loaded onto a 0.7% agarose gel and, following electrophoretic separation, the DNA is capillary-transferred to a nylon membrane by standard methods (Sambrook, J. *et al.* (1989) Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory).

20 EXAMPLE 10: Detection Of Protein-Protein Interaction Through Phage Display

Materials And Methods

Phage display provides a method for isolating molecular interactions based on affinity for a desired bait. cDNA fragments cloned as fusions to phage coat proteins are displayed on the surface of the phage. Phage(s) interacting with a bait are enriched by affinity purification and the insert DNA from individual clones is analyzed.

T7 Phage Display Libraries

All libraries were constructed in the T7Select1-1b vector (Novagen) according to the manufacturer's directions.

Bait Presentation

Protein domains to be used as baits are generated as C-terminal fusions to
5 GST and expressed in *E. coli*. Peptides are chemically synthesized and biotinylated at the N-terminus using a long chain spacer biotin reagent.

Selection

Aliquots of refreshed libraries (10^{10} - 10^{12} pfu) supplemented with PanMix and a cocktail of *E. coli* inhibitors (Sigma P-8465) are incubated for 1-2 hrs at room
10 temperature with the immobilized baits. Unbound phage is extensively washed (at least 4 times) with wash buffer.

After 3-4 rounds of selection, bound phage is eluted in 100 μ L of 1% SDS and plated on agarose plates to obtain single plaques.

Identification of insert DNAs

15 Individual plaques are picked into 25 μ L of 10 mM EDTA and the phage is disrupted by heating at 70 °C for 10 min. 2 μ L of the disrupted phage are added to 50 μ L PCR reaction mix. The insert DNA is amplified by 35 rounds of thermal cycling (94 °C, 50 sec; 50 °C, 1min; 72 °C, 1min).

Composition of Buffer

20 10x PanMix
5% Triton X-100
10% non-fat dry milk (Carnation)
10 mM EGTA
250 mM NaF
25 250 μ g/mL Heparin (sigma)
250 μ g/mL sheared, boiled salmon sperm DNA (sigma)
0.05% Na azide
Prepared in PBS

Wash Buffer

PBS supplemented with:

0.5% NP-40

25 μ l g/mL heparin

5 PCR reaction mix

1.0 mL 10x PCR buffer (Perkin-Elmer, with 15 mM Mg)

0.2 mL each dNTPs (10 mM stock)

0.1 mL T7UP primer (15 pmol/ μ L) GGAGCTGTCGTATTCCAGTC

0.1 mL T7DN primer (15 pmol/ μ L)

10 AACCCCTCAAGACCCGTTTAG

0.2 mL 25 mM $MgCl_2$ or $MgSO_4$ to compensate for EDTA

Q.S. to 10 mL with distilled water

Add 1 unit of Taq polymerase per 50 μ L reaction

LIBRARY: T7 Select1-H441

15

Example 11: Gene Expression based on Incyte and Public ESTs

Materials and Methods

The nucleic acid sequences for the proteases were used as queries in a BLASTN search of the Incyte and public dbEST databases of expressed sequences.

- 20 The tissue sources of the libraries in which the protease was represented are listed below, along with the frequency the gene occurred in specific tissues. The frequency is determined by the number of clones representing the gene within a given tissue source. The Incyte gene identification number or public NCBI accession number is given, followed by the tissue source. A brief summary of the
- 25 tissue specificity is then given for each gene.

Results

SGPr397, SEQID:1,

Incyte 366783.1 Clones: 2 prostate, 3 colon, retina and small intestine

366783.3 1 prostate clone

Selective expression in prostate (3/8 clones) and colon (3/8 clones)

5

SGPr413, SEQID:2,

Incyte 475365.6 5 clones, 3 in small intestine, plus prostate, breast tumor

475365.5 10 clones: 8 in small intestine, plus brain (2)

Highly selective expression in small intestine (11/15 clones)

10

SGPr404, SEQID:3,

Incyte 1129157.1 213 clones, highest in brain (18), m/f genitalia (21/23), breast (14), and digestive (25)

1129157.2 1: mixed

15 Broad expression, some elevation in brain (18/214 clones), digestive tissues (25/118) and male/female genitalia (21, 23 clones) and breast (14 clones)

SGPr536_1, SEQID:4,

Incyte 233762.17 149 clones: no tissue >21 hits

20 233762.15 15 clones, mixed

Broad expression seen in 164 clones

SGPr414, SEQID:5,

Incyte 399773.5 669 clones, 404 libraries, broadly distributed

25 Expressed broadly and strongly (669 clones)

SGPr430, SEQID:6,

Incyte 407823.1 21 clones (4 testis, 3 brain, 3 prostate)

1136483.1 1 Prostate

- 411246.1 2 ea lung tumor, sm intestine, fetal liver, and 1 heart
 322700.1 T cells
 407823.2 Fetal liver/spleen
 Mixed expression, highest in testis (4/31 clones), brain (3/31) and prostate (4/31)
 5 and fetal liver/spleen (4/31)
- SGPr496_1, SEQID:7, 12 clones: 2 ea lung, brain, adrenal tumor
 Incyte 986031.1
 Selective expression in lung (2/12 clones), adrenal tumor (2/12) and brain (2/12)
- 10 SGPr495, SEQID:8, 22 clones: 7 liver, 1 fetal liver/spleen, 3 lung
 Incyte 350921.2 16 clones: 2 thymus, 3 colon, 3 brain
 350921.7 Adrenal tumor
 350921.10 14 clones: 2 colon, 1 adrenal tumor
 15 350921.6 10 clones: 2 adrenal (1 tumor), 2 brain
 350921.1 Adrenal
 350921.9 Colon (2) Sm intestine, lung tumor
 Selectively expressed in adrenal gland (5/46 clones) and colon (7/46)
- 20 SGPr407, SEQID:9, No ESTs
 SGPr453, SEQID:10,
 Incyte 428428.1 17 clones: 3 lung (2 tumors), 2 prostate, 4 testis, 2 teratoma
 25 (hNT2)
 428428.5 brain, lung, teratoma
 428428.6 teratoma (2), lung, kidney
 Highly expressed in hNT2 teratoma cell line (5/24 clones), and selective for lung (5 clones) and testis (4 clones)

SGPr445, SEQID:11,

- Incyte 350921.7 1 adrenal tumor
 350921.10 14 incl 1 adrenal tumor
 5 350921.6 10 clones: normal and tumor adrenal (1 ea) colon tumor (2)
 350921.1 Adrenal gland
 350921.8 2 prostate tumor, 1 retina

Highest in adrenal gland (5/28 clones), indicates a possible involvement in adrenal hormone processing

10

SGPr401_1, SEQID:12,

- Incyte 232414.1 169 clones: 69 NS, 18 male genitalia, (10 prostate), 11 female genitalia, 11 respiratory system, 5 kidney, 9 in one glioblastoma library.
 232414.2 testis
 15 Selective for nervous system (69/170 clones), especially glioblastoma

SGPr408, SEQID:13,

- Incyte 233660.2 357 clones, 248 libraries: 54 brain, 26 hemic/immune 24/21 f/m genitalia, 21 digestive, 20 cardiovascular
 20 233660.11 13 clones, broad expression.
 233660.10 7 clones, broad expression
 Expressed broadly and strongly (377 clones)

SGPr480, SEQID:14,

- 25 Incyte 1326256.3 274 clones, broad but highest in NS (59), hemic (35) genitalia (24/10, m/f). 6 clones in one pituitary gland library
 1326256.8 4 mixed clones
 1326256.1 26 clones, 13 in male genitalia
 1326256.10 10 clones mixed

Broad, strong expression (over 300 clones)

SGPr431, SEQID:15,

Incyte 236368.1 151 clones, 110 libraries, highest in digestive, nervous, hemic

5 (18, 17, 18): 5, 4 hits each in two fetal liver/spleen libraries

236368.2: 1 fetal heart

236368.14: 7: mixed

Broad and moderately strong expression (159 clones total)

10 SGPr429, SEQID:16, 41 clones, broad

Incyte 890540.9, 41 clones, broad

890540.1: 125 clones, broad

890540.8: 15 clones, broad

Broad and moderately strong expression (181 clones)

15 SGPr503, SEQID:17, 107 clones highest in NS (15), male genitalia (11) and

digestive tissue (15)

1447357.3 107 clones highest in NS (15), male genitalia (11) and

digestive tissue (15)

1447357.1: Dendritic cells

20 245045.1 16 mixed

Broad expression (124 clones), highest in nervous system (16), male genitalia (11)

and digestive tissue (15)

SGPr427, SEQID:18,

25 Incyte 903092.31 41 clones, 35 libraries; 9 clones in prostate, otherwise very

broad

903092.23 1 brain

Expression elevated in prostate (9/42 clones)

SGPr092, SEQID:19,

Incyte 339251.1 4/5 uterus, 1 mixed tissue

339251.2 1/1 uterus; Highly selective expression in uterus (5/6 clones)

5 SGPr359, SEQID:20,

Incyte 391133.1 mixed tissue (fetal lung, testis, B-cell)

gi|7280399 = same as Incyte

Mixed tissues, one EST only

10 SGPr104_1, SEQID:21,

Incyte 12/23 clones in brain

232015.5 6/7 clones in brain

232015.2 2/4 clones in brain

232015.1 1/1 brain

15 232015.6 1/1 brain

Brain specific

SGPr303, SEQID:22,

Incyte 323846.15 38 samples, highest in brain(8) breast(4), uterus and ovary(7)

20 323846.1 304 clones, high in nervous sys(72) and genitalia (28 f, 24 m),
other tissue

414048.34 45 clones, highest in NS

323846.11 8 clones

Broad expression

25

SGPr402_1, SEQID:23,

Incyte 244407.4 25 clones, highest in testis (8) brain (4), uterus (2; 1 tumor)

244407.2 uterus

244407.1 testis

- 244407.6 uterus tumor
- 244407.5 fallopian tube tumor
- 244407.9 mixed tissue incl tumor, nasal tumor
- Enriched in genital samples.
- 5
- SGPr434, SEQID:24,
- Incyte 110154.4 no clone origin
- 110154.6 2 prostate
- 110154.12 1 prostate, 1 pituitary
- 10 110154.11 1 pituitary
- 110154.8 fallopian tube tumor (2), mixed (1)
- 110154.7 Pituitary
- 110154.5 Thigh muscle (2) – tissue-specific splicing
- 110154.10 3 heart, 2 brain, pituitary (61/62 match)
- 15 Selective expression in prostate (3/17 clones), 4 pituitary gland (4/17 clones) and fallopian tube tumor (2/17 clones). May indicate a role in hormone processing in pituitary and prostate.- hormone processing.
- SGPr446_1, SEQID:25,
- 20 Incyte 1040641.1 Heart, Muscle
- 1388371.1 2 Heart
- Specific for muscle (3/3 clones) especially heart muscle (2/3 clones)
- SGPr447, SEQID:26,
- 25 Incyte 1352932.1 pancreas tumor
- Single clone from pancreas tumor
- SGPr432_1, SEQID:27,
- Incyte 474674.15 29 clones, mixed

474674.30 90 clones, mixed
 474674.1 82 clones, mixed
 Broad and strong expression (201 clones total)

- 5 SGPr529, SEQID:28,
 Incyte 988019.3 71 clones, 23 in f genitalia. 9 from 1 ovary tumor library, 2 from another, 2 from another, and one from yet another (no normal ovaries). 5 from one pancreatic tumor line, 4 from pancreas tumor library
 988019.1 breast skin
- 10 Selective expression in pancreas (4/72 clones from one pancreatic tumor library and 5 from a pancreatic tumor line) and ovary (14 from ovary tumors, none from normal ovary).

- SGPr428_1, SEQID:29,
- 15 Incyte 891146.1 4: brain, pituitary, blood, thymus
 Broad, low-level expression (4 clones all from different tissues)

- SGPr425, SEQID:30,
 Incyte 400833.1 25 clones, mixed (<4 from any tissue, except 5 from 'fetus')
- 20 Expressed broadly but not strongly (25 clones total)

- SGPr548, SEQID:31,
 Incyte 971236.1 2 clones from mixed testis, fetal lung, B cells
 Rare transcript, just two clones from a mixed library of testis, fetal lung and B cells
- 25

SGPr396, SEQID:32,
 Incyte 209051.1 Lung (1)
 889126.1 Brain (1)
 Only 2 ESTs – lung and brain

- SGPr426, SEQID:33,
Incye No ESTs
- 5 SGPr552, SEQID:34,
Incye 1510512.1 tonsil, spinal cord
1511222.1 tonsil
406221.1 83 clones: 16 in NS, 10 in hemic/immune; 9 in male genitalia,
and several other tissues: 1 tonsil,
10 981355.3 8 clones, 2 ovary tumor; 1 tonsil, varied
Of 94 clones, see some selectivity in tonsil (3/94, but tonsil not usually seen as an
expression source), and nervous system (17 clones)
- SGPr405, SEQID:35,
15 Incye 134360.10 1 kidney
One clone, from kidney
- SGPr485_1, SEQID:36,
Incye 180576.2 5/5 clones in testis
20 Testis specific (5/5 clones)
- SGPr534, SEQID:37,
Incye 1383391.20 112/114, matches well at start (103-165 = perfect match) but
maybe template artefact
- 25 1450812.1 1/1 pancreas, few mismatches are N's
1383391.13 5/5 pancreas
1045834.1 1/1 pancreas
Almost completely pancreas-specific (118/120 clones from pancreas).

SGPr390, SEQID:38,

Incye 199428.9 Bone tumor, small intestine

199428.3 382 clones: 41 brain, 34/23 genitalia (m/f), 22 hemic/immune,
27 digestive

- 5 Broad tissue distribution, highest in brain (41/382 clones), male and female genitalia (34 and 23 clones, respectively) and digestive system (27 clones)

SGPr521, SEQID:39,

Incye 427826.1 28 clones, most in sm intestine tumor (5, 1 library), neonatal

- 10 keratinocytes (3 ea from 2 libraries), 8 ovary tumors, 5 breast skin

Selective expression in ovarian tumors (8/28 clones), neonatal keratinocytes (6/28), breast keratinocytes (5) and in a small intestine tumor library (5 clones from one library)

- 15 SGPr530_1, SEQID:40,
No ESTs

SGPr520, SEQID:41,

Incye 405947.1 4/4 clones adrenal tumor (pheochromocytoma) (3 from one
20 library, 1 from another)

1338652.1 1/1 clones from adrenal tumor (pheochromocytoma)

1477189.1 1/1 clones from adrenal (mixed normal and
pheochromocytoma)

- 25 Specific to pheochromocytoma (adrenal gland tumor): 4/5 clones from
pheochromocytoma and 1/5 from mixed normal adrenal gland and
pheochromocytoma.

SGPr455, SEQID:42,

Incye 1115833.1 mixed fetal lung/testis/Bcell

- 987279.1 Brain (1), mixed tissues incl tumor (1)
Three clones, only one (brain) with a specific source
- SGPr507_2, SEQID:43,
- 5 Incyte 403891.1 10 clones: 6 in testis and 6 in mixed (testis, lung, Bcell)
403891.2 1 brain
Testis-selective: 6/11 clones from testis and 5/11 from mixed libraries including
testis samples
- 10 SGPr559, SEQID:44,
Incyte 475100.1 35 clones, 11 in genitalia, 8 in digestive, 7 uterus tumors
(none normal), 4 in breast, 2 ovary tumors, 1 HeLa cervical tumor
475100.6 Th1 cells, HeLa cells
Selective expression in tumors of the uterus (7/37 clones), ovary (2/37), cervix (2/37)
15 from HeLa cervical tumor cell line), as well as breast (4)
- SGPr567_1, SEQID:45,
- Incyte 981355.3 Mixed (2/8 clones from ovary tumor library, 1 ea from tonsil,
brain, lung tumor, heart, placenta, dorsal root ganglion)
- 20 Rare broad expression (8 clones from 7 different tissues).
- SGPr479_1, SEQID:46,
- Incyte 219214.1 1 testis
Single EST, expressed in testis
- 25 SGPr489_1, SEQID:47,
Incyte 338956.1 2 kidney, 1 placenta
1042306.1 1 mouth tumor, 2 fallopian tube tumor
1384824.1 Sm intestine, kidney

Rare but broad expression, selective to kidney (3/8 clones) and fallopian tube tumor (2/8 from one library)

SGPr465_1, SEQID:48,

5 No ESTs

SGPr524_1, SEQID:49,

Incyte 952182.3 1 testis

952182.2 1 testis

10 952182.4 1 prostate

Specific to male genitalia (2/3 clones in testis, 1/3 in prostate)

SGPr422, SEQID:50,

Incyte 1511284.1 1 tonsil

15 1351259.1 1 brain

Rare transcript seen only in tonsil (1/2 clones) and brain (1/2 clones)

SGPr538, SEQID:51,

Incyte 903092.29 4 clones, 3 brain, 1 breast

20 903092.19 1 brain

903092.22 2 brain

903092.28 38 clones, 24 in brain

903092.24 2 brain, 1 sm intestine

Selective expression in nervous system (32/48 clones)

25

SGPr527_1, SEQID:52,

Incyte 65450.1 1 prostate tumor

103554.2 mixed tissues incl tumor

228456.2 11 clones: mixed (2 brain, 2 blood)

103554.1 Mixed (3)
Broad low-level expression (16 clones)

SGPr542, SEQID:53,

- 5 Incyte 244085.1 Expression is selective to hemopoetic cells: All 11 clones are from hemopoetic tissues: 6 from fetal liver/spleen, 4 of which are mast cells, 2 from umbilical cord blood, 1 from CD34+ bone marrow, and two clones from leukemias: 1 from AML blast cells and one from CML.
- 10 SGPr551, SEQID:54,
Incyte 319529.1 22 clones: 7 liver, 1 fetal liver/spleen, 3 lung,
319529.2 1 liver
319529.3 2 mixed tissues incl testis, 1 testis (tissue-specific splice)
Selectively expressed in liver (9/26 clones), may have a testis-specific splice form
15 (3/3 clones of one template)
- 20 SGPr451, SEQID:55,
Incyte 1471541.1 1 mixed)
- 20 SGPr452_1, SEQID:56,
Incyte 446374.1 Mixed (melanocytes, uterus, fetal heart).
No expression data (single EST from mixed tissues)
- 25 SGPr504, SEQID:57,
Incyte 244085.1 Expression is selective to hemopoetic cells: All 11 clones are from hemopoetic tissues: 6 from fetal liver/spleen, 4 of which are mast cells, 2 from umbilical cord blood, 1 from CD34+ bone marrow, and two clones from leukemias: 1 from AML blast cells and one from CML.

SGPr469, SEQID:58,

Incyte 110154.10 7 clones: 3 heart, 2 brain, 1 pituitary

110154.3 heart, muscle, testis

Selective expression in heart (4/10 clones)

5

SGPr400, SEQID:59,

Incyte 889126.1 Brain

Only one EST, in brain

10

CONCLUSION

One skilled in the art would readily appreciate that the present invention is well adapted to carry out the objects and obtain the ends and advantages mentioned, as well as those inherent therein. The molecular complexes and the methods, procedures, treatments, molecules, specific compounds described herein are presently representative of preferred embodiments, are exemplary, and are not intended as limitations on the scope of the invention. It will be readily apparent to one skilled in the art that varying substitutions and modifications may be made to the invention disclosed herein without departing from the scope and spirit of the invention.

20 All patents and publications mentioned in the specification are indicative of the levels of those skilled in the art to which the invention pertains. All patents and publications are herein incorporated by reference to the same extent as if each individual publication was specifically and individually indicated to be incorporated by reference.

25 The invention illustratively described herein suitably may be practiced in the absence of any element or elements, limitation or limitations which is not specifically disclosed herein. Thus, for example, in each instance herein any of the terms "comprising," "consisting essentially of" and "consisting of" may be replaced

with either of the other two terms. The terms and expressions which have been employed are used as terms of description and not of limitation, and there is no intention that in the use of such terms and expressions of excluding any equivalents of the features shown and described or portions thereof, but it is recognized that
5 various modifications are possible within the scope of the invention claimed. Thus, it should be understood that although the present invention has been specifically disclosed by preferred embodiments and optional features, modification and variation of the concepts herein disclosed may be resorted to by those skilled in the art, and that such modifications and variations are considered to be within the scope
10 of this invention as defined by the appended claims.

In addition, where features or aspects of the invention are described in terms of Markush groups, those skilled in the art will recognize that the invention is also thereby described in terms of any individual member or subgroup of members of the Markush group. For example, if X is described as selected from the group
15 consisting of bromine, chlorine, and iodine, claims for X being bromine and claims for X being bromine and chlorine are fully described.

In view of the degeneracy of the genetic code, other combinations of nucleic acids also encode the claimed peptides and proteins of the invention. For example, all four nucleic acid sequences GCT, GCC, GCA, and GCG encode the amino acid
20 alanine. Therefore, if for an amino acid there exists an average of three codons, a polypeptide of 100 amino acids in length will, on average, be encoded by 3100, or 5×10^4 , nucleic acid sequences. Thus, a nucleic acid sequence can be modified to form a second nucleic acid sequence, encoding the same polypeptide as encoded by the first nucleic acid sequences, using routine procedures and without undue
25 experimentation. Thus, all possible nucleic acids that encode the claimed peptides and proteins are also fully described herein, as if all were written out in full taking into account the codon usage, especially that preferred in humans. Furthermore, changes in the amino acid sequences of polypeptides, or in the corresponding nucleic acid sequence encoding such polypeptide, may be designed or selected to

take place in an area of the sequence where the significant activity of the polypeptide remains unchanged. For example, an amino acid change may take place within a β -turn, away from the active site of the polypeptide. Also changes such as deletions (*e.g.* removal of a segment of the polypeptide, or in the corresponding
5 nucleic acid sequence encoding such polypeptide, which does not affect the active site) and additions (*e.g.* addition of more amino acids to the polypeptide sequence without affecting the function of the active site, such as the formation of GST-fusion proteins, or additions in the corresponding nucleic acid sequence encoding such polypeptide without affecting the function of the active site) are also within the
10 scope of the present invention. Such changes to the polypeptides can be performed by those with ordinary skill in the art using routine procedures and without undue experimentation. Thus, all possible nucleic and/or amino acid sequences that can readily be determined not to affect a significant activity of the peptide or protein of the invention are also fully described herein.

15 The invention has been described broadly and generically herein. Each of the narrower species and subgeneric groupings falling within the generic disclosure also form part of the invention. This includes the generic description of the invention with a proviso or negative limitation removing any subject matter from the genus, regardless of whether or not the excised material is specifically recited
20 herein.

Other embodiments are within the following claims.

What is claimed is:

CLAIMS

1. An isolated, enriched or purified nucleic acid molecule, wherein said nucleic acid molecule comprises a nucleotide sequence that:
- 5 (a) encodes a polypeptide having an amino acid sequence selected from the group consisting of those set forth in SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118 and biological domains thereof;
- 10 (b) is the complement of the nucleotide sequence of (a); or
- 15 (c) hybridizes under stringent conditions to the nucleotide molecule of (a) and encodes a protease polypeptide.
- 20
- 25
2. The nucleic acid molecule of claim 1, further comprising a vector or promoter operatively linked to the nucleotide sequence.

3. The nucleic acid molecule of claim 1, wherein said nucleic acid molecule is isolated, enriched, or purified from a mammal.
- 5 4. The nucleic acid molecule of claim 3, wherein said mammal is a human.
- 10 5. The nucleic acid molecule of claim 1 comprising a nucleic acid comprising a nucleotide sequence which hybridizes under stringent conditions to a nucleotide sequence encoding a protease polypeptide having an amino acid sequence selected from the group consisting of those set forth in SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118.
- 15
- 20
- 25

- 5 6. An isolated, enriched, or purified protease polypeptide, wherein said polypeptide comprises an amino acid sequence at least about 90% identical to a sequence selected from the group consisting of those set forth in SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118 and biological domains thereof.
- 10 7. The protease polypeptide of claim 6, wherein said polypeptide is isolated, purified, or enriched from a mammal.
- 15 8. The protease polypeptide of claim 7, wherein said mammal is a human.
- 20 9. An antibody or antibody fragment having specific binding affinity to a protease polypeptide or to a domain of said polypeptide, wherein said polypeptide comprises an amino acid sequence selected from the
- 25

5 group consisting of those set forth in SEQ ID NO:60, SEQ ID NO:61,
SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65,
SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69,
SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73,
SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77,
SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81,
SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85,
SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89,
SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93,
10 SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97,
SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101,
SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID
NO:105, SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:108, SEQ
ID NO:109, SEQ ID NO:110, SEQ ID NO:111, SEQ ID NO:112,
15 SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID
NO:116, SEQ ID NO:117 and SEQ ID NO:118.

10. A hybridoma which produces the antibody of Claim 9.
- 20 11. A kit comprising an antibody which binds to a polypeptide of claim 6
and a negative control antibody.
12. A method for identifying a substance that modulates the activity of a
protease polypeptide comprising the steps of:
25 (a) contacting a protease polypeptide substantially identical to an
amino acid sequence selected from the group consisting of
those set forth in SEQ ID NO:60, SEQ ID NO:61, SEQ ID
NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65,
SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID

- 5 NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72,
SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID
NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79,
SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID
NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86,
SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID
NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93,
SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID
NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100,
10 SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID
NO:104, SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:107,
SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID
NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114,
SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ
15 ID NO:118 with a test substance;
(b) measuring the activity of said polypeptide; and
(c) determining whether said substance modulates the activity of
said polypeptide.
- 20 13. A method for identifying a substance that modulates the activity of a
protease polypeptide in a cell comprising the steps of:
(a) expressing a protease polypeptide in a cell, wherein said
polypeptide comprises a sequence substantially identical to an
25 amino acid sequence selected from the group consisting of
those set forth in SEQ ID NO:60, SEQ ID NO:61, SEQ ID
NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65,
SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID
NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72,
SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID

- 5 NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79,
SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID
NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86,
SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID
NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93,
SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID
NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100,
SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID
NO:104, SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:107,
10 SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID
NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114,
SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ
ID NO:118;
- 15 (b) adding a test substance to said cell; and
(c) monitoring a change in cell phenotype, cell proliferation, cell
differentiation or the interaction between said polypeptide and
a natural binding partner.
- 20 14. A method for treating a disease or disorder by administering to a
patient in need of such treatment a substance that modulates the
activity of a protease substantially identical to an amino acid
sequence selected from the group consisting of those set forth in SEQ
ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID
NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID
NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID
25 NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID
NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID
NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID
NO:84, SEQ ID NO:85, SEQ ID NO:86, SEQ ID NO:87, SEQ ID

NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118.

- 5
- 10
- 15
- 15
- 20
- 25
15. The method of claim 14, wherein said disease or disorder is selected from the group consisting of cancers, immune-related diseases and disorders, cardiovascular disease, brain or neuronal-associated diseases, metabolic disorders and inflammatory disorders.
16. The method of claim 15, wherein said disease or disorder is selected from the group consisting of cancers of tissues; cancers of blood or hematopoietic origin; cancers of the breast, colon, lung, prostate, cervical, brain, ovarian, bladder or kidney.
17. The method of claim 15, wherein said disease or disorder is selected from the group consisting of central or peripheral nervous system diseases, migraines; pain; sexual dysfunction; mood disorders; attention disorders; cognition disorders; hypotension; hypertension; psychotic disorders; neurological disorders and dyskinesias.
18. The method of claim 15, wherein said substance modulates protease activity *in vitro*.

19. The method of claim 18, wherein said substance is a protease inhibitor.
20. A method for detection of a protease polypeptide in a sample as a diagnostic tool for a disease or disorder, wherein said method comprises:
- (a) contacting said sample with a nucleic acid probe which hybridizes under hybridization assay conditions to a nucleic acid target region of a protease polypeptide having an amino acid sequence selected from the group consisting of those set forth in SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118, said probe comprising the nucleic acid sequence encoding the polypeptide, fragments thereof, and the complements of the sequences and fragments; and

(b) detecting the presence or amount of the probe:target region hybrid as an indication of the disease.

- 5 21. The method of claim 20, wherein said disease or disorder is selected from the group consisting of cancers, immune-related diseases and disorders, cardiovascular disease, brain or neuronal-associated diseases, metabolic disorders and inflammatory disorders.
- 10 22. The method of claim 21, wherein said disease or disorder is selected from the group consisting of cancers of tissues; cancers of hematopoietic cancers of blood or hematopoietic origin; cancers of the breast, colon, lung, prostate, cervical, brain, ovarian, bladder or kidney.
- 15 23. The method of claim 21, wherein said disease or disorder is selected from the group consisting of central or peripheral nervous systems disease, migraines, pain; sexual dysfunction; mood disorders; attention disorders; cognition disorders; hypotension; hypertension; psychotic disorders; neurological disorders; and dyskinesias.
- 20 24. The isolated, enriched or purified nucleic acid molecule of claim 1 comprising a nucleic molecule encoding a biological domain of a protease polypeptide having a sequence selected from the group consisting of SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID
- 25

5 NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86, SEQ ID
 NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID
 NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID
 NO:95, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:98, SEQ ID
 NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ ID
 NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106, SEQ
 ID NO:107, SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110,
 SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID
 NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and
 10 SEQ ID NO:118.

25. The nucleic acid molecule of claim 1 comprising a nucleic acid
 sequence encoding a protease polypeptide having an amino acid
 sequence that has least 90 % identity to a polypeptide selected from
 15 the group consisting of those set forth in SEQ ID NO:60, SEQ ID
 NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID
 NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID
 NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID
 NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID
 20 NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:80, SEQ ID
 NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID
 NO:85, SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, SEQ ID
 NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID
 NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID
 25 NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID
 NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ
 ID NO:105, SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:108,
 SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:111, SEQ ID

NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118.

- 5 26. The nucleic acid molecule of Claim 1 wherein the molecule comprises a nucleotide sequence substantially identical to a sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:44, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:58, and SEQ ID NO:59.
- 10 27. An isolated, enriched or purified nucleic acid molecule consisting essentially of about 10-30 contiguous nucleotide bases of a nucleic acid sequence that encodes a polypeptide that is selected from the group consisting of SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74,
- 15 20 25

5 SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78,
SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82,
SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86,
SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90,
SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94,
SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:98,
SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102,
SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID
10 NO:106, SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:109, SEQ
ID NO:110, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113,
SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID
NO:117 and SEQ ID NO:118.

28. 15 The isolated, enriched or purified nucleic acid molecule of Claim 27
consisting essentially of about 10-30 contiguous nucleotide bases of a
nucleic acid sequence selected from the group consisting of SEQ ID
NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5,
SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID
NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID
20 NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID
NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID
NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID
NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID
NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID
25 NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID
NO:38, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:41, SEQ ID
NO:42, SEQ ID NO:43, SEQ ID NO:44, SEQ ID NO:45, SEQ ID
NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID
NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID

NO:54, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:58, and SEQ ID NO:59.

29. A recombinant cell comprising the nucleic acid molecule of claim 1.

30. A method for detecting the presence or amount of protease polypeptide in a sample comprising

(a) contacting the sample with the antibody of Claim 9 under conditions suitable for protease-antibody immunocomplex formation; and

(b) detecting the presence or amount of the antibody conjugated to the protease polypeptide.

FIG. 1A

>SGPR397_SEQID_1
 ATGAAGTGTCTCGGGAAGCGCAGGGGCCAGGCTGCTTTCCTGCCCTCTTTGCTGGCTCTTTTGAAGATTCT
 GCAACCGGGGCACAGCCACCTTTTATAACAACCGCTATGCTGGTGATAAGTGATAAGATTATTCCTCAAAACAGA
 AGAGGAAGCATATGCACCTGAAGAAAATATCCTATCAACTTAAGGTGGACCTGTGGCAGCCAGCAGTATCTCCTA
 TGTATCAGAGGGAACAGTTACTGATGTCCATATCCCCAAAATGTTCCCGAGCCCTGTAGCCTTCTTACAGGA
 AGCCAACATCCAGTACAAGGTCTCTATAGAAGATCTTCAGAAAACACTGGAGAAAGGGAAGCAGCTTGCACACCC
 AGAGAAACCGAAGATCCCTCTCTGGATATAATTATGAAGTTTATCACTCCTTAGAAGAAATTCAAAATGGATGCA
 TCATCTGAATAAAACTCACTCAGGCCCTCATTCACATGTTCTCTATTGGAAGATCATATGAGGGAAGATGCTCTTTT
 ATTTAAAGCTGGCAGACGATCACGACTCAAAGAGCTGTTGGATAGACTGTGGTATTCATGCAAGAGAATGG
 ATTGGTCTGCCCTTTGTCAGTGGTTGTAAAGAAGCTCTTCTAACATATAAGAGTGACCCAGCCATGAGAAAAA
 TGCTGAATCATCTATATTTCTATATCATGCTGTGTTAACGTCGATGGATACCATTTTAGTTGGACCAATGATCGA
 TTTTGGAGAAAAACAAGGTCAAGGAACCAAGGTTTCGCTGCCGTGGAGTGGATGCCAATAGAAACTGGAAAGT
 GAAGTGGTGGTAAGTTTGGGACCAACTGGGATCCAGATCCAAAGGTTTCTGCAGGTTTACTCTGCAAAATAT
 GAGTCCAGAGGACTCTCATGGGAGACTCATGTTTTTCTGTATGTGA

>SGPR413_SEQID_2
 ATGAAGCCTCTGCTTGAAACCCCTTATCTTTGGGGATGCTGGTTCCTGGAGGGCTGGGATATGATAGATCCTTA
 GCCCAACACAGACAGAGATTGTGGACAAGTCAGTGAGTCCATGGAGCCTGGAGACGTATTCCTATAACATATAC
 CACCCCATGGGAGAGATCTATGAGTGGATGAGAGAGATCAGTGAGAAGTACAAGGAAGTGGTGACACAGCATTT
 CCTAGGAGTGACCTATGAGACCCACCCCATGTATTATCTGAAGATCAGCCCAACCATCTGGTAATCCCAAGAAAT
 CATTGGATGGACTGTGGAATTCACGCCAGAGAAATGGATTGCTCCTGCTTTTCCCAATGGTTCGTCAAAGAAAT
 TCTACAAAACCATAAAGACAACCTCAAGTATACGCAAGCTCCTTAGGAACCTGGACTTCTATGTCTCCAGTTCTT
 AACATAGATGGTTATATCTACACTTGGACAACTGATCGTCTTTGGAGGAAATCCCGTTCACCCCATATAATGGCA
 CATGTTTTGGGACGGATCTCAATCGAAATTTCAATGCATCTTGGTGTAGTATTGGTGCCTCTAGAAACTGCCAAGA
 TCAAAACATCTGTGGACAGGGCCAGTGTCTGAACCCAGAGACTAAAGCTGTTGCCAGCTTCATAGAGAGCAAGA
 AGGATGATATTTTGTGCTTCTGACCATGCACCTCTTATGGCAGTTAATTCACACCTTACGGCTACACCAAAA
 TAAATCAAGTAACCAACCCAGAAATGATTCAAGTTGGACAGAAAGGAGCAAAATGCATTGAAGCAAGTATGGAAC
 CAATTATAGAGTTGGATCGAGTGCAGATATTTTATATGCCCTCATCAGGCTCTTCAAGAGATTGGCCCGAGACAT
 TGGGATCCCTTCTCATATACGTTTGGAGCTGAGGACAGTGAACATATGGGTTTGTCTGCCAGAAAGCTCAGAT
 CCAGCCCACTGTGAGGAGACCATGGAGGCTGTGCTGTAGTCTGATGTGTATGCGAAACACTGGCACT
 CGGACAGTGTGGAAGGGTGACATCTGCCACTATGCTGCTGGCCCTGCTGGTGTCTGCTGCTCTCTCTCTAA

>SGPR536 1 SEQID 4

ATGTGGCGATGTC¹CACTGGGCTACTGCTGTTGGCTGCCGCTGGCTGGCACTTGGCTCTGGCTGCCCAGCAGG
GTCTGGCGCGCGGAGCTAGCACCGGCTCTGCACCTGCGGGGCAATCGGGACGCGGGAGCGCGGTACTGC
CAGGAGCAGGACCTGTGCTGCCGCGCGCTGCCGACGACTGTGCCCTGCCCTACCTGGCGGCCATCTGTTACT
GTGACCTCTTCTGCAACCGCACGGTCTCCGACTGCTGCCCTGACTTCTGGGACTTCTGCCTCGCGTGCCACCC
CCTTTTCCCGGATCCAGGATGTATGCATGGAGTCTGTATCTATCCAGTCTGGGAACGTACTGGGACAACCTGT
AAACCGTTGCACCTGCCAGGAGAACAGGCAGTGGCAGTGTGACCAAGAACCATGCCTGTGTGATCCAGACATGA
TCAAAGCCATCAACGAGGGCAACTATGGCTGGCAGGCTGGGAACCAACAGCGGCTTCTGGGGCATGACCCCTGGA

FIG. 1C

TGAGGGCATTGCTACCGCCTGGGCACCATCCGCCCATCTTCCTCGGTCATGAACATGCATGAAATTTATACAGT
 GCTGAACCCAGGGAGGTGCTTCCACAGCCTTCGAGGCCCTCTGAGAACTGGCCCAACCTGATTCATGAGCCT
 CTTGACCAAGGCAACTGTGCAGGCTCCTGGCCCTTCTCCACAGCAGCTGTGGCATCCGATCGTGTCTCAATCCA
 TTCTCTGGACACATGACGCTGTCTGTGCCCCAGAACCTGCTGTCTTGACACCCACAGCAGCAGGGCT
 GCCGCGTGGCGTCTCGATGGTGGTGGTCTGCTGCGCCGAGGGTGGTGTCTGACCACTGCTACC
 CCTTCTGGGCGTGAACGAGACGAGGCTGGCCCTGCGCCCTGTATGATGCACAGCCGAGCCATGGGTCG
 GGGCAAGCGCCAGGCCACTGCCCACTGCCCAACAGCTATGTTAATAACAATGACATCTACCAGGTCACCTCTG
 TCTACCGCTCGGCTCCAAACGACGAGGATCATGAAGGAGCTGATGGAGATGGCCCTGTCCAAGCCCTCATG
 GAGGTGCATGAGGACTTCTTCTATACAGGGAGGCTACAGCCACACGCCAGTGGCCCTTGGAGGCCAG
 AGAGATACCGCGCATGGACCCACTCAGTCAAGATCACAGGATGGGAGAGGAGACGCTGCCAGATGGAAG
 GACGCTCAATACTGACTCGGCCCACTCTGGGCCAGCCCTGGGCGAGAGGGCCACTTCCGCATCGT
 GCGCGCGTCAATGAGTGCAGACATCGAGAGCTTCGTGGCGCTCTGGGCGCGCTGGGCATGGAGGACAT
 GGGTCATCACTGA

>SGPR414_SEQID_5

ATGTGCGAAGAACTGCGCAGACCTGGTGGAGGTAAATGAATATCAGATGTAGAAGGTGGTGATGGACTGCA
 GCTCAGAAAGGAACATACTCTCAAAATATTTACTTACATCAATTCCTGGACACAGAGGCAATGTCTATGCTGCTTC
 AAGGAATATAAGCAATTTGGAGATTTTAAATCAAGTAGTGTGACCTTATTAACCTTAGTGATTGCCCAAGTTCAGT
 GCTCCGGGACCAGCTTTGTAACATTTGTACTACCATTAACATAGATCCACGTGGCAAGATGAGAGTAATCAAGC
 AGAAGAACCACCTGAATATAGATAGAGAGTGTAAATGAAGGAAGTACAGAAAGACAAAATCAATAGAAAAAAATCA
 AACTCTACAAGAAATTTGTAATCTGACTGAGGAGGAATCTTCAAAGAGTTCTGATCCCTTTAGTTTATGGAGTACAG
 ATGAGAAGGAAAAACTCTTACTATGTGTGGCAAAATTTTCAAATTCAGTTCCCTTATATACTGCTTACAAGCAT
 AATACTACCCCTACTATTGAGGATATATCAACTCAAGAAAGTAACATATTAGGGCATTCTGTGATATGAATGATG
 TAGAAGTACCATTGCAATTTGCTTCGTTATGTATGTTTGTGGGAAAAATGGCCCTTCTCTCATGAAGGATTG
 CTTTGAATATGGAACCTCTGAACTTTGCCATTTCTATAGCACATGCGTTTATTACAGTTGTCTAATATTAGAA
 TATGGCTACATAATCCCGCTGTATGCAGCACATTATACCTTTTAGACCTATGTTATTAGGTATTTATGCAAGCT
 CTCGGATCAGGAGTTACGACAGAGTGCAGCTCGTAACATGGCTGACTTAATGTGGACACAGTCAAAGAACCAT
 TGGATACAACATTTATGCTTTGATAAAGAAAGCCTAGATCTTGCAATTAAGTACTTTATGTCACTACTTTGACTATG
 AGTTGGCTGGATTGAGTCAGATAACAAATCAACTCCATACCTTCAATGATGTGTGCAATAATGAATCATTAGTAT
 CGGACACAGAAACGTCCATTGCAAAAGAACTTGACAGCTGGCTTATTAGCAACAATGTGGTGAGCATATATTG
 GACCAAAATTTACATATTGAGATTATCAACAGTGCCTAAGTATTTGAATTTTGGCAGCAGAGGGCGACTGAG
 TACTCAACATATTGACTGTATTTGGGCTGCAGCACAGTTGAACATTTGTAGTCGGTATATACATGACTTATTTCCCTT

FIG. 1D

CACTCATCAAGAAATTGGATCCCGTACCACCTTAGACATCTACTTAATCTGGTCTCAGCTCTTGAGCCAAAGTGTTCA
TACTGAACAGACACTGTACTTGGCATCCATGTTAATTAAAGCACTGTGGAATAACGCCTAGCAGCTAAGGCTCA
GTTATCTAAACAGAGTCTTTTGCATCTTTTAAATACTAATAATCCCATTTGGAATAAGAAAGAGGAAGAGC
TTAGAAGAACAGCTCCATCACCTTGGTCACCTGCAGCTAGTCTCAAAGCAGTGATAATAGCGATACACATCAAA
GTGGAGGTAGTGACATTGAAATGGATGAGCAACTTATTAATAGAACCACACATGTGCAACACGACTTTCAGACA
CAGAGGAATCCATGCGAGGGAAGTCTGACGAACTGCCAACAGTGGTGAAGATGGAAGCAGTGGTCTCTGGTAG
CAGTAGTGGGCATAGTGATGGATCTAGCAATGAGGTTAATCTAGCCACGCAAGCCAGTCAAGTGGAGCCCTG
GCAGTGAGGTACAGTCAGAGACATTGCAGATATTGAAGCCCTCAAAGAGGGAAGATGAAGACGATGATCATGGT
CATAATCTCCCAAAGCAGTTGTGGTACAGATCTTCGGAATAGAAAGTTAGAGAGTCAAGCAGGCAATTTGCCTG
GGGACTCCCAAGGCACGTGAGAAAGAAATGGACAAAGCAGGGAACAGGAAAGGACCTGGTTTTTAACACTG
AATCATGCCATCAGTAGATAATCGAATGCGAATGCTGGATGCTTGTTCCACTCTGAAGACCCAGAACATGATAT
TTCAGGGGAAATGAATGCTACTCATATAGCACAAAGGCTCAGGAGTCTTGATCACACGAACCTGGGACTTCTCT
TGGGAGACTATTGGGAATGAATATTATTTAATTGTCGACAAATTTATGGTCCACAGCATCACACCACCACCA
CCATCACCCACCACGATGGGCATATGGTTGATGATATGCTAAGTGCAGATGATGTCAGTTGTAGTAGCTCCCA
GGTTAGTGCAAAATCAGAAAAAATATGGCTGATTTGATGGTGAAGAAATCTGGATGTGAAGAGGAGCTAGTTCA
GATTAATTCACATCGGGAACCTGACATCTCACCTCCAACACATCTTCCCAATTTAGCTTCCATTTACCATGAACAT
CTTAGTCAAGGACCTGTAGTTTCAATAACATCAATTCACAGTAATGCTGTTACAGACATTAATTTGGATAATGTTT
CAAGAAAGGAAATACTTTGTTGGGATATAGTCCAAAGATGAAGATGCAGTTAATCTTCTGAAGGATTAATAAT
GAAGCAGAGAAACTCTTTGTTGTTAGTATGTTGGTTTACAGATAGACAAATTCGAATGAGATTCATTGAAGGTT
GCCTTGAAACTTGGGAACAACAGATCAGTAGTAATTTCACTTCGCTCTTCCAAACTATTTGGTACTTTTCAG
CAGTTTGGAGCAGTTACGATACACACTGGATAACAATGTGGGCAGAAAAAGAACTGAACATGATGAAGCTTTTC
TTTGATAATTTGGTACTACATTCAAACCTGTGAGAGAGGAGAACAAACATGCACGTACAGCCATAGTGCTG
AAGTTCAAGTTCGCTTCAATCTTGACTTGTTAGTAGAAGATTCTGAAATGTTATGATGATGCACCTCCATTGTTTTAA
GAGCAAGTTGACATCTTATGGCATTGTTAGTAGAAGATTCTGAAATGTTATGATGATGCACCTCCATTGTTTTAA
TCAAGTTGGAAGTAAAGATCAACATGCTATGGGTATGGAACCTACAACATCTTTCTGGAGAAGATGCCCCA
GCTAAAACCTGAAACAAATTAGCATGACTGGCTTAAACCTGTTTCAGCATCTCTGTAACCTGGCTCGATTGGCTACC
AGTGCCCTATGATGGTTGTTCAAATCTGAGCTGTGGTATGGACCAATTTTGGGCATTGCTTTAAGAGCACAA
TCTGGTGATGTCAGTCGAGCAGCTATCCAGTATATTAACTCCTATTATTAATGTTAAACAGGTTTGGAGAAG
AGCAAGAATTTATTAGTAAGTGCATGGAGAGTCTTATGATAGCTTCTAGCAGTCTTGAACAGGAATCACACTCAAG
TCTCATGTTATAGAAGAGGACTCCTTATGCTGAAGACACATCTGGAAGCGTTTAGGAGAAGGTTTGCAATCA
TCTGAGACAGTGGCAAATTTGAAGGCACCTGTTAGTAGTCAATTTGAAGGCACTGAGTGACAAACAGTCTCTGCC
GCTAAGGGTTGTATGCCAGCCAGCTGGACTTCCTGACAAGATGACTATTGAAATGTATCTCTAGTGACCAAGGTAGC

FIG. 1E

AGATCTTAGGGCTGAAGTAACTCATTGGTATGAAAATTTACAGAAAGAACAAATAAATCAACAAGCTCAGCTTCAG
GAGTTTGGTCAAGCAACCGAAAGGAGAGTTTCTGGAGGCTCATGGACCTGTGAGGATGATTTTCATCTGG
ACACGAGTTAACAAACAGATTATGATGAAAAGCACCTTCATGAGCTTGTTTTAAGGATATGCAGATGGTATTTGTA
TCTTTGGGTGCACCAAGGAGAGAGCGGAAAGGTTTCAGCTGCCAGCATCTTGCCTCCACCCCTC
AGAAGGACAACATTCCAATGCTTTTGTCTTTACAGAGCCTCATTTAACTACTCTTTTGTATTTAGAGATGCTT
GCATCATTTAAACCACCTCAGGAAAAGTGCGAGTGATGATAGTGAGAGCTTACGATGTGAAGAATTCATCTT
CATGCAGAAAATCTGTCTAGCGGCTGGAGCTACTGATGCTTCTTCCATCATGTCTTAATGTTGATGGCA
TTCCAGAAATATCTCAGATGAGCAGAGTTTTAAAGCTCAGTCTGATCACAGGCTAGACATGAAGTTTCACATTAT
CAATGTGGCTCTTGGTGAAGTTGGCTCATTGCTTCTTCTAGTGAATCTAGCCTTGCTGATAGCGATCATTTACA
AGATTGGCTAAGAAATTGACTCTCCTTATTCCTGAGACTGCAGTTCTGATCAATCAATCGTTCTTTCTGCTATTGGCTGCCCTCAACATTATTGA
TTATCCCTGTCAGGGCTGGATGGAGGAGACTCAATCAATCGTTCTTTCTGCTATTGGCTGCCCTCAACATTATTGA
AATTTCTCTGATGCTCAAGCACTCAAACTATTAGGATAGATGATTAGGAAAGAACCAATATTAAACCAGG
ATGTAAGAGTATTTTGGTTTATGCAAAATTAGTTGACAACATACATATAAAGGACGCTAGTCAGACAACGCTC
CTCGACTTAGATGCCTTGGCAAGACATTTGGCTGACTGATTCGAAGTAGGGAGATCCTTGATCATCAGGATGGT
AATGTAGAAGATGATGGGCTTACAGGACTCCTAAGGCTTGCAACAAGTGTGTTAAACACAACCAACCCCTTTAAAT
TTTCAAGGGAAGCACAGGAATTTTGAGAGATATCTTCAATCTCCTGTTTTGTTGCCAAGTCTAAAGGACCCGACA
ACAGCCAAAGTGCAAAATCACATTTCTTCAAGAGCTGCCGCTTACGATTTGTTAGTAGAGATGGTAAAGGGTCTGT
TGAGAACTACAGGCTAATACACAACCTGGTTATGGCACACACATGCAGTCCCATGCCCTTATAAATGGGATTA
CTGGCCTCATGAAGATGTCGCTGAATGTAGATTTGTTGGCTTACTAACCTTGAGCTACTTGTACTTAGCT
TCTACTATTAGCACTTTATATGATACCTGAGGCAAGACAGGCTGCTTCACTGCCAAGTATTCAGAGGATATGA
AGCACAAAGACCCTCTTCTGGAGCTTCAGAAAATGTTTACATATTTAATGGAGATGAATGCAAGCATATAATCC
TAGACCTTTCTGTAAACATACACCATGGATAAGCAGCCTCTGAATACTGGGAAACAGAAAGATATGACAGAGTT
TTTTACTGATCTAATTACCAAAATCGAAGAAATGTCTCCGAACTGAAAAATACCGTCAAAAGTTTATTGGAGGT
GTAATTACAAACAATGTTGTATCTTGGATTGTGAACATGTTAGTCAAACTGCTGAAGAGTTTTTACTGTGAGGT
GCCAAGTGGCTGATATGAAGAACAATTTATGAATCTCTTGATGAAGTTACTATAAAGACACCTTTGGAAGGTGATAA
CATGTATACTTGTCTCAATGTGGGAAGAAAGTACGAGCTGAAAAAAGGGCATGTTTTAAGAAATTCCTCGCAT
TTNAGTTTCAATACTATGAGATACACATTTAATATGGTCACGATGATGAAGAGAAAGTGAATACACACTTTTTCCT
CCCATACGTTTGGACATGACGCCCTATACAGAAGATTTTCTTATGGAAAGAGTGAGAGGAAAGGTTTTAA
AGAAGTCAGTGATCATTTCAAAAGACTCAGAGAGCTATGAATATGACTTGATAGGAGTGACTGTTACACAGGAAC
GGCAGATGGTGACACTATTATAGCTTTATCAGAGATATAGTAAATCCCCATGCTTATAAACAATAAATGAT
CTTTTAAATGATGCTGAGGTAAACCTTTTGATCTGCTCAACTTGCAATGTTTGGTGAGAGATGACGA
CCAAGACCTATGATCTGTTACAGATAAATTTATGGACTTCTCTTTGAAAGACACACAGTGCTATATGCTGTTT

FIG. 1F

TACAAACGCATGGAACACAGAGGAAGAAATGGCAGAGAATACAAATTTGATGTTTCGTCAGAGTTACTAGAGTGG
ATTTGGCATGATAACATGCAGTTTCTTCAAGACAAACAAATTTTGACATACATAATTTTGATTATGTGGCAATT
GTGTAGTTGATTTCCAGTACATTACAGATCCTAAAGCTGTGCTTAAATGACAGCAAAAGTTAAGCACCTTCCTTT
GTCTAGAGACATTTATTCATTTCTAAAGAAAGCCACGATGCTTCAGTGGATTGAAGTGTGACGAAACAGTTTA
ATAATAGTCAGGCAGCTTGTGAGTGGTTTTAGATCGTATGGCTGATGACGACTGGTGGCCAAATGCAGATACTAA
TTAAGTGCCCTAATCAAAATTGTGAGACAGATGTTTCAGCGTTTGTGTATCCATGTGATTCAGAGGCTGAGACCTGT
GCATGCTCATCTCTATTTGCAGCCAGGAATGGAAGATGGGTGAGATGATATGGATACCTCAGTAGAAGATATTGG
TGGTCGTTTCATGTCTACCTCGCTTGTGAGAACCTGTATTAAATTTGGAACATGGTGTAAACCTCACAGTAA
CATCTTACAGAGTATTTTGGCTTCTTACGAATTTGCAAAATGGGTGAAGAGAGAGCCAAATTTTGTCTTTCATT
GCAAGCTATATCTACAATGTTACATTTTACATGGGAACAAAGGACCTGAAATCCCTCAAGTTGAAGTGTATCA
GAGGAAGAGGGGAGAGAGAGAGGAGGAAGATATCCTCTCTCTGGCAGAGAAATACAGGCCAGCTG
CCCTTGAAGAGATGATAGCTTTAGTTGCTCTTTGGTTGAACAGTCTCGATCAGAAAGGCATTTGACATTTATCACA
GACTGACATGGCAGCATTAACAGGAGGAAGGATTTCCCTTCTTGTTCACACATATTCGTGATGGCATCAATATA
AGACAAACTTGTATCTGATTTTCAGCTGTGTCGATACAAATATCGACTTGCAGAACATATTTGATCTATGCTTTT
CACATCAATAGCAAGTTGACTCCTGAGGAGCCAAATCCTTTCTTTAAGTTGTTGACTATGCTAATGGAGTTTGCT
GGTGACCTCCAGGAATGCCCTCCCTTTCATCTTATTTCTGCAGAGGATATGGAGGTGATTTGAATACAAATCCT
TCTCAGTGTCTAGATTGGTTGGCAGTGCAGACACCCCGAAATAAAGTGGCACACAGCTGGTCTTACAGAAATATG
GAAACTGGGTCGAGCGGTTCTTTGGCTCACAAATATCCTAGAGTGAGGACTTCTGCAGCTTATCTTCTGGTG
TCCCTTATACCAAGCAATTCATCCGTCAGATGTTCCGGTCAACAAGGCTTTGCACATCCCAACCCGTGACCTT
CCACTCAGTCCAGACACACAGTAGTCTACATCAGGCTACACAGTGTCTCTGGTTTGTCTCAAGAGCCAAA
CTTTATGTTGATGCTGCTGTTTCATGGCACTACAAAGCTAGTGCCCTATTTAGCTTTATGACTTACTGTTTAAATTC
CAAACTGAGAAGCTGATGTTTCCACATATTTTCATGGATTTGTGGAACCTTTTCCAGCCTAAACTTTCTGAGCCA
GCAATAGCTACAAATCACAATAAACAGGCTTTGCTTTCAATTTGGTACAATGTCTGTGCTGACTGTCCAGAGAATA
TCCGCCCTTATTGTTTCAGAACCCAGTGGTAACCAAGAACATTGCCCTCAATTACATCCTTGTGCTGACCATGATGATCA
GGATGTGGTCTTTTAAACCGTGGATGCTGCCAGCGTACTATGGCATTTCTGAGGCTCTGCTGTGAGCAGTCTC
CTGCATTCACACGACAACTGGCTTCTCACAGAACATCCAGTGGCCCTTAAAGAAATCTTACACCCACATGCCAGCC
AATACCTTGGAGCAGTAGAAGAACTGTTTAACTGATGCAGCTGTTTATAGCTCAGAGGCCAGATATGAGAGAAG
AAGAAATTAGAAGATATTAAACAGTTCAAGAAACAAACCAATAGTTGTTACTTACGTTGCTTAGATGGCCGCTCCTG
CTGGACTACTTTAATAAGTGCCTTCAGAAATCTATTAGAATCTGATGAAGACAGACTTCTTGTGTTAATTCGAG
GATTGATTTCAATGACAGAGTCTTTCAACACTTTCACATGATGTATCACGAAGCTACAGCTTGCCATGTGACTGG
AGATTTAGTAGAACTTCTGTCAATATTTCTTTCGGTTTTGAAGTCTACAGCCCTTATCTTCAGAGAAAGATGTGA
AACAGCATTAATCCAGTGGCAGGAGCGAATTGAATTTGCCCATAAACTGTTAACTCTTCTTAATTCCTATAGTCC

FIG. 1G

TCCAGAACTTAGAAATGCCTGTATAGATGTCTCAAGGAACCTGTACTTTTGAGTCCCCATGATTTCTTCATACT
 CTGGTCCCTTTCTACAACACAACCATTTGTACTTACCATCACAGTAATATACCAATGTCTCTTGGACCTTATTTCCC
 TTGTCGAGAAAATATCAAGCTAATAGGAGGAAAGCAATATTCGGCTCCGGCCCTGAACCTCAATATGTGCCT
 CTTGCCACAATGGTGGAAACCAGTAAGGGCAAGATGACGTTTATGATCGTATGCTGCTAGACTACTTCTTTTC
 TTATCATCAGTTTCATCCATCTATTATGCCGAGTTGCAATCAACTGTGAAAAATTTACTGAACATTAGTTAAGCTGA
 GTGTCCTAGTTGCCTATGAAGGTTTGGCACCTTCATCTTGCACTGTTCGCCAACTTTGGACTGAGCTATGCCAGA
 CTCAGTCTGCTATGTCAAAAACCTGCATCAAGCTTTTGTGTGAAGATCCTGTTTTCGAGAAATATAATTAATGTATC
 CTAATGGATGAAGAACTTTTAAACAACAACATTTGTCTACACGTTTCATGACACATTTCTTCTAAAGTTCAAAG
 TCAAGTGTCTTCTGAAGCAAACTGTGCCAATTTGATCAGCACTCTTATTACAACTTGATAAGCCAGTATCAGAAC
 CTACAGTCTGATTTCTCCAACCGAGTTGAAATTTCCAAGCAAGTGCTTCTTTAAATGGGACCTGAGGGCACTC
 GCTTGTCTCCTGTCACTACACACTCCCAACAGTTAAACCCAGCTCTAATTCCAACTCTGCAAGAGCTTTTAAAGCA
 AATGCAGGACTTGTCTGCAACAGAGAACTCACTCCAAGAGCAAGAACCCAAAGAAAGAACTAAAGATGATG
 AAGGAGCAACTCCCATTAAGGCGCGTGTAGCAGTGATGAGGAGCACACTGTAGACAGCTGCATCAGTGAC
 ATGAAAACAGAAACCCAGGAGGTCTGACCCCAACGAGCACTTCTGACAAATGAGACCAGAGACTCCTCAATTAT
 GATCCAGGAACTGAGCAAGATCTTCTTCCCTGAAAATAGTTCTGTTAAAGAAATACCGAATGGAAGTTCCATCTT
 CGTTTTCAGAAACATGTCAATATCAGGTCAAGCATGCAGAAAGCAAGTCCAAACAATGGTAGATATGACGATT
 GTAAAGAAATTAAGACCTCCACTGTTCCAAGATTCTACCCCTAGCTGAGGAAGAACTGAGTTCCCTTCTACTTC
 TATCTCTGCAGTTCTGTCTGACTTAGCTGACTTGAGAAGCTGTGATGGCCAAAGCTTTGCCCTCCAGGACCCCTGA
 GGTGCTTTATCTCTCAGTTGTGGCCATTCAGAGGACTCTTTAGTCAATATGCAAGCAACATGACATTTTAGATACC
 CTGTGTAGGACCATTGAATCTACAAATCCATGTCTCACAAAGGATATCTGGCAAGGAAACCAAGCTGCTTCTTGA

>SGPR430_SEQID_6

ATGTCTCCTCTGAAGATACATGGTCTCTATCAGAAATTCGAAGTATGCAGACTGGGATTACAAAGTGGAAGAGGA
 TCCTTTGAAATTGTAGAAAAGAGAAATAAAGTCAGCCTAGTAGTTCACTACAATACCTGGAGGAAATCCAAGGATAT
 TTCAGCTAAGTCAATACATTAAAAATGTGGTCTTCGACCCAGTGGAGCGAAACAAGCCGCTAATGTTAACTC
 TGCAAGATAACAGCTTCTTGCTATTGACAAAGTACCAAGTAAGGATGCAGAGGAAATGAGGTTGTTCTAGATG
 CAGTCCATCAAAACAGACTTCTGTCAGCCATGAAACCGTCTCAGGGGTCTGGTAGTTTGGAGCCATTCTGGGC
 AGCAGGACCTCACAGAAGGAAACCCAGCAGGCGACTTCTTACTCAGACAAATCAGGCTTCTGCAAAAAGAGGAAG
 TTTGGAAACTAAAGATGATATTCATTTGAAAAGTTCTTGTAATCCGGGTAGAGGATCGATTAAGACTGTAGCA
 GGAAGTGGAATAGCTCGGACGATTCTTCTTTGACATCTACTTCAACACCTCTTAGATCAGGGTTGCTAGAAAAT
 CGTACTGAAAAGAGGAAAGAAATGATATCAACTGGCTCAGAAATGAATGAAGATTACCCCTAAGGAAAATGATTAT
 CATCGAACACAAGGCCATGACAGATCCCTCCAGAAAAGTATTTAACCAGCAGTAGAGAAAGCAGCTGAGTTTGA

FIG. 1H

AACAGTCAGAAGAAATAGGACATCAGGTGGGCTTTACCTTTACAGTCATCATCCTTTTATGGTAGCAGAGCTG
GATCCAAGGAACACTCTTCTGGTGCACTAACTTAGACAGGACTAATGTTTCAAGCCAGACTCCCTCTGCCAAA
GAAGTTGGGATTTCTCTCAGCCAGTTCCTCTTCTGTTAAAACTGAGGTGTAACCAAGATTACACTGGCTG
GAATAACCAAGAGTGCCCTTTCTCTCACCAACAGCAGCAACTGCAGGCTTCTCCAATTTGGGAAATACCTG
CTATATGAATGCTATTCTACAATCTCTATTTCACTCCAGTCATTTGCAATGACTTGCTTAAACAAGGTATCCCAT
GGAAGAAAATCCACTCAATGCACCTATCAGACGCTTTGCACACTTGCTTGTAAAAAGATATCTGTAAATTCAGA
GACCAAAAAGGATTTACTCAAGAAGGTTAAAAATGCCATTTTCAGCTACAGCAGAGAGATTCTCTGGTTATATGCAG
AATGATGCTCATGAATTTTAAAGTCAGTGTGACCCAGCTGAAGAAGATATGGAAAAATTAATAAAACTTTGGA
AGACTGAACCTGTTCTGGAGAAGAAAATTCACCAGATAATTCAGCTACCAGAGCATACACTTGCCCTGTTATTAC
TAATTTGGAGTTTGAGGTTACGACTCCATCATTTGTAAAGCATGTGGAGAGATTATCCCAAAAGAGAACAGTTT
AATGACCTCTCTATTGACCTTCCCTAGGAAAAAACCACTCCCTCCCTCGTTCAATTCAGATTCTCTTGATCTTTT
CTTTAGGGCCGAAGAACTGGAGTATTTCTGTGAGAAAGTGTGGGGAAGTGTCTCTTGTCAAGCACAAATTTAA
CAGGCTTCTAGGCTCTCATTTCTCCATTTGAACGATATAGCTTCAATGTGGCTCTCTCGCTTAAACAATAAGATT
GGGCAGCAAGTCATCATTTCCAAGATACCTGACCCCTGTCATCTCATTTGCACCTGAAAAATACAAAACCACTTTTACCC
TTGGTTGGAGTGACATATGGCAATGTCTAGACCATTTGAAAGCCCTCTCAATGGTGAATTCCTGCATCACAGCC
CTTCTACACCTTCAAGAAATTCACCTTCAATCCAAAGAGCTCTTGGCTTTATGCCCTTATGCTTCAAGACAGTGAGGA
TGAGCTAAAACGTTCTGTGGCCCTCAGCCAGAGACTTTGTGAATGTGAGCAAGTCTGAATGGAAAACTCAGGATTTGACAGAAATGA
TGGAAGAAGATTCAAAATTTATGCCCAATAGAGCCCTGACAAAGTCTGAATGGAAATGTGAGAGATGCTTCAACCATCTCTGAGTCATGAAGATG
GCGAAGAAGAGCTTCTAGCAGCTGTCTTGGAGATAAGTAAGAGAGATGCTTCAAGAAATGCCAGAAATCCAGACA
ATGATAAGCCCACTAGCAGCCAGATACCGGATTTGCAAGAGATGATATTTCAAGAAATGCCAGAAATCCAGACA
CTATGGAACTGAGAAGCCCAAAACAATCAGAGAGCTGGATCCTGCCAGTTTACTGAGATAACTAAAGACTGTG
ATGAGATAAAGAAACAACAACTCCAAAGGATCTCAGGGAGAAAGTTGATTGGCTCCAGCAGTATGATATGGAGC
GTGAAAGGGAAGCAAGAGCTTCAAGAGGCACTGGCTCAGAGCCTTCAAGAGCAAGAGGCTTGGGAACAGAA
AGAAGATGATGACCTCAAAAGAGCTACCGAGTTAAGTCTTCAAGAGTTTAAACAACCTCTTGTGGATGCATTGGG
TTCTGATGAGGACTCTGGAATGAGGATGTTTTGATATGGAGTACACAGAGCTGAAGCTGAGGAACCTGAAAG
AAATGCTGAGACAGGAATCTGCCCTCATTCGTACCGGCTCATCAGTGTGTCACTCAGTCACTTGTGAGCACTTCTTC
TTCAGGTCATTACATTAGTGATGTATATGACATTAAAGCAAGCGGTGTTTACTTACATGACCTGGAGGTATCA
AAAATCCAAGAGGCTGCCGTGCAGAGTGATCGAGATCGGAGTGGCTACATCTCTTTTATATGCACAAGGAGATC
TTTGATGAGCTGCTGGAACACAGAAAAAGAACTCTCAGTCACTTAGCACGGAAAGTGGGAAGACTACCCGTCAGGC
CTCGTGA

FIG. 11

>SGPR496_1_SEQID_7
ATGACACTACTTGCCTCCCTGGTACACAGGCCCCATGATCCCCATGGATGTTAATGAGCCCCAGCTCCGTGACCCAC
GGCTCCTACCCCTCAGCTCTAGCCTGAGCATATCTCCTCATTCTCGCCACTGGTAAGAACTTTCCCTCCATTT
TGGTCATCCACGTGAGTGAAGTCAACAGGATTGATGACAAAATAGAGAGGATTGGAAGACAGTGAGCCAG
GTGCCAACTCTTCAATAATGATGAGTCTGTTGCTGCTGCAAAAACGGGGCCAGTGAACATTACATCAGTGT
GTGTGAGTCCCAGGACCTTACAAATATCAGTTTTGTGTTATCAGAGAAATACGAGGTTATTGTTAAATTTGAATC
GGATGAATTACCTTTTGGTGAATTGGTTCTAATATTGGTGATGCACATTTTCAAGAAATTCAGGGCTGGAATCTCC
TGGAGCCTGTGGTAGATCCTGATGACCCCATCTCAGTTCCTGATTGCTGCAGCAGCAGCAGCAGGAT
TCCTTCAGTGAGTGTGCTAGTTGCAGTTCTCTGTTGCAGGCCACAAAGGCGAGGCAATTTATTGAAGGATGCT
GGGTGCTTCAAGGAATTGAAGCAAGAGCTGACTCAGGAAGGCCGGGGGACACCCAGGTCTGCGTG
GCCCCCGCCGACGCCAGTGCCGCCGAGCCCTGCAGCAGGGGAGGAGTGGAGCGGAGCGGAAAGGTGGA
CGAGGAGGTAGAGGCGGAGACGGCGGAGAGCGGAGAGGAGGAGTGGAGCGGAGCGGAGCGGAAAGGTGGA
GGGGAAGGCGGAGCGGCGGAGGCGGAGGCGGAGGCGGAGGAGTGGAGCGGAGGAGTGGAGCGGAGGAGG
GCGGGGAAGGTGACGCCCTGGGAAGTGGAGACGCGGAGGTCGCGGCCCGGCTGAGCTGAGCTCAAGCT
GGAGCCGAACCCGAGCCGTCGCGGAGCGGAGCAGGAGCCGAAGCAGGAGCTGGAGGATGAGAACCAG
CGCGAGCGCGGTGGCGCAACAGCAGGAGTCTCTCCCCCACCCTTCCCTCCGATCCACCGGCCCC
CCGATCCCTCTCCGCTCGCAGTCGTGCGCGCGCGCCGACCCCGGCCCGGCCCGGCGGCGGCTCCGTA
CCCCCGCAGCCTAGCCCCCGGCCCGGCCCGGCCCGGCCCGGCCCGGCCCGGCCCGGCCCGGCCCGGCCG
GATGGGATTTTGGCGTGGGCCACAGGCTGTTCTACGTGAACAGCTTTAAGGTGGAGAGAACTGGAGGC
AGGAAGTGGGTTATCTACAGTGTCTCGTGTGTGTGGAAACCCAGAGACGAGAAAGGCAAGTCC
TGCATCTGCCATGTGTGGCACCCTCTGAACAGACTCCACTTTGCCCTTCTGTGTCTTTGGCTGCTTC
ACGGAGAAACACATTACGAGCAGCAGAGAGAGAAACACAACTTAGCAGTAGACCTGTATTACGGAGGTAT
ATACTGCTTTATGTGAAGGACTATGTATATGACAAAGACATTGAGCAATTTGCCAAAGAGAGAGAGAGC
TTTGAATTACAAGCTCCACCTCAACAGAGTTTCTACCCAGCAGTGTTCAGTGCCAGGCCCTGGTGAGAAAT
CCCAACCTGGGAAACAAACCAACAGAAATTAGAAGTGTGGGCAACACCCGAGGAGAGAAATCACCTCCA
GCTTACGATCGTTAAGAGGACTCATCAATCTTGGCAACACGTGCTTTATGAAGTGTGTCAGGCCCTCA
CCACACGCCGATAGAGATTCTTCTCTGACAGGACCGATGTGAGATGCCGAGTCCCGAGTTGTGT
CTGGTCTGTGAGATGTGCTGCTGTTTCGGGAGTTGATTCTGGAACCCGCTCTCTCATGTGCCCTATAAGTTA
CTGCACCTGTTGGATACATGCCCGCATTTAGCAGGTTACAGGCAACAGGATGCCCCACGAGTTCCTCATTGC
AGCGTTAGATGCTGACAGGCACTGCAAGGTGATGTGCGGAAGCGGCGCAACAATCCCAACCACTGTA
ACTGCATCATAGACCAAACTTTCACAGGTGGCCTGCAGTCTGATGTACACCTGTCAAGCCTGCCATGGCGTCTCCA

FIG. 1J

CCACGATAGACCCATGCTGGGACATTAGTTGGACTTGCCTGGCTCTTGACACCTCCTTCTGGCCCATGAGCCCA
GGGAGGAGAGCAGTGTGAACGGGAAAGCCACATACCAGGAATCACCAACCTCACGGACTGCTTGGCGAGGT
TTACGAGGCCAGAGCACTTAGGAAGCAGTGCCAAAATCAAATGTGGTAGTTGCCAAAGCTACCAGGAATCTACC
AAACAGCTACAATGAATAAATTACCTGTCGTTGCCCTGTTTCATTTCAAACGGTTTGAACATTCAGCGAAACAGA
GGCGAAGATCACTACATACATTTCCCTTCCCTCGAGCTGGATATGACGCCGTTTATGGCCTCAAGTAAAGAGA
GCAGAAATGAATGGACAATTGCAGCTGCCAACCAATAGTGGAAACAACGAATAAGTATTCTTGTGCTGTGG
TTAATCACCAAGGAACCTTGGAGAGTGGCCACTATACAGCTTCATCCGGCACCAAGGACCGAGTGGTTCAAG
TGTGATGATGCCGTCACTAAGGCCAGTATTAAAGACGTACTGGACAGTGAAGGTAATTTACTGTCTATCAC
AAACAGGTGCTAGAACATGAGTCAGAAAGTGAAAGAAATGAACACACACAAAGCCTACTGA

>SGPR495_SEQID_8

ATCGGGTGAAAGATCCAACATAAGCTTTACCTGAGAAAGCCAAAGAAAGTAAAGGCCCTACTGTACCTCATGAT
GAAGACTCTTCAGATGATATTGCTGTAGGTTAACTTGCCAACATGTAAGTCATGCTATCAGCGTGAATCATGTAA
AGAGAGCAATAGCTGAGAATCTGTGGTCAGTTTCTCAGAAATGTTAGAAAGAAAGAAATCTATGATGGCAGC
TAGTACTTACTTCTGATATTGGTTGTGCTCAAGTGTGGCTTCCAGGGATGTGTTAAACTCAGAAAGCCAAAC
ATTCAATTGAAGCACTTAAAGATTCCAGAACAGAGCCCCATTGTATTATAATTAATCTGAGCACATGGATTATATG
GTGTTATGAATGTGATGAAATAATATCAACGCATTGTAATAAGAAAGTTTTGGCTCAGATAGTTGATTTCTCCAG
AAACATGCTTCTAAACACAAACAAGTGCAATTTCTAGAATCATGAACCTTTGTGAAGAAATGTGAACACAGATG
AAATACAGAAGGGAGGAAATGCAGAAATTTATCTGTAAGAGGAATTAACAAATTTAGGAATACCTTTTCTTTTAAAT
GCAGTCATGCAGAACTTGGCACAGACTTATCTCTTACTGATCTGATGAATGAGATCAAGAAAGTAGTACAAAAC
TCAAGATTTTCCCTCAGACTCTCAGCTGGACCCATTGGTGTGGAACCTTCAAGGCTGGACCACTGACCT
CAGCCTTGTTCCTGTTCTTACAGCATGAAGGAGACTGAAAGGACCACTTCTCCTAAAGTTCTTTTAAATCA
GCTTTGTCAGAGGCACCTCGATTAAAGATTTCCAGCAACAGGACAGTCAAGGACTTCTTCAATTATCTTCTGGAT
GCAGTGAGGACAGAAACAAAGCGAATACAAAGCTAGCACTTCTAAAGCATTTAACAACCCAACTACTAAACT
GCTGATGATGAAACTAGAAAAAAGTCAAGATCTCCACGGTGAAGATCCATTGATATTGATATTTCACTTCTATAA
TAGAAGAAAGGGTTCAAAACCTTACTTTGGGAAGAAATGAATAATATAGAAGTTTACGGGAGACAGATCATGA
TCGATACAGTGGCAATGTTACTATAGAAAATATTCAACACCTAGAGCTGCCAAGAACATTTCTCATCTAAAGAT
AAGAGTCAACTAATTCATGACCGAAAATGTATTAGAAAATTTGCATCTGGAGAACTGTACATACAGAAAAATG
AAAACTTGAATGAATGGGATTCTTTAATGTTTCCAGCCTCATGAATCTGAGTCACGCTGATGAATGAAGCCC
TACTGATGACAGTGAAGAAAGCCAGCCATTCTGAAAGCAATGTTGATGCTGACAGTGAGCCTTCAGAACTCTGA
AAGTGCTTCAAAGCAGACTGGGCTGTTCCAGATCCAGTAGTGATCCGGTGTGCAGCCAGATGGACCCCTTTTACC
CTCTGTCAGCAGGTAACTGCTGTACACCAAGGAGACTGACAGTGGTGATAAGGAATGCGAGAAGCTATTTCT

FIG. 1K

GAACCTCGTTTGAGCAGCACTGTAACCTGGGATCAAGATTTTGACAGAGAAAAATCAGCCACTAAATATTTCAAATA
ATTTATGTTTTTAGAGGGGAAGCAATTTGAGGCTTATAGTCCCCAAAATGCTTTTCAGACCCCTTTCTCAGAGCTA
TATAACTACTTCTAAAGAAATGTTCAATTCAGTCCTGTCTCTACCAAGTTTACATCTATGGAATTAATAATGGGAATA
ATAAGCTTCTATGTGAGAAATGTACTAAAAACAACAGAAAGTACCAAGAGAAACCAAGTTTGCAGAAAAGAAAGT
AGAAGGAGTTTATACTAATGCCAGGAAGCAATTTGCTCATTTCTGCTGTCCAGCTGTCTTAATCTCCACCTGAAA
AGATTTATCAGGCTGGCTTGAGTCTTGTAAGTAACAGACATGTAGATTTTCCACTTATGCTCGATTTAGCAC
CATTCTGCTCTACTTGTAGAAATGCAAGTGTGGAGATAAAGTTCTCTACGGTCTCTATGGCATAGTGGAAAC
ATAGTGGCTCGATGAGAGAAGGCCACTACACTGCTTATGTGAAAGTGAGAACACCCCTCCAGGAAATATCGGAA
CATAACACTAAAAGAAAATGTGCCTGTTTGAAGCGGCTGATAGTGAATCAGCAGGCCAGTGGTCCATGTT
AGTGACACTTACTACAGGTGGTTCCAGAATCAAGAGCACTTAGTGACAAAGCCTACCTTCTTTCTATGAAAGAG
TATTATAA

>SGPR407_SEQID_9

ATGGAGTATCCAGTCCCATACCTTAGATCCCCGAACAGGACTCTGATCCCAGAGAGAAATTTGGTCAAAACCCATTA
CTTGCTTTGGTCATCGCATACAAGACTGTGAGTTGGCCAAAGACAGCAGCTGCTTGCAAGCAAGCTAATAAGTGG
ATGCCCTTTGTGATACCGAGCAAAACCTTGCCATGGGACCCCACTGGAAGTCAAGATTTGTTATCAGCAAAATCGC
CCATATCCCTCCCCGACCCATCAAACTTTCCCTACCTTCTACGCTGTGAATGCTTCTCTGCAGCTGCTCTTCT
ATCTCCACAGCCCTCATGGCATAAGCCCGAGGGCTTAAAGCCAGCAGGATACCCAAAGAGTTCTGACATTCCT
TATGGAGCGGCTACACCTTGAAATCAACCAAGGAGCGCGGGCTCCACAGTCCCTGCCCATGGTCCAGC
TCCCTCTCCACCCCAAGGGGAGTGCTCTGTAAAGAGTCTGAGTTAAATGATGCTGACTGGGCCAACCTA
ATGTGGAAGCGTTATCTGGAAGAACAAAGAGGACAGCAAGATGGTGGATCTGTTGTGGCCAGATGAAAAGTTA
TCTCAAGTCCCAGGCTGTGGGTACCACTCTATGACCTTCAAGGTTTTTTTTTTGTGACCTCTCCCTGACCATC
CCCAAGAAAGGATTTGCTGGGGCAAGGTGCTCTGCGGATTTTAAGCCCTTTTACCAGGAAGAGAGCT
AGAGTTAGAGAAATGCCCTCAGGGACTTTGCCAGTGACAAAGTCGGAAGTCTGTCTACCAAGCTGTGCCCCCTTG
GAACCACTCAGGCAGCATCCACTGTGCCACTACACAAACCTGTGCCAGTCCAGACTGGTTGGCACGTTTACA
ATGACTCTTGCTCCCTAAACACGCTGCGGGACACAGAAAGGAATAGAACTCACAGTTATGAAGGCTCTAGTT
CTAGACATTCGTCAAAGCTTCCACAGATATTAATTTATTAATCATGACTCCAGCTCTGGAAACAAATGGAGGA
AGTTACCAGAACCTGGAGGTTTGAAAGAAACATGAAGAGCTGAGACTCAGACCTCTGAAGGAGGAGTACCAT
TGGCTGGTGTGCTGCTGAACTGACAGGAAGTCCCAAGATGGAGGCCAGGAGGCGCTGGCCA
GCTGCAGCTGGTGTCTCCAAAGGCTCACCATGAGGCGGTTATGGGTGTGAGGACAAAGCTGGAACAGGAA
CCAGATGCTGCTGGGGCAAGACCTGTGATAGGTGATACAGTCAGCAACAGCCAGACAACTAGGGACAAG
GCTTGACAGCGGCCACCTTCTCACTCTGCTTACACAGTCCCTCTCTGGGCATGTCTGGATCCTGATCTCTTC

FIG. 1L

TTCTATGGACACAGTCATATTGGATGAAGGCCACCTTAATGACCTCATTTTAAGGAGGGCCCTGTGACACAA
ATGCCCCAGAGTTTTACTGGGGTTTTCTGTGGAGGAACTTGCTGCTTTAGAAATGCTGCCTGATGGACCA
GCACCAAGGACGTTTCTTCAGAAAGAGCTGCTCTTTCCCTGTTCTCTTACATTTTTCATAAGGCAGGTA
AACTCTCCAGCCTGATGCTCATGGATTTCTAGTGAAGAAAGTTTCATGCTCCAACAAGGGCATCGTGTTCATCAT
GGAACCAAGACAGCTGGTGGGAGGGCTCCCTGTCAAACTCCAACCGCTGTGCACTGGGAGGAATGAAC
AGTGGATGGAGCCACAGAGTCTGCACCATTTGCAGCAGGGAAGGCTGCCCCCTCCTCTTCTGTGTGCAAA
CCTGAGATTCAAACCTACGAGTTTACAAATTTGAGGAAGAGCTTTGGTCCAGGGCAGGCTTGGGGAAGAAAAGTG
ACAACCACTCATCTAGGCAGATGCCCTGGGTGCCGTGGGTGGCATGCCAGCATCCATGTAACCTGCCCCAG
AATTGTTGCAGAGTTGACACCTCCAAATTTGCTCAATTTGGTTTCTGAAACACAGTTCAAGTTCACTTCTACT
TCCCTGTCTCAGTTTTCTCAATGATTTCAACCCAGAGGAAGCAATACCTCCTCAATCCCTGCTCCCGGGTTCC
CCAAGGACAAATTCATTTCCCAAGGACAAATTTGTCCCCAAGGACAAATTTGAAGGTGATATTGTCCCTGCTGACA
ATGTATGAACTAGACCGATTATT

>SGPR453_SEQID_10

ATGCTAGCAATGGATACGTGCAAAACATGTTGGCAGCTGCAGCTTGCTCAAGACCATTCAGCCTCAACCTCA
GAAATGGCACTGTGTGGACTGCAACACGACCGAGTCCATTTGGGCTTGCCCTTAGCTGCTCCCATGTTGCCCTGTG
GAAGATATATTGAAGAGCATGCACCTCAAGCACCTTCAAGAAAGCAGTCATCCTGTTGCATTTGGAGGTGAATGAGA
TGTACGTTTTTGTACCTTTGTGATGATTATGTTCTGAATGATAACGCAACTGGAGACCTGAAGTTACTACGACG
TACATTAGTGCCATCAAAGTCAAATTTATCACTGCACAACTGCTAGTGGAGGTTTTACGGTCCATGGGTACA
GGTGATGATTTCTTATTCTTACATGACGGTGCCCAATCTCTGCTTCAAAGTGAAGATCAACTGTACTGCTCTTT
GGCACAGGAGAAGGATACATAATGGGTAAATCTTTGAAACATGTTTGAACAATCACCCATTGGAAGAAAAGC
AAGAAGAACCATTTCAGGAGAAAATAGTAGTAAAGAGAAAGTAAAGAAAGACGGCAGGAATTTGGAGTATCAAG
TTAAGCAGAAATTGAAAGTATGCCCTCCAAGAAAGAGTTTACGTTTACAAGGGCTCGCTCAGTCGACCATATAG
AAATAGTTTCTGTTCCAGGTGCCAGCACAAACGCCAGCATCACCAAGAAAGATAAGTACTCTCTACCTCAGAAA
ATGAAATATCTCAAAGTCAGTGACTCCTCAGTTAACGAAGGCCAATAGTAACCTCTGTTGTAACAGGATTGA
GAAATTTGGAAATACCTTGCTATATGAATTTCTGTTCTCAGGTGTTGAGTCAATTTACTTATTTTCGACAATGTTTT
TAAAGCTTGATCTGAACCAATGGCTGGCTATGACTGTAGCGAGAAGACAAGATCTTTGTAAGCATCCACAGTCA
CAGATACAGTAGTATATCAATGAATGAATGTACGAAAGATACAGGTTTTGTTGCTCCAGACAATCAAGTCT
GTACATCAGGACTAAGTGGAGCATCAAAGGTAGAAAGATGGAACCTTATTCAAGCAAGGAGCCAACTTCACA
GTACATTTCTCTTTGTGATGAATTCATACCTTTGTTCCAGTCAATGTTGTTGAAAGTGGCGTTGGTCTCACCA
TTTGCTATGCTACACTCAGTGTGGAGACTCATTCCTGCTTTCTGTTGTTACGCCCAACAGACGCTCAGGAATTT
CTTTGTGAACTTTATAGATAAAATACAACGTGAATTAGAGACAACCTGGTACCAGTTTACCAGCTCTTATCCCCACTT

FIG. 1M

CTCAAGGAACTCATCAACAAGTTCTGAATGTTGTAATAACATTTTTTCATGGACAACCTTCTTAGTCAGGTTACA
 TGCTTGCAATGTGACAAACAATCAAAATACCATAGAACCTTTCTGGGACTTGTCATGGAGTTTCCAGAAAGGTATC
 AATGCAGTGGAAAAGATATTGCTTCCAGCCATGCTGGTTACTGAAATGTTGGCCAAATTTACAGAAAAGTGAAG
 CTTTAGAAGGAAAATCTACGTATGTGACCAAGTGAACCTCAAGCGTAGAAGGTTTTCTCCAAACCAGTTGTACT
 CACAGAAGCCAGAAACAACCTTATGATATGCCACCTACCTCAGGTTCTCAGACTGCACCTCAAAACGATTACGGTG
 GTCAGGACGTAATAACCGAGAGAGATTGGTGTTCAATGTTGGCTTTGAGGAAATCTTAAACATGGAGCCCTATTG
 CTGACGGAGACCCCTGAAATCCCTCAGACCAGAAATGCTTTATCTATGACTTGTCGCGGTGGTGTATGCACCATG
 GGAAAGGATTTGGCTCAGGGCCTACACTGCCTACTGCTATAATCTGAAGGAGGTTCTGGGTACACTGCAAT
 GATTCCAAACTAAGCATGTGCACTATGGATGAAGTATGCAAGGCTCAAGCTTATATCTTGTTTATACCCAAACGAG
 TTAAGTGAATGGACATTTCTAAACCTTTTGCCCTCCAGAGCTCCTGTTGGGGAGCCCAACATCCCAATGAAGACGCTG
 ATACCTCGTCTAATGAAATCCTTAGCTGA

>SGPR445_SEQID_11

ATGCGGGTGAAAGATCCAACTAAAGCTTTACCTGAGAAAGCCAAAGAAGTAAAGGCCTACTGTACCTCATGAT
 GAAGACTCTTCAGATGATATTGCTGTAGGTTTAACTTGCCAAACATGTAAGTCATGCTATCAGCGTGATCATGTAA
 AGAGAGCAATAGCTGAGAACTCTGTGGTCAGTTTGCTCAGAAATGTTAGAGAAGAAGATTCTATGATGGGCAGC
 TAGTACTTACTCTGATATTGGTTGTGCTCAAGTGTGGCTTCCAGGATGTGTAATAAAGTCAAGAAAGCCAAAC
 ATTCATTGAAGCACTTAAAGATTCCAGAACAGAGCCCACTTGTTATTAATTAATCTGAGCACATGGATTATATG
 GTGTTATGAATGTGATGAAAATTTATCAACGCATTGTAATAAGAAGGTTTGGCTCAGATAGTTGATTTTCTCCAG
 AAACATGCTTCTAAACACAAACAAGTGCAATTTCTAGAAATCATGAACCTTTGTGAAGAAAATGTGAACACAGATG
 AAATACAGAAGGGAGGAAAATGCAGAAATTTATCTGTAAAGAGGAATACAAATTTAGGAAATACCTGCTTTTTTAAT
 GCAGTCATGCAGAACTTGGCACAGACTTATACTCTTACTGATCTGATGAATGAGATCAAGAAGTAGTACAAAAC
 TCAAGATTTTCCCTCCCTCAGACTCTCAGCTGGACCCATTGGTGTGGAACCTTCAAGGCCTGGACCACTGACCT
 CAGCCTTGTTCTGTTCTTCCAGCATGAAGGAGACTGAAAAGGACCACCTTCTCCTAAAGTTCTTTTAAATCA
 GCTTTGTCAGAAGCGGTGCATCTACATTTAATATAA

>SGPR401_1_SEQID_12

ATGACTGTCCGAAACATCGCCTCCATCTGTAATATGGGCACCAATGCTCTGCTCTGGAAAAGACATTGGTCCA
 GAGCAGTTTCCAAATCAATGAACACTATTTCCGATTGGTCAATTTTGGAAACACATGCTACTGTAACTCCGTGCTTC
 AGGCATTGTACTTCTGCCGTCCATTCGGGAGAAATGTTGGCATACAAGGCCAGCAAAAGAAGGAAAGGAAAAC
 TTGCTGACGTGCCTGGCGGACCTTTTCCACAGCATTTGCCACACAGAGAAGGTTGGCGTCATCCCACCAA
 GAAGTTCAATTCAGGCTGAGAAAAGAGAATGATCTCTTTGATAACTACATGCAGCAGGATGCTCATGAATTTTAA

FIG. 1N

AATTATTGCTAAACACTATTGGGACATCCTTCAGGAGGAGAGAAACAGGAAACAAATGGAATTAATAA
ATGGCAACATGAACGAACTCGGAAATAATAAACAGAACTCACCTGGTCCATGAGATTTTCAGGGAACGC
TTACCAATGAACTCGATGCTTGAAGTGAAGTGTAGTAAAGATGAAGATTTCTTGACCTTTCTGTTGA
TGTGGAGCAGAAACATCCATTACCCACTGTCTAAGAGACTTCAGCAACACAGAAACACTGTGTAGTGAACAAA
ATATTATTGTGAACATGCTGCAGCAACAAAGAGCCAGAAAGGATGAGGTAAAGCTGCCCATGGTCTT
GGCCCTGCACCTAAAGCGGTTCAAGTACATGGAGCAGCTGCCAGATACACCAAGCTGTCTTACCGTGTGGTCT
TCCCTCTGGAACTCCGGCTCTTCAACACCTCCAGTGATGCAGTGAACCTGGACCGCATGTATGACTTGGTTGCG
GTGGTCGTTCACTGTGGCAGTGGTCCCTAATCGTGGCATTTATCACTATTGTGAAAGTCAACGGCTTCTGGCTT
TTGTTTGATGATGACATTGTAGAGAAAATAGATGCTCAAGCTATTGAAGAAATCTATGGCCTGACGTCAGATATAT
CAAAAATTCAGAACTCTGGATATATTTTATTCTATCAGTCAAGAGATTA

>SGPR408_SEQID_13

ATGGTCCCCGGGAGAGAACCAACTGGTCCCGAAAGAGGCACCACTGGATCATACCAGTGACAAGTCACTTCT
CGACGCTAATTTTGAGCCAGGAAAGAACTTTCTGCATTTGACAGATAAGATGGTGAACAACCTCAATACT
GCTGGAGGATTCAGTGCTGGGGAAGACAGTGTTCATGACAGGTTTATAGTCCGCTTCCAAGAGAAGGTTCTG
TGGTCTACCAGTGATTATGTGACGCCAAAGCTACTCTACTCATCTATTTGAATAAATCAGAAACTGGATATGT
GGACTAGTAAACCAAGCAATGACTTGTCTATTGATAGCCTTTTGCAACACTTTTATGACTCCTGAAATTTAGG
AATGCATTATATAAGTGGGAATTTGAAGATCTGAAGAAGATCCAGTGACAAGTATTCATACCAACTTCAAAGGC
TTTTGTTTTGTTACAACCAAGCAAAAGAGAGCAATTGAACCAACAGATGTTACAAGGAGCTTTGGATGGGATAG
TAGTGAGGCTTGGCAGCAGCATGATGTACAAGAACTATGCAGAGTCATGTTTGTGCTTTGGAACAGAAATGGAA
GCAACAGAAACAGGCTGATCTTATAAATGAGCTATATCAAGGCAAGCTGAAGGACTACGTGAGATGCTGGAAATG
TGGTTATGAGGGCTGGCGAATCGACACATATCTTGATATCCCATTTGGTCAATCCGACCTTATGGTCCAGCCAAGC
ATTTGCTAGTGTGGAAGCAATTCATGCAATTTATTAGCCAGAGATCTGGATGGCCCAATCAGTATTTTGT
GAACGTTGTAAGAAGAGTGTGATGCACGGAAGGCTTCGGTTTTTGCAATTTCTTATCTGCTGACCTTACAG
CTGAAAAGATTTCGATTTGATTATACAAACCATGCATAGGATTAAGTGAATGATCGAATGACATTTCCCGAGGAAC
TAGATATGAGTACTTTTATTGATGTTGAAGATGAGAAATCTCCTCAGACTGAAAGTTGCACGTGACAGTGGAGCAGA
AAATGAAGGTAGTTGTCACAGTGATCAGATGAGCAACGATTTCTCCAATGATGATGGTGTGATGAAGGAATCTG
TCTTGAACCAATAGTGAACCTGAAAGATCTCAAAATCTGGACTTGAAGAATTCCTTGATCTATGAACTTTTCT
CTGTTATGGCTCATTTCTGGGAGCGCTGCTGGTGCTATTATATGCATGTATAAGTCAATTCAGTGATGAGCAGT
GGTACAGCTTCGATGATCAACATGTCAGCAGGATAACACAAGAGGACATTAAGAAACACATGGTGGATCTTCAG
GAAGCAGAGGATATATTCTAGTCTTTCGCAAGTTCCACAAATGCATATATGCTGATCTATAGACTGAAGGATCC
AGCCAGAAATGCAAAATTTCTAGAAAGTGGTGAAATACCCAGAACATATTAAGAACTTGGTGCAGAAAGAGAGAGA

FIG. 10

GTTGGAACAAGAAAAGAGACAACGAGAAATTGAGCGCAATACATGCAAGATAAAATTATCTGTTTGCATCCT
ACAAAACAAGTAATGATGAAAAATAAATTGGAGGTTTCAAGGATAAGACATTAAAGGAAGCAGTAGAAATGGCTT
ATAAGATGATGATTTAGAAGAGGTAATACCCCTGGATTGCTGCGCTTGTAAATATGATGAGTTTCATGATTA
TCTAGAACGGTCATATGAAGGAGAGAAGATACACCAATGGGGCTTCTACTAGGTGGCTCAAGTCAACATATAT
GTTTGATCTGCTGTTGGAGACGAGAAAGCCCTGATCAGGTTTTCCAACTTATAAACCTGGAGAAAGTATGTTGAA
AGTTCATGTTGTTGATCTAAAGGCAGAACTGTAGCTGCTCTATAAAGTGTTCGTGCTTACTTAAATCAGACAGTT
ACAGAATTCAAAACAACCTGATTTCAAAGGCCATCCATTACCTGCTGAACAATAGAAATAGTCTGGAACGCTGC
TACAATGATTTGCGTCTTCTCAGTGTCTCCAGTAAACCCCTGAAAGCTGAAGGATTTTTAGAAAGTAAACAAGGTGT
TTGTTGAAAGCTCCGAGACTTTGGATTACAGATGGCTTTGCGAGACTCTCATTTATGAAACTCCTGGATCGGC
ATGCAAATACAATCAGATTATTTGTTTGTACCTGAACAATCCCGAGTATCTTATCCAAAAGGACAGCATACCA
GAAAGCTGGAGGCGATTCTGGTAATGTGGATGATGACTGTAAAGAGTCAAGGACCTGTAGGAAGCCCTAAAGT
CTGTGGAAGCTATTCTAGAAGAAAGCACTGAAAACCTCAAAAGCTTGTCACTGCAGCAACAGCAGGATGGAGATA
ATGGGACAGCAGCAAAAGTACTGAGACAAGTGACTTTGAAAACATCGAATCACCTCTCAATGAGAGGACTCTT
CAGCATCAGTGGATAATAGAGAACTTGAACAGCATATTCAGACTTCTGATCCAGAAAATTTTCAGTCTGAAGAACG
ATCAGACTCAGATGTGAATAATGACAGGAGTACAAGTTCAAGTGCAGAGTGATATCTTAGCTCCAGTCATAGCAG
TGATACTTTGTGCAATGCAGACAATGCTCAGATCCCTTTGGCTAATGGACTTGACTCTCACAGTATCACAAAGTAGT
AGAAGAACGAAAGCAAAATGAAGGGAAGAAAAGAAACATGGGATACAGCAGAGAAGAACTCTGGAAGTATGATGA
ATATGATGAGAGTGGCAAGAGTAGGGGAGAAATGCAGTACATGATTTCAAAGCTGAACCTTATGCTGCAGATGA
AGGTTCTGGGAAGGACATAAATGGTTGATGGTGATGTTGATAAAAGAAATTAAGTCTGGCAGCTTTCAAAACAACA
TTTAGAGCCCTTTGTTGGAGTTTGTCTCTCCTCAAGGCTTTTCGAGTGATGCCAGCAATCAAGAGTTTGAG
AGCGTCCGGCTGAATGAGACACTTTTCATCATTTCTGATGACAATAAGATTACAATTAGACTGGGAGAGCACTT
AAAAAGGAGAAATACAGAGTTAAAGTATACCAAGCTTTTGGTCAATGAACAAGAGCCATGCAAGTTTCTGCTAGAT
GCTGTGTTTGTCTAAAGGAATGACTGTACGGCAATCAAAAGAGGAATTAATTCCTCAGCTCAGGGAGCAATGTGGT
TTAGAGCTCAGTATTGACAGGTTTCGTCTAAGGAAAAAACAATGGAAGATCCTGGCACTGTCTTTTGGATTATC
ATATTTATGAAGAAGATATTAATATTTCCAGCAACTGGAGGTTTTCTTGAAGTTCTTGATGGGTAGAGAAGAT
GAAGTCCATGTCACAGCTTGAGTTTTGTCAAGACGGTGAAGCCCTCAGAGATGAAGTTGGATCCCTTCCAGG
AGGTTGATTGGAAAGCAGTAGTGTGGACGAATTCGAGAGAAAGCTTAGTGAATCAGTGGGATTCCTTTGGATG
ATATTGAATTTGCTAAGGTAGAGGAACATTTCCCTGTGATATTTCTGCTTGTATTCATCAGGATTTAGACTG
GAATCCTAAAGTTTCTACCCCTGAATGTCTGGCCTCTTTATATCTGTGATGATGTTGCGGTCAATTTTATAGGGAT
AAACAGAAAGAAATTAATGGAATTGACAGATGAGCAAAAGAAATGAAGTGAATGAAGAAAGCAAGTCCGACTCCAG
AAGACTGGACATCGTGAACATACTCACCTCGTAAAGAGAAAGCACTAAATAATATCTGGATGGAGCAACCAAT
AAAGATCTGACTCAAGACTGA

FIG. 1P

>SGPR480_SEQID_14
ATGGGTGCCAAGGAGTCACGGATCGGATTCCTCAGCTACGAGGAGCGCTGAGGAGAGTTACAGATGTAGAGC
TAAACGACTGAAGGATGCTTTCAAGAGGACCTGTGGACTCTCATATTACATGGGCCAGCACTGCTTCATCCGGG
AAGTCTTGGGATGGAGTGCCTCCAAAGTTGCTGAGGTGATTTACTGTTCTTTGGTGAACATCCAAAGGG
CTGCACTTCAATAATTAATAGTTGGACTTGTCTCCTTACAGAGGCAAGATGAAGAGAAAGCAAAATACATTT
TTAGTCTTTTTCAAGTGAATCTGGGAACCTATGTTATACGGGAAGAAATGGAAGAAATGCTCCACGTGGTGATG
GTAAAGTCCAGATACACTCAGGAAGTGTCTCAGAGGTGAAAGGTAACTATGAAGTTTAGAAATTGGC
TTTTCTAAACAAGATGCTTTTACTTTCTCTCGATGGCTTCTATCTGGAGGTGTATGTTACCCCTCACTGATGAT
AGTGATACTCTACTTTCTACCAAACTCTGGCTGGAGTCACACATTTGGAGGAATCAGACATCATTTGATCTTGAGA
AACGCTATTGTTATTGAAGGCTCAATCCCGGACTGGACGATTTGATTTAGAGACATTTGGCCCATTTGGTTTCAC
CACCTATTCGTCCATCTTAAGTGAAGGTTGTTAATGCTTTTGTGAATCGTGACAAATCAGATAGATTTTAAG
GAGATATCCTGTGGTTATCAGCCTGTTGCAGGGGACCCCTGGCTGAAGAGACAAAATTTTGCTTCAAGGTATTT
GATGTTGACCGTGATGGAGTTCTCTCCAGGTTGAACCTGAGAGACATGGTGGTTGCACITTTAGAACTCTGGAA
GGACAACCGCACTGATGATATTCCTGAATTACATATGATCTCTCTGATATTGTAGAAGGCATACCTGAATGCACAT
GACACCACAAGATGGGTCACTTACTCTGGAAAGACTATCAGATCTGGAGTGTGAAAATGTTCTTGCCCAATGAG
TTTTTGAACCTCCTTTTCCAGGTGTGCACATAGTTCTGGGGTTAAGACCAGCTACTCCGGAGAAAGAGACAA
ATTATTAGAGGATGTTAGAACGAGAGCAGGTATGGTCTGCAAGCAGGACACAACTGGTTTATCATCTCCATG
CAGTGTGGCAACAGTGGAAAGAAATATGTCAATACGATGCCAACCCCTGTGGTAAATTGAGCCATCATCTGTTTG
AATGGAGGAAAATACTCAATTTGGAACTGCAGCCCATCCTATGGAGCAGGTGGAAGATAGAAATTGGAGCAGCCT
CAGTTACGTGAATACTACAGAAAGAGAAAATTTTACAGACAACATTTCTACTGCATCTGAAGCCTCAGAACTGCTGGC
AGCGGCTTTCGTATTTCTGCCACACCCAGGGGCAGATGTTTGGCTCGACAACATAACACTTCTGACAATAAC
AACCAGTGTGCTGGAGCCAATGGGAATATTTTGTGCACCTTAACCCCTCAGAAACCAGGGCTATTGATAAT
CAGCCATTAGTAACCAAGAACCAGTAAAGGCTACATCATTAACACTAGAAGGAGGACGATTAACAGCACTCCA
CAGCTGATTATGGAAGAGACTATGAAATGGTCCAGAACCTGTGTGGAGAGCACTTTATCAGTGGTATGGAGCA
AACCTGGCCTTACCTAGACCAGTTATCAAGAACAGCAAGACAGACATCCAGAGCTGGAAATATTTCCCGCTAT
CTTCTCTTCTGAGACAGCAGCCTGCCACTCGGACACAGCAGTCTAACATCTGGTGAATATGGGAATGTACCT
TCTCCGAATGCACCTTTAAAGCGGGTATTAGCCTATACAGGCTGTTTGTAGTCGAATGCAGACCATCAAGGAAAT
CAGGAATATCTATCTCAAGGCTGCGCATTAAGAGGAAGATATGCGCCTGTGGCTATACACAGTGAAGAACTAC
CTTACTCTTCTGGATGATGAGGATCATAAATTGGAATATTTGAAAATCCAGGATGAACAACACCTGGTAAATTGAAG
TTCGCAACAAGATATGAGTTGGCCTGAGGAGATGCTTTTATAGCAATAGTAGTAAATAGATAGACACAAGGT
TCCACAGAAAAGGAGGCCACAGGTCTAAGCAATCTGGGAAACACATGCTTCATGAACCTCAAGCATCCAGTGTG

FIG. 1Q

TTAGTAACACAGCCACTGACACAGTATTTTATCTCAGGGAGACATCTTTATGAACTCAACAGGACAAATCCCAT
TGGTATGAAGGGGCATATGGCTAAATGCTATGGTGATTTAGTGCAGGAACCTTTGGAGTGAACCTCAGAAGAATGT
TGCCCCATTAAAGCTTCGGTGGACCATAGCAAAAATATGCTCCAGGTTTAAATGGGTTTCAGCAACAGGACTCCCA
AGAACTTCIGGCTTTTCCTTGGATGGTCTTCATGAAGATCTTAATCGAGTCCATGAAAAGCCATATGTGGAACCTG
AAGGACAGTATGGCGCAGACTGGGAAGTAGCTGCAGAGGCTGGACAAACCATCTAAGAAGAAATAGAT
CAATTGTTGGATTGTTCCATGGCAGCTAAGATCTCAAGTAAATGCAAGACATGTGGGCATATAAGTGTC
GATTTGACCCTTTCAATTTTGTCTTTGCCACTACCAATGGACAGTTATATGCACCTTAGAATAACAGTGATTAAG
TTAGATGGTACTACCCCTGTACGGTATGGACTAAGACTGAATATGGATGAAAGTACACAGGTTTAAAAAACAG
CTGAGTGATCTCTGTGGACTTAATTCAGAACAAATCCTTCTAGCAGAAGTACATGGTTCCACACATAAAGAACTTTC
CTCAGGACAACCAAAAGTACGACTCTCAGTGAGTGGATTTTGTGTGCAATTTGAAATTCCTGTCCCTGTGTCTCC
AATTCAGCTTCTAGTCCAACACAGACAGATTTCTCCTCTCGCCATCTACAAATGAAATGTTACCCCTAACTACC
AATGGGACCTACCCGACCAATATTATCCCAATGGAATGCCAAACACTGTTGTGCCATGTGGAACTGAGAA
GAACTTCACAAATGGAATGGTAAATGGTCACATGCCATCTCTTCTGACAGCCCTTTACAGGTTACATCATTGCA
GTCCACCGAAAAATGATGAGGACAGAACTGTATTTCTGTCTATCTCAGAAGAAATCGCCACAGCCTCTTTGGAATG
CCATTGATTGTTCCATGTACTGTGCATACCCGGAAGAAAGACCTATATGATGCGGTTTGATTCAAGTATCCCGG
TTAGCGAGCCCACTCCCACTCAGGAAGCTAGTAATCATGCCAGGATTGTGACGACAGTATGGGCTATCAATAT
CCATTCACTACGAGTTGTGCAGAAAGATGGAACTCCTGTGCTTGGTCCCATGTATAGATTTGCAGAGGC
TGTAATATTGATTGTGGGAAGACAGAGCTTTTATTGGAATGCCATATCGCTGTGATTGGGATCCACAGCC
CTTCACCTTCGCTATCAACATCCAGGAAGGGTTGTAGATGAGCATGAGAGTGTGGAGCAGAGTCGGCGAGC
GCAAGCCGAGCCCATCAACCTGGACAGCTGTCTCCGTGCTTTACCCAGTGAGGAAGAGCTAGGGGAAATGAG
ATGTACTACTGTTCCAAGTGTAAAGACCCACTGCTTAGCAACAAGAAAGCTGGATCTCTGGAGGCTTCCACCCATC
CTGATTTACACCTTAAGCGATTTCAATTTGTAATGGTCGGTGGATAAAATCACAGAAAATTTGTCAAATTTCCCTCG
GGAAAGTTTGTATCCAAGTGTCTTTTGGTACCAAGAGACCCGGCTCTCTGCCAGCATAAACCACTCACACCCCA
GGGGATGAGCTCTCTGAGCCGAGGATTCTGCAAGGGAGGTGAAGAAAGTGGATGCGCAGAGTTCGGCTGG
GGAAGAGGACGTCTCTGAGCAAAAGCCCATCCTCACTAGCGCTAACATCATCAGCAGCCCGGAAAGTTCTC
CTTCTTCATCAGAAAAAGTGAACCAAGCTGTCCCTCCAGCAAAACAGCAGCCCTAATAGCAGCCACGGACTT
TGGGAGGAGCAAGGGAGGCTCCGGCTGCCGCTGCCCAGATTGGCAGCAAAATAAAGTGTCAAGTAGTAAAGAGAA
CTTGATGCCAGCAAGAAATGGGCTGGGAGATATGTGAGCTGGCTGACGCCCTTGAGTCGAGGGCATGTG
CTGGGGGCGAGCCAAACAGAGTTGGTCACTCTCAGGACCATGAGGTAGCTTTGGCCAAATGGATTCCTTTATGA
GCATGAAGCATGTGGCAATGGCTACAGCAATGGTCAGCTTGGAAACCAAGTGAAGAACAGCAGCAGTATGACC
AAAGAGAAGATCTCGTATTAAAGCCTATTTATATCTATATGCAATTTCTGTCCTTCAAGGAAATCTGGTGGGG
CCATTACGTCACTTATGCCAAAAACCCAAACTGCAAGTGGTACTGTACAAATGACAGCAGCTGTAAAGGAACTTCA

FIG. 1R

CCCGGATGAAATTGACACCGACTCTGCCCTACATTCTTTCTATGAGCAGCAGGGGATAGACTATGCACAATTTCT
GCCAAAGACTGATGGCAAAAGATGGCAGACACAGCAGTATGATGAAGACTTTGAGTCTGATTACAAAAAGTA
CTGTGTGTTACAGTAA

>SGPR431_SEQID_15

ATGGACAAGATCCTGGAGGGCCTTGAGTTCTCGCATCCCTGCCCTCAAGCGGTGATTGTGCGGAAGG
TGGTGAATCGGCGGAGCACTGGCTAGACGAGGCGCAGTGCAGGCCATGTTGACCTGACGACCCGGCTCAT
CCTGGAGGGCCAGACCTTTCCAGCGCAGGTGGGCACCAAGTCTGGAGGCCCTAGCAGCATACCAACCG
GCCAGAGTTCGAGTCCTTCAACAAGACCTTCGTGTGGCCTCCTTCATCAGGGCTACCACTCTCTGGACAG
GAAGGATGTAGCCATCCTGGACTACATTCACAACGGCCTGAAGCTGATTATGAGCTGCCGTGGTGGTGGATC
TCTTAGCCTCCTGCAGGTAGAGGTGTTACGGATGTTGTGTGAGAGCGGAGCCGAGCTCTGTGCCCGACT
GAGCGACCTCTGACCGACTTTGTGCAATGCATCCCAAGGGAAATTGTCCATCACGTTCTGTCAACAGCTGGT
TCGAACGATAGGCCATTTCCAGTGCCTGTCACCCAGGAAGAGAGCTGCGGGAATATGTCTCCAGGTGACAA
AAGTGAGTAAGTCTGCAGAACATCTGGAAGGCCGAGCCTGCCACACTACTGCCCTCCCTGCAAGAAGTTTTG
CAAGCATCTCTCCACAGATGCATCATTTGAACCTTCTGTAGCATTTGGCAAGCCTTGTGCAGCATATTCTCTTCA
GATGATTACAGTTCTCATCAGGAGCCTTACTACGGATCCAAATGTAAAGATGCAAGTATGACCCAAAGCCCTTG
CAGAAATGATTGACTGGCTATCCTGGCCATTGGCTCAGCATGTGGATACATGGTAATTGCACCTCCTGAAAGGACT
GGCAGCTGCCAGAGTTACTATTTGATAGATGTTACTTTGCTGAAATAGAACTGGTTTTAATCGACTTTGGT
TTCCCTTTGTGAGACCTGGTCTCTTGCAGTTCTTTCTCACATGCTGCTTAGCTTTCAGCATTTCTCCAGAGGCGTT
CCATTTGATTGTTCTCATGTGGTTAATTTGGTTCAATCTTTCAAAATGATGGTCTGCCCTTCAAGTACAGCCTTCT
TAGTACAATTACAGAAATTGATACACTGTATGATGATCATTAATCTGGATTTCAGATCTCTATGAACCTATTCTG
GAGGCAATAAAGGATTTCTAAGCCAGTGAAGAGAAGATTAAAGTTAATTTCTCAATCAAAGTGCCTGGACTTCTC
AATCCAATTTCTGGCGTCTTGCTGTAGACTTTCTGGAAATCTGAAACTGGGAAACTGGTCTTATTAACTT
AGGAAATACATGTTATGAACAGTGTATACAAGCCTGTTTATGGCCACAGATTCAGGAGACAAAGTATTATCT
TTAAATCTAAATGGTGCAATTCATTAATGAAAAAATTACAGCATCTTTTGGCCTTTCTGCCCCATACACAGAGGG
AAGCATACGCACCTCGGATATTTCTTGAGGCTTCCAGACCTCCATGGTTTACTCCAGATCACAGCAAGACTGTT
CTGAATACCTCAGATTTCTCCTTGACAGGCTCCATGAAGAAGAAAGATCTTGAAAGTTTCAGGCCCTCACACAAGC
CTTCTGAAATTCGGAATGCAGTGAACCTTTTACAGGAAGTAGCTAGTAAAGCAGCAGTACTAAACAGAGACCC
CTCGTACAAAGTGACGGTGAGAAGACTTTAATAGAAAAAATGTTTGAGGAAAACTACGAACTCACATACGTTGTTT
GAACTGCAGGAGTACCTCACAAAAAGTGAAGCCTTTACAGATCTTTTCGCTTGGCTTTTGTCTTCTCTCTTTG
GAAAAATGTCTGTCCAAAGATCCAGCATCATCACCCAGTATACAAGATGGTGGTCTAATGCAAGCCTCTGTACCC
GGTCCCTTCAGAAGAACCAGTAGTTTATAATCCAACAACAGCTGCCCTTCATCTGTGACTCACTGTGAATGAAAAA

FIG. 1S

CCATAGGCAGTCTCCTAATGAGTTTTACTGTTCTGAAACACACTTCTGTCCCTAACGAATCTAACAGATTCTTGT
TAATAAGATGTACCTCAGAAACCCAGGAGGTGAACCCACACACCTTCAGTAACCTGACTTACTAAATATTTTTGGCT
CCAGAGATTCTTACTGGTGATAACCAATATTAATTGTGAAACCTGTGCCCTCTCTGCAAAATGCTGAGAAACTATGC
AAATCACGGAGAACCTGAATACCTTATCTTACTCTCTGAGATTTTCATATGATCAGAAGTATCATGTGAGAAG
GAAATTTTAGACAATGTATCACTGCCACTGGTTTTGGAGTTGCCAGTTAAAGAAATTAATCTTCTTCTTCAATTGT
CAGAAAGTTGGTCTGTAGATGTTGACTTCACTGATCTTAGTGAGAACCTTGTAAATAATTAAGCCTTCAGGGAC
TGATGAAGCTTCTGCACAAATTTGGTGCCCTATCTATTAAGTTCCGTTGTGTTCACTCTGTGTATATCCTCTGAA
AGTGGCATTACTATTCTTATGCCAGAAATATCACAAGTACAGACTCTTCATATCAGATGTACCACAGCTGAGG
CTCTGGCATTAGCATCTCCAGAGTCACTTACTAGGGAGAGATAGTCCAGTGCAGTTTTTGAACAGGATTTGG
AAATAAGGAAATGTCAAAGAATGGTTTTTATTAATGACAGTAGAGTGACATTTACTTCAATTCAGTCAGTCCAG
AAATTACGAGCAGGTTTCCAAAGGACACAGCTTATGTCTTTGTATAAAACAGCATAGTACTAATGGTTTAA
GTGGTAATAACCCAAACAGTGGACTCTGGATAAATGGAGACCCACCTCTACAGAAAGAACTTATGGATGCTATAA
CAAAAGACAATAAATAATTTACAGGAACAAGAGTTGAATGCTCGAGCCCGGCCCTCCAAAGCTGCATCTGCTT
CATGTTCAATTCGGCCCAATGGATTTGATGACAACGACCCACCAAGGAAGCTGTGGACCAACTGGTGAGGGGGT
GGAGGAGGATTTAATACAGTTGGCAGACTCGTATTTTGA

>SGPR429_SEQID_16

ATGGCCCCGGGCTGCAGCTGGAGAAAGCGGCGCTGGCGCTGGCGGAGACGGTGGCGCCGAGGAGGTGTC
GCAGGAGCATATCGAGACCGCTTACCGCATCTGGCTGGAGCCCTGCATTGCGGCGGTGTCAGACGAAACTGC
AAAGGAAATCCGAATTGCTTGGTGTATGGTGAGCATATTTGGTTAGGAGAAATAGATGAAATAGTTTTTCATA
ACATCGATGATCCCAACTGTGAGAGGAGAAAGAACTCAATTTGTGGGCTGACTAACCTTGGAGCCACTTGT
ATGTCAACACATTTCTCAAGTGTGGTTTCTCAACTTGGAGCTTCGGCAGGCACTCTACTTATGTCCAAGCACTTG
TAGTGACTACATGCTGGGAGACGGCATCCAAGAAAGAAAGATTATGAGCCTCAACAAATTTGTGAGCATCTCCA
GTACTTGTTCCTTGTGCAAAACAGTAATAGGCGATACATTGATCCATCAGGATTTGTTAAAGCCTTGGGCGCTG
GACACTGGACAACAGCAGGATGCTCAAGAAATTTCAAGCTCTTTATGCTCTATTGGAAGATACTTTGTCTAAAC
AAAAGAAATCCAGATGTGCGCAATATTGTTCAACAGCAGTCTGTGGAGAAATATGCCCTATGTAACTGTTTGAACCA
GTGTGGCAGAGTCTAAGCTTTGTCAAATTTTATGAGCTGGAGTTAAATATCCAAGGCCACAAACAGTTAACA
GATTGTATCTCGGAATTTTGAAGGAAGAAATTAGAAGGAGACAAATCGCTATTTTGGAGAACTGTCAAAGCA
AACAGAAATGCAACAAGAAAGATTGACTTCTTAGCCTTCCCTGCACTCTGAACCTGCAGCTAATGCGTTTTGTCTT
TGACAGGCAAACTGGACATAAGAAAGCTGAATACCTACATTGGCTTCTCAGAAATTTTGGATATGGAGCCTTAT
GTGGAACATAAAGGTGGTCTACGTGTATGAACCTCAGCGCAGTCTCATACAGAGGAGTGAGTGCTTATCT
GGCCACTACATCGCCCACTGAAAGATCCACAGTCTGGTGAATGGTATAAGTTAATGATGAAGACATAGAAAAG

FIG. 1T

ATGGAGGGAAGAAATTACAACCTAGGGATTGAGGAAGATCTAGAACCTTCTAAGTCTCAGACACGTAAACCCAAAG
 TGTGCAAGGAACTCATTGCTCTCGAAATGCATATATGTTGTTTATAGACTGCAAACTCAAGAAAGCCCAACA
 CTAAGTTCAGATTCCAGCCTTCTTCAAGAGCTGGTAGTCGGGATAATCCAAATTTGAGGAGTGGTGATTGA
 AATGGCTGAGATGCGTAAGCAAAAGTGTGGATAAAGGAAAGCAAAACACGAAGAGTTAAGGAGCTGTACCAAA
 GGTTACCTGCTGGAGCTGAGCCCTATGAGTTTGTCTCTCTGGAATGGCTGCAAAAGTGGTTGGATGAATCAACAC
 CTACCAAACCTATTGATAATCACGCTTGCCCTGTGTTCCCATGACAAGCTTCACCCGGATAAATATCAATTATGAA
 GAGGATATCTGAATATGCAGCTGACATTTTCTATAGTAGATATGGAGGAGTCCAAAGACTAACTGTGAAAGCCCT
 GTGTAGGAATGTGTAGTAGAACGTTGTGCGATATTGCGTCTGAAGAACCAACTAAATGAAGATTATAAACTGTT
 AATAATCTGCTGAAAGCAGCAGTAAGGGCGATGGATTTTGGTGGGGAAGTCTCTCTTGGGAGTTGGCGCA
 GCTAGCTCTTGAACAGCTGGATGAGCAAGATGGTGATGCAGAACAAAGCAACGGAAGATGAACGGTAGCACCT
 TAAATAAGATGAATCAAGGAAGAAAGAAAGAGAGGAGGAATTAATTTTAATGAAGATATTCTGTGTCCACA
 TGGTGAGTTATGCATATCTGAAAATGAAGAAGGCTTGTTTCTAAAGAGGCTTGGAGCAAACTGCAGCAGTACTT
 TCCAAAGGCTCCTGAGTTTCCAAAGTTACAAGAGTGTCTACAGTGCAGATTTTAGAAAGAGAGGGGAAGA
 AATGAAGCCTTACATAAGATGATTGCAACGAGCAAAAGACTTCTCTCCCAAATTTGTTCCAGGATAAAACAGA
 CCGTGTCTCAGTAAGTGGCCAGAGGATACGGATGTCCTCTACATCGTGTCTCAGTCTCTTTGTAGAAGAGTGGCG
 GAAATTTGTTAGAAAGCCTACAAGATGCAGCCCTGTGTCTCAGTTGGGAACAGTGTCTTTTGTGTCCACGG
 GGGCCTCATGTTACATTTGCTTCCATGACCAAGAAAGATTCTAACTTATAGCTCTCATATGCCCAGTGAGTGG
 CAAATGATACAAAGCTCTTTGTTGTGGATCATGTAATTAATCAACGAGAAATGAAGTGGGAGATGTAACCCCTT
 CAGAAACACAGTATATTTCTGAGCCCAACTCTGTCCAGAAATGCAGAGAGGCTTATTTGTGTAGCAGCAGAGGG
 ACCTGCGTGAATACACTCAAGCCACCCTATATGTCCATAAAGTTGTGGATAATAAAAGGTGATGAAGGATTCCGG
 CTCCGGAAGTGAATGTGAGTAGTTCTGAAACAGAGGAGGACAAAGGAAGCTAAACCCAGATGGAGAAAAGAT
 CCAGATTTTAAATCAAGCAATGGTGGACAAAGCGGCAAAAGATATCCCATCAAAATATATAGCCTATCAAAAGC
 AAGTTATTCGCCGAAGTATCGGACATAGAAAAGTTCTGTGGTGAAGAACACTTCTCGTTTCTGCTAATCAGACGT
 TAAAGAAATTGAAAATTCAGATCATGCATGCAATTTTCAGTTGCTCCTTTTGACCAGAAATTTGTCAATTGATGGAAG
 ATTTTAAGTGATGACTGTGCCACCCTAGGCACCCCTTGGCGTCATTCCTGAATCTGTCTATTTTGAAGGCTGATG
 AACCAATTGCAGATTATGCTGCAATGGATGATGTCATGCAAGTTTGTATGCCAGAAAGGGTTTAAAGGTACTG
 GTCTTCTTGGACATTAA

>SGPR503_SEQID_17

ATGCTGAGCTCCCGGGCCGAGGCGCGGCGATGACCGCGCGCCGACAGGGCCATCCAGCGCTTCTCGGACCGGG
 GCGGCCGTGAGATATAAGTCAAGAACTGGGAGTTATAGGTGGAATTTGCTGCTGCTCTTGCAGCAGGAAT
 ATATGTTATTTGGGTCCCATACAGAAAAGAAAGCGTAGAAAAGGGCTTGTGCTGGCCTTGTTAATTTAGG

FIG. 1U

GAACACCTGCTTCAATGAACCTCCCTGCTACAAGGCCCTGCTGCCCTGTCTCTCATCAGGTGGCTGGAAGAGTT
CACCTCCCACTACTCCAGGGATCAGAAGGAGGCCCTCACACCAGTATTTATCCTTAACACTCTTGCACTTCT
GAAAGCCTTGCTCTGCCAAGAGTTACTGATGATGAGGTCTTAGATGCAAGCTGCTTGTTGGATGCTTTAAGAAAT
GTACAGATGGCAGATCTCATCATTTGAAGAACAGGATGCTCACGAAATTATCCATGTCATTACCTCGTCATTGGAA
GATGAGCGAGACCGCCAGCCTCGGGTCACACATTTGTTTGATGTGCATTCCTGGAGCAGCAGTCAGAAATAAC
TCCCAACAAATTACCTGCCGCACAAGGGTCACCTCACCCACATCCAATCACTGGAAGTCTCAACATCCTTT
TCATGGAAGACTCACTAGTAATATGCTCGCAACACTGTGAACACACAGAGTCCCTGTTTCGATTTGATACCTTTGAT
AGCCTTTCACTAAGTATTCAGCCGCCACATGGGGTCACCCATTGACCCCTGGACCACTGCCCTTACCACCTTCATC
TCATCAGAAATCAGTGGGGATGTTGTGTGACAACTGTACAAAGATTGAAGCCAAAGGAAACGTTGAACGGGGA
AAAGGTGGAACACCAGAGGACCACCTTTTGTAAACAGTTAAACTAGGGAAGCTCCCTCAGTGTCTCTGCATCCA
CCTACAGCGCTGAGCTGGTCCAGCCACGCGCCCTCTGAAGCGGCATGAGCACGTGCAGTTCAATGAGTTC
CTGATGATGGACATTTACAAGTACCACCTCCTTGGACATAAACCTAGTCAACACAACCCCTAACTGAACAAGAAC
CCAGGGCCTACACTGGAGCTGCAGGATGGCGCGGAGCCCCACACCAAGTTCTGAATCAGCCAGGGGCCCCC
AAACACACAGATTTTATGAATGGCGCTGCTCCCCATCTTTAATTGCCAACGCTGTACGCGCCGATGCCCTTCCCT
CTCCCAGTTGTTCCCGACTACAGCTCCTCCACATACCTCTTCGGCTGATGGCAGTTGTCTGCCACCATGGAGA
CATGCACCTGGACACTTTGTCACTTACCGACGGTCCCCACCTTCTGCCAGGAACCCCTCTCTCAACTAGCAATCA
GTGGCTGTGGTCTCCGATGACACTGTCCGCAAGGCCAGCCTGCAGGAGGTCTCTGTCTCCAGCGCCTACCTG
CTGTTCTACGAGCGCTCCTTTCCAGGATGCAGCACCAAGCCAGGAGTGCAAGTCTGAAGAATGA

>SGPR427_SEQID 18

ATGGACCTGGGCCCGGGACGCGGACGGGAGGGGACCGCTCGCGCCCGGCCCGCCGCGCTCCCT
GCGCGCTGTTAGCCGCTTCTGCTGGCGTGGGACCGCTCAGCCCCGGGACTACCGCCCCGGCC
CCAGCCGGACACTGTATGGCGACGTGAGGGGCTTCGCTGCGCCCGGGCCAGTTCCAGCGGCCCC
CGGAGCCCCGGGAGAACGCCCGCCGACCCAGCTCCAGCTCCCGCGCGCATGGGGCGC
GCCGCCGGCGCTCAGGGCTTGAAGAACACGGCAACACTGTTTCATGAACGCGGTGGTGCAGTCTCAG
CAACACGACCTGCTGGCCGAGTTCCTGGCGTGGGCGCTACCGGGCGCTCCGGCCGCGCAGGTCAC
CGAGCAGCTGGCGGCGCTGGTGC GCGCTCTGGA CTGCGAATACGCCCCAACTTTCGCGGAGTTCAAG
AATGCAGTTTCCAAGTACGGCTCTCAGTTCCAAGGCAATCCAGCACGACGCCCTGGAATTCCTGCTGGTTG
CTGGATCGTGATCAGGACCTGGAGGTTTCATCCCGAGGGCGGTGTCGGAGAAGCTTCGCGCTGAAGCCA
CTAAACCTCTGAGAACTGCCTGTCAACCATCAGCTCAGCTTCCTCTAGGTCAAAGCTTGTGCAAAGCCACTTTC
AAGCACAAATAGATCTTCCTTGACTTGTCCTCCCACTGCCCTGAACAGAGCAACACCTTTGATCCTTTCTGTGTG
GTCCCTACCTATCCCCCTTGGCGCCAGACGAGGTTCTTGAGTGTCACTTGGTCTTCCCTCTAAGAGCCACGGT

FIG. 1V

TCCTGCGGTTGGCCGTGCCGATCCTCAGCACAGTGGCAGCCCTGAGGAAGATGGTTGCAGAGGAAGG
AGCGTCCCTGCAGATGAGGTGATCTTGGTTGAACGTGATCCAGTGGATCCAGCGGTCTTTCTTTGATGAAGA
GGACCTGAATACCATCGCAGAGGGAGATAATGTGTATGCCCTTCAAGTTCTCCCTCACCAGCCAGGGGACTC
TCTCAGCTCATCCACTGGGTCTGTGGCCCTCCACAGCCTGGCAGCCCGTGAGGGCCAGCGATTCTCCCTCTCT
CTCCACAGTGAGAGCAAGTGCTAATCCTCTTCTGTAACTTGGTGGGTGAGGCGCAGGCTAGCAGGTTTGG
GCCACCCCTTCTGATAAGGGAAGACAGAGCTGTTCTTGGCCCAAGTCCAGCAGTCTATCCTCAGCAAGTCC
GCCATCTTATGAAGAGTGAGGCCCTGTACAGAACCTGGGTCTCTGTTCTCCATCCGTGTTGTGGACTCTCT
GTGGCCTGCAGCTATTTGTCTCCGAAGGACAGTCCGCCCTCTGTCACTGGCAGTTGACAGGGTTTGCATCT
CAGGAGGCCAGGAGGCCCTCCACATGTCAAGCTGGCGTGGAGTGGATAGCTCTGTCAAGGAGCGCTGTTT
GGGAGCCTCCAGGAGGAGCGCAGGATGCCACAGTGTGTGGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGC
AGCTGTACCTTGGATGAATGTTTTCAGTTCTACCAAGGAGGAGCAGCTGGCCCAAGGATGACGCCCTGGAAGTG
TCCTCACTGCCAAGTCTGCAGCAGGGGATGGTGAAGCTGAGTTGTGGACGCTGCCTGACATCCTCATCATCC
ACCTCAAAAGGTTCTGCCAGGTGGCGGAGAGAAAGCAAGCTCTCCACGCTGGTGAAGTTCCGCTCTCATCC
CTCAACATGGCTCCCATGTGGCCAGAGAAAGCAGCCCTGAGGCGAGCTGGCCCTGGCCCTTCCCTGGA
AGCAGCCGACTGCCTGCCACAGTTACCCGCTGGACTTCTGTACGACCTGTATGCCGTCTGCAACCCACCAT
GGCAACCTGCAAGGTGGCATTACACAGCCTACTGCCGGAATCTCTGTGATGCCAGTGTACAGTTATGATGA
CAGCAGGTGGAACCGCTTCGAGAAGATGAGGTCAACACCAAGAGGGCTTATATCCTGTTCTATCAGAACGGA
ACAGCATCCCTCCCTGTCAGCCAGCAGCTCCATGAGAGGCTTACCAGCTCCTCCCTGTCTGATCACTGGCTC
TTACGGCTCGGAGCCACGCTGGCAGCACAAAGGGAAGCCTGTCTCTGGAGCTCTGCCCTGCCCCCTCCC
TGCCCCAGGTTCTGACTCTCCCATCTTCAACCAACAGCCTCTGCAATCAGGAAAGGGAGGTTGGAGCCCAAG
CGTTTGTACGGCGGTGAAGGCAGAGCATTAGCATGAAGGCACCCACCACTTCCCGAGCCAGCAGGAC
CATTCAAGACCATTGCCTCTGCGGTGTTCTTGGATCCAAGGAGAAACCAACCAAGTGCCTCCGTCGAGTTGGTG
GAGTACTTGGATCCAGACGAGACCTCGGTCCAGGAGCCAGTCCATTGTGTCGTGTTGACGGCACTGCGG
GTGAGGATGAGAAATCAGCATCGCCGAGGTCCAAAGTCCGCTTCTGCTAACAGCGAAGATGGTGGCGGGC
CATTGAAAGAGGTCCAGCGGGTGCCTGTCCCTCGGCTCAACCCCAACCACTGTCTGGCCCCCTGAAACTCA
GATGGTCCAAACACAGCAAGGAACTCAAGGAAATGCAGGGCAGGACATCAAGCTTCCAGAAAGTTTGACCT
GCCTCTACTGTGATGCCCTTCAAGTGGAGCATGAGAAACCAAGTCCAGCGGAGGCGCAGAGCCATGAACCTG
AAGGAGAGCTTCCAGATGGGAAGCAAAAGCAGCCCAACCTTCCCTTATATGGATTCTCTGAAACAGCAAGA
CAGTCGCCGAGGCACCTCTGAGCTAGACAGACCCCTGCAGGGGACACTCACCTTCTGAGGTCCGTGTTTCGG
AAGAAGGAGAACAGGAGGAATGAGAGGCGAGAGTCTCTCCACAGGTGCCCCCGTCTCCCTGGTGGTGGCG
GGCTGAGCCCTGCCATGGACGGCAGGCTCCAGGCTCACCTCTCCCTCAGGATCCAGAGGGCCCTGGCCA
GGGCTGGCAGCCGCTCGAGAGGGATGTCTGGTCAGCCCCCAGCTCTCTCCGCTCCCTCGTAAAGCCA

FIG. 1W

GCAGGGCCCCGAGAGGCAGTGCACCTGGCATGTACAAAGGACTGTTCCAGGGGAGCAGGCTTCTTATGGCAC
CTTTCAGAGAGTCAATATCACACTCTTTCTTTAGGTCGAAAGAAACCTTACCGGAGTCCAGCTTTTGA

>SGPR092_SEQID_19

ATGCAGCTCGTCATCTTAAGAGTTACTATCTTCCCTGGTGTTCGCCGTTCCAGTGCCCCCTGCTGCAGAC
CATAAAGGATGGACTTTGTTGAGGGCTATTTCCATCAATTTTCCTGACCAAGAAGGAGTCGCCACTCCTTACC
CAGGAGACACAAACACAGCTCCTGTCAACAATTCCATCGGAATGGACAGACCTACTTGACATGCAGATGCATGC
TCTGCTACACCAAGCCCCACTGTGGGTGCTGATGGTCCGACACCTCCATCTCGCCAGGAAGATGCAAGTGG
AATAAGCACACTCTAACTTACAGGATTATCAATTACCCACATGATATGAAGCCATCCGCAGTGAAGACAGTATAT
ATAATGCAGTTTCCATCTGGAGCAATGTGACCCCTTTGATATTCAGCAAGTGCAGAAATGGAGATGCAGACATCA
AGGTTTCTTCTGCGAGTGGGCCCATGAAGATGGTTGGCCCTTTGATGGCCAGGTGGTATCTTAGGCCATGCC
TTTTTACCAAAATCTGGAATCCTGGAGTTGTCCATTTTGACAAGAAATGAACACTGGTCAGCTTCAGACACTGGAT
ATAATCTGTTCTGTTGCAACTCATGAGATTGGCAATCTTTGGCCCTGCAGCACTCTGGGAATCAGAGCTCCA
TAATGTACCCCACTTACTGTATCACGACCCCTAGAACCTTCCAGCTCAGTGCCGATGATATCCAAAGGATCCAGC
ATTTGTATGGAGAAATGTTTCATCTGACATACCTTAA

>SGPR359_SEQID_20

ATGAAGGTGCTCCTCGCATCTGGCCTTGCTGTCTTCTCATCATGGCTTTGAAGTTTCCACTGCAGCCCCCTCC
CTAGTTGCAGCCTCCCCCAGGACCTGGAGGAACAACACTACGCCCTGCACAGGCGTATCTTGACAAATATTACAC
AAATAAGAAAGGACACACAGATTGGTGAGATGGTTGCAAGAGGAAGCAATTCATGATAAGGAAGATTAAAGGAGCT
ACAAGCGTTCTTTGGCCTCCAAGTACCGGGAAGTTAGACAGACCAACAATGAACGTGATCAAGAAGCCTCGCT
GTGGAGTTCCTGATGTGGCCAATTATCGCCTCTTCCCTGGTGAACCCAAATGGAAAAAAATACTTTGACATACA
GAATATCTAAATACACACCTTCCATGAGTTCTGTCGAGGTGGACAAAGCAGTGGAGATGGCCTTGACGCCCTGG
AGTAGCGCCGTCCTGAGCTTTGTAGAAATAAATCAGGAGAGCGGATATTATGATATCTTTTGAAAATGGA
GATCAGCGGGATTCTCATTCGATGGCCCTCGGGGACTCTAGCCCATGCATTTGCTCCTGGAGAAAGCCT
GGGAGGAGATACACATTCGACAAATCCTGAGAAGTGGACTATGGGAACGAATGGTTTAATTTGTTTACCGTTGC
TGCTCATGAATTTGGCCATGCCCTGGCCTGGCCATTCCACAGACCCATCAGCACTGATGTACCCAACTTATAA
GTACAAGAAATCCCTATGATTCCACCTCCCAAGATGATGTGAAGGGATCCAGGCATTATACGGACCTCGGAA
AGTATTCTGGGAAGCCCACTCTGCCCATGCCCCCATCACAAGCCATCCATCCCTGACCTCTGTGACTCCA
GCTCATCCTTTGACGCTGTGACAAATGCTGGGGAAGGAGCTCTGCTCTTCAAGGACCGGATTTTCTGGAGACGG
CAGGTTCACTTGGGACAGGAATTCGGCCCAAGCACTATTACAGCTCCTTCCCCCAGCTCATGTCCAATGTGGA
TGCAGCTTACGAAGTGGCTGAGAGGGGCACTGCTTACTTCTTCAAAGGTCCCCCACTACTGGATAACAAGAGAT

FIG. 1X

TCCAAATGCAAGGTCCTCGGACTATTTATGACTTTGGATTTCCAAAGGCACGTGCAGCAATAGATGCTGCTG
TCTACCTCAGGGAGCCACAGAAGACCCCTTTCTTTGTGGAGATGAATACTACAGCTACGACGAAAGGAAAGG
AAATGGAAAAGACTATCCAAAAGAACTACTGAAGAAATTTTCAGGAGTAATGGCCAAATCGATGCTGCTGTA
GAATTAATGGCTACATTTACTTCTTTTCAGGACCAAAACATACAAAGTATGACACAGAGAAAGGAGATGTGGTTA
GTGTGTGAAATCTAGTTCTCTGATTGGTTGCTAA

>SGPR104_1_SEQID_21

ATGAACGTCCGCTGCAGGAGCTGGGAGCTGGCAGCAACATGGTGGAGTACAAACGGGCCACGCTTCGGGATG
AAGACGCACCCGAGACCCCGTAGAGGGCGGGCTCCCGGACGCCATGGAGGTGGATTCCAGAGGGGA
CAAGACAGCTGTAGGCTCAGCACGCAGCTGGAGCTGGTCTTAGCAGTGCCTCTCTACTGCTGGCTGCACCTG
CTTCTGGCTGCCCTGTGGCCCTAGGGTCCAGTACCACAGAGACCCATCCACAGCACCTGCCTTACAGAGG
CCTGCATTGAGTGGCTGGAAAATCCTGGAGTCCCTGGACCGAGGGGTGAGCCCTGTGAGGACTTTTACCA
GTTCTCCTGTGGGGCTGGATTCGGAGGAACCCCTGCCGATGGCGTTCTCGCTGGAACACCTTCAACAGC
CTCTGGACCAAAACCAGGCCATCTAGAGCACCTGCTTGAAAACACACCTTCAACTCCAGCAGTGAAGCTGA
GCAGAAACACAGCGCTTCTACCTATCTTGCTACAGGTGAGCGCATTTGAGGAGCTGGGAGCCAGCCACTG
AGAGACCTCATTGAGAAAGATTGGTGTGGAACATTACGGGGCCCTGGACCCAGGACAACTTTATGGAGGTGT
GAAGGCAGTAGCAGGGACCTACAGGGCCACCCCTTCTCACCGTCTACATCAGTGTGACTCTAAGAGTTCCA
ACAGCAATGTTATCCAGGTGGACCAAGTCTGGCTCTTTCTGCCCTCTCGGATTACTACTTAAACAGAACTGCCA
ATGAGAAAGTGTCACTGCCTATCTGGATTACATGGAGGAACCTGGGATGCTGCTGGTGGCGGCCACCTC
CACGAGGGAGCAGATGCAGCAGGTGCTGGAGTTGGAGATACAGCTGGCCAACTACAGTCCCCCAGGACCCAG
CGCGCGACGAGGAGAAGATCTACCACAAGATGAGCAATTCGAGCTGCAGGCTCTGGCGCCCTCCATGGACT
GGCTTGAGTTCCTGCTTTCTGCTGTCAACATTGGAGTTGAGTGACTCTGAGCCTGTGGTGGTATGGGATGG
ATTATTTGCAGCAGGTGTCAGAGCTCATCAACCGCACGGAACCAAGCATCTGAACAATTACCTGATCTGGAACC
TGGTGCAAAAGACAACTCAAGCCTGGACCGACGCTTTGAGTCTGCACAAGAGAGCTGCTGGAGACCCCTAT
GGCACTAAGAAGTCTGTGTCGAGGTGGCAGACCTGCATCTCCAACACGGATGACGCCCTTGGCTTTGCTTT
GGGTCCCTCTTCGTGAAGGCCACGTTTGACCGGCAAGCAAGAAATTGCAGAGGGGATGATCAGCGAAATCC
GGACCGCATTTGAGGAGGCCCTGGACAGCTGTTTGGATGGATGAGAAGACCCGCCAGGAGCAAGGAGAA
AGCAGATGCCATCTATGATATGATTGGTTCCAGACTTTATCCTGGAGCCCAAGAGCTGGATGATGTTTATGA
CGGGTACGAAATTTCTGAAGATTCTTTCTTCCAAACATGTTGAATTTGTACAACCTCTCTGCCAAGGTTATGGCT
GACCAGCTCCGCAAGCCTCCAGCCGAGACCAGTGGAGCATGACCCCCCAGACAGTGAATGCCTACTACCTTC
CAACTAAGATGAGATCGTCTTCCCGCTGGCATCCTGCAGGCCCTTCTATGCCCGCAACCAAGGAGCC
CTGAACCTCGGTGGCATCGGTGTGTCATGGCCATGAGTTGACGCCATGCCCTTTGATGACCAAGGGCGCGAGTA

FIG. 1Y

TGACAAAGGAACCTGCGGCCCTGGTGGCAGAATGAGTCCCTGGCAGCCTTCCGGAACCACACGCGCTGC
 ATGAGGAACAGTACAATCAATACCAGGTCAATGGGAGAGGCTCAACGGCCGACAGCCTGGGGAGAACAA
 TTGCTGACAACGGGGCTGAAGGCTGCCTACAATGCTTACAAAGCATGGCTGAGAAAGCATGGGGAGGAGCA
 GCAACTGCCAGCCGTGGGCTCACCACCACAGCTCTCTTCTGGGATTTGCCAGGTGTGTGCTCGGTC
 CGCACACAGAGAGCTCTCACGAGGGCTGTGACCGAGCCCAACAGCCCTGCCCGCTTCCGCGTGTGGGC
 ACTCTCTCCAACCTCCGTGACTTCTCGGGCACCTTCGGCTGCCCTGTGCGCTCCCCCATGAACCCAGGGCAGCT
 GTGTGAGGTGTGGTAG

>SGPR303_SEQID_22

ATGCCGGAGAAGAGGCCCTTCGAGCGGCTGCCTGCCGATGTCTCCCCCATCACTGCAGCCTTTGCCTCAAGC
 CCGACTTGCTGGACTTCACCTTCGAGGGCAAGCTGGAGGCCGCCAGGTGAGGCGGACTAATCAGAT
 TGTGATGAATTGTGCTGATATTGATATTATTACAGCTTCATATGCACCAGAAGGAGATGAAGAAATACATGCTACA
 GGATTTAACTATCAGAATGAAGATGAAAAGTCACCTTGCTTTCCCTAGTACTCTGCAACAGGTACGGAACTT
 TAAAGATAGATTTTGTGGAGAGCTGAATGACAAATGAAGGTTTCTATAGAAGTAAGTATACTACCCCTTCTGG
 AGAGGTGCGCTATGCTGTAAACACAGTTTGAAGCTACTGATGCCGAAGGGCTTTTCTTGTCTGGGATGAGC
 GTGCTATCAAGCAACTTTTGATATCTCATTTGTTGTTCTAAAGACAGAGTAGCTTTATCAACATGAATGTAAAT
 GACCGGAAACCATACCCTGATGATGAAATTTAGTGGAAGTGAAGTTGCCGACACCTGTTACATCTACATAT
 CTGGTGGCATTGTTGTGGTGAAATATGACTTTGTAGAAACAAGGTCAAAAGATGGTGTGCTGTGTTTAC
 ACTCCTGTTGGCAAAGCAGAACAAAGGAAATTTGCATTAGAGGTTGCTGCTAAACCTTGCCTTTTTATAACGACT
 ACTTCAATGTTCTTATCCTCTACCTAAATTTGATCTCATTTGCTATTGCAGACTTTCAGCTGGTGCCATGGAGAA
 CTGGGACCTTGTTACTTATAGGGAGACTGCATTGCTTATTGATCCAAAATTCCTGTTCTCATCCCGCCAGTGG
 GTTGCTCTGTTGTGGACATGAACCTTGCCCATCAATGGTTTGAAATCTTGTTACTATGGAATGGTGGACTCAT
 CTTTGGTTAAATGAAGGTTTGCATCCTGGATTGAATATCTGTGTAGACCACCTGCTTCCAGAGTATGATATTT
 GGACTCAGTTTGTCTGTGATTACACCCGTGCCAGGAGCTTGACGCCCTTAGATAACAGCCATCCTATTGAAG
 TCAGTGTGGGCCATCCATCTGAGGTGATGAGATATTGATGCTATATCATATAGCAAGGTGCATCTGTCTATCC
 GAATGCTGCATGACTACATTGGGGATAAGGTAAAAAAAACCTTAAAGTATT

>SGPR402_1_SEQID_23

ATCGGGCCCGCCCCGATTGCGCTGTGGCTGCGCCTGGTCTTGCCCTGGCCCTTGCCGCCCGGGCTGTG
 GGGTGGGGCCCGTCCGAGCCCCCATCTATGTACGAGCTGGCCGTCCAGGTGTCCAGGGTAACCGGGAG
 GTCGAGCGCTGGCACGCAAAATTCGGCTTCGTCAACCTGGGGCCGATCTCCCTGACGGGCAGTACTTTCACCT
 GCGGCACCGGGCGTGTCTCAGCAGTCCCTGACCCCGCACTGGGGCCACCACTGCACCTGAAGAAAAACCC

FIG. 1Z

CAAGGTGCAGTGGTTCCAGCAGCAGCGCTGCAGCGGGGGTGAAACGCTCTGTGCTGGTGCCACGGACCC
CTGTTCTCCAAGCAGTGGTACATGAACAGCGAGGCCCAACAGACCTGAGCATCTCTGAGGCCCTGGAGTCAG
GGGCTGTACAGGCCAGGGCATCGTGGTCTCTGTGCTGGACGATGGCATCGAGAAGGACCAACCCGGACCTCTGG
GCCAACTACGACCCCTGGCCAGCTATGACTTCAATGACTACGACCCGACCCAGCCCGCTACACCCCA
GCAAGAGAACCGCAGCGGACCCGCTGTCTGGGAGGTGGCCGCGATGGCCACAATGGCTTCTGTGTG
TGGGGTCTGCTTCAACGCCCGGAATCGAGGCGTACGGATGCTGGACGGTACCATCACCGATGTCATCGAGGC
CCAGTCGCTGAGCCTGACGCCGACACATCCACATTTACAGCCGACGCTGGGTCCGAGGACGACGGCGG
CACGGTGGACGGCCCGCATCTCACCCGAGGGCTTCCGGCGTGGTGTACCAAGGGCCGCGGGGCT
GGCACGCTCTTCATCTGGGCTCGGGCAACGGGCGCTGCATACGACAACGAACTGCAACTGCGACGGCTACACC
AACAGCATCCACACGCTTCCGTGGGAGCACCACCCAGAGGCGCGTGCCTGGTACAGCGAAGCCTGCG
CCTCACCTCACCACTACAGCAGCGGCTGGCCACCCAGACCCCGATCGTACACGAGACCTGCATCA
CGGGTGACAGACGACACGGGCACTCGGCTCAGCCCACTGGCGCGGCGCATGATCGCCCTAGCGCT
GGAGGCCAACCCGTTCTGACGTGGAGAGACATGCAGCACCTGGTGTCCGCGCTCCAAGCCGGCGCACCT
GCAGCCGAGGACTGGAGGACCAACGGCGTGGGGCGCAAGTGAGCCATCACTACGGATACGGGCTGCTGGA
CGCCGGGCTGCTGTGGACACCGCCCGACCTGGTGGCCACCCAGCCGAGAGAAAGTGGCGCTCCGGGT
CCAGAGCCGCCACCCCATCTGCCGCTGATCTACATCAGGGAACGATCGGCTGCGCCGCGCTCCAC
AACTCCATCCGCTCGTGGAGCAGTGCAGGCGCAGCTGACGCTGTCTACAGCCGCGCGGAGACCTGGAG
ATCTCGCTACACGCCCCATGGCAGCGCTCCACACTCGTGGCCATACGACCCCTGGACGTCAGCACTGAAG
GCTACAACAACTGGTCTTCATGTCCACCCACTTCTGGGATGAGAACCCACAGGCGTGTGGACCCCTGGGCCCTA
GAGAACAAAGGCTACTATTTCAACACGGGGACGTTGTACCGCTACACGCTGTGCTCTATGGACGGCCGAGG
ACATGACAGCGCGGCTACAGGCCCGCAGGTGACGAGCGCGTGTGTGACGGGACACAGAGGGGCTGT
GCCAGGCGTGTGACGGCCCGCTACATCCTGGGACAGCTCTGCCCTGGCTACTGCCCCCGGGTCTTCAA
CCACACAAGGCTGGTACCGCTGGGCTGGGACACAGGCGGCGCGCTGAGGCTGTGCTCCAGCTGCCA
TGCCTCCTGTACACCTGCCGCGGCTCCCGAGGACTGCACCTCTGTCCCCCATCTCCACGCTGGAC
CAGCAGAGGCTCCTGATGGAGCCACACCCCGACAGCCCGCCGCTTAGAGCTGCCGCTGTCCC
CACCACCGCTGCCAGCCTCGGCCATGGTGTGAGCCTCCTGGCGGTGACCCCTCGGAGGCCCGCTCCTCTGC
GGCATGTCCATGGACCTCCCACTATACGCTGTGGCTCTCCCGTGCCAGGGCCACCCCAACCCCAAGTCT
GGCTGCCAGCTGGAACTGA

>SGPR434_SEQID_24

GGCCCGGCGCAGGCGGTGGTGCGCAGGGAGGCGTAGCACTGCTCTTCCCTCCGCGCTCCCTCAGGGCC
AGGCGCCAGGACCCCGAGCGGATGGAGCCGCCACCTGTAGGGGCTCCAGGATCCCCAGCGGGCC

FIG. 1AA

CCCAGTCCAGGGGAACGCAGTGCGCCCGCTTCGGTGTTACTTCCCTCAGCCTGTGGCCAGCGGACTTCAAG
 GATAACTGGAGGATTGCCGGCTCCAGACAGGAAGTGGCCCTGGCAGGTGAGCCTGCAGACCAGCAACAGACAC
 ATCTCGGAGGCTCCCTTATCGCCAGACACTGGGTATAAGAGGACACAACTCAAGTTGTGTGAGCCCC
 TGGTGTGCGAGGATTTGGAAATGACCCGCCATTGGCCCTGGAGGTGAGCCTCCGGATGGAATGAGCACG
 TGTGTGAGGGGCCCTCATTGACCCAGCTGGGTGGTGACTGCGGCCACTGCAGCCAAGGCACCAAGAGTA
 CTCAGTGGTCTTGCCACCTCCAAGCTGCAGCCCATGAACCTCAGCAGGGCCCTCTGGTCCCTGTGAGGGAC
 ATCATTATGCACCCCAAGTACTGGGCCGGCCCTTCATCATGGTGAGCTTGCCCTTGTCACCTTCAAACACCT
 GTCACCTTCAGTGAGTACGTGCAGCCCATCTGCCCTCCGGAGCCCAATTTCAACCTGAAGGTTGGGACGCAGTG
 TTGGGTGACTGGCTGGAGCCAGGTTAAGCAGCGCTTTTCAGGCTCCACAGCCAACCTCATGCTGACCCAGAGC
 TGCAGGAGGCTGAGGTGTTATCATGGACAACAAGAGGTGACCCGGCATTAACAAGAACTCCTTCTCCCCCA
 GTTGTCCCTTGCTGCTGGGGACATGATCTGTGCCACCAATTTATGGGAAACCTTGCTATGGGATTCTGG
 AGGGCCATTGGCTTGTAAGTTGAGGGCAGATGGATTCTGGCTGGGTGTTGCTGGGAAAGGCTGCGTG
 AAGGCACAGATCCAGGTGTGTACACCCGCTACCCAAATACACCAATGGATCAAGAAGCAATGAGCAATGG
 AGCCTTCTCAGGTCCCTGTGCCCTCTGCCCTGCTGCTGGCCGCTGCAGCCCCAGATGGGCTCCT
 GA

>SGPR446_1_SEQID_25

ATCCTCACCCAGTGTGTGGCCGAACCCCTCTGAGATCGTGGGAGGAGTGACGCGGAGGAAGGGAGGTGG
 CCCTGGCAGGTGAGCGTGAGGACCAAGGCAGGCACATCTCGCGCGCACCTGGTCACCGCCACGTGGGTG
 CTGACGGCAGGCCACTGCATTTCCAGCCGTTTCCATTACAGTGTCAAGATGGGAGATCGGAGTGTCTATAATGAA
 AACACAAGTGTGTCTCAGTCCAAGAGCTTTGTCCACCCCTAAGTTCTCAACAGTTACAACCATTCGAAAT
 GACCTTGCCCTTCTCCAGCTCCAACATCCTGTGAATTTACCTCAACATCCAGCCTATCTGCATCCCTCAGGAG
 AATTTCCAGGTGGAAGTAGGACCAAGGTGCTGGTGACCGGATGGGGCAAAACACCAGAACGTGGAGAGAAAC
 TTGCATCAGAAATCTTCAGGATGTGGACCAATACATCATGTGCTATGAGGAATGTAATAAGATAATACAGAAGGC
 CTGTGCTACTACTAAGGATGTAATAATAAAGGGATGGTCTGTGGCTATAAGAACAAAGGAAGGATTCTTGTCAA
 GGAGATTCTGGGGCCGCTTGCCCTGTGAATAATGACACATGGGTCCAGGTAGGATTGTGAGCTGGGGCA
 TCGGCTGTGGTCGC

>SGPR447_SEQID_26

ATGGCGCGCGCGGGCGCTGCTGCTGGCGCTGCTGCTGGCTCGGGCTGGACTCGGGAAGCCGGAGGCCCTG
 CGGCCACCGGAAATTCACGCGCTGGTGGCGGGCGGAGTGGAGTCCGCGCGGCGCTGGCCATGGCAGG
 CCAGCCTGCGCCTGAGGAGACGCCACCGATGTGGAGGGAGCCTGCTCAGCCGCCGCTGGGTGCTCTCGGCTG

FIG. 1BB

CGCACTGCTTCCAAAAGCACTACTATCCCTCCGAGTGGACGGTCCAGCTGGCGAGCTGACTTCCAGGCCAACT
CCTTGGAACCTGCGGCCCTACAGCAGTCGTTACAAAGTGCAGGACATCATTTGTGAACCCCTGACGCACCTTGGGGT
TTTACGCAATGACATTGCCCTGCTGAGACTGGCCTCTTCTGTACCTACAATGCGTACATCCAGCCCCATTGTCAT
CGAGCTTCCACCTTCAACTTCGTGCACCGCGGACTGCTGGTGACCGCTGGGGTTAATCAGCCCCCAGT
GGACACCTCTGCCACCTCCTTACAACCTCCGGGAAGCACAGGTCAACCATCTTAAACAACACCAGGTGTAATTAC
CTGTTTGAACAGCCCTCTAGCCGTAGTATGATCTGGGATTCATGTTTGTGCTGGTCTGAGGATGGCAGTGTA
GACACCTGCAAAAGGTGACTCAGGTGGACCCCTTGTCTGTGACAAGGATGGACTGTGGTATCAGGTTGGAATCGT
GAGCTGGGAATGACTGCGGTCAACCCAAATCGGCCTGGTGTCTACACCAACATCAGTGTGTACTTCCACTGGA
TCCGGAGGGTGATGTCCACACAGTACACCCAGGCCAAACCCCTCCAGCTGTGCTGCTCCTTGCCCTGCTGTGG
GCTCCC

>SGPR432_1_SEQID_27

ATGGGCAGCACCTGGGGAGCCCTGGCTGGGTGCGGCTCGCTCTTTGCCTGACGGGCTTAGTGCTCTCGCTCT
ACGCGCTGCACGTGAAGCGCGCGCGCGCGGACCGGATTACCGCGGCTCTGCGACGTGGCACCAGCC
ATCAGCTGTTCCGCGCTTCTCCTCCAGGTGGGCAGGGGTTTCGGGCTGGTGAGCATGTGCTGGGACAGG
ACAGCATCTCAATCAATCCAACAGCATATTGCGTTGCATCTTACACACTACAGCTATTGTTAGTCTTCAAGC
CGCTACGCTGCGCTGTGGACAGCGTGGCCCGCGCCCGCCCAAGCCTCAGGAGGCAACACAGTCCCTGGCGA
GTGGCCCTGGCAGGCCAGTGTGAGGAGGCAAGGAGCCACATCTGCAGCGGCTCCCTGGTGCCAGACACCTG
GGTCTCACTGCTGCCACTGCTTTGAAAGGCAGCAGCAACAGAACTGAATCTGTGTCAGTGTCTGGTT
CTCTGACGCTGAGGACTCAGCCCTGGGCGCGAAGAGTGGGGTGGCTGCCCTGCAGTTGCCAGGGCCT
ATAACCACTACAGCAGGGCTCAGACCTGGCCCTGCTGCAGCTCGCCCAACCCACGACCCACACACCCCTCTG
CCTGCCCCAGCCCGCCATCGCTTCCCTTTGGAGCCTCCTGTCTGGGCCACTGGCTGGATCAGGACACCACT
GATGCTCCTGGGACCCCTACGCAATCTGCGCCTGCGTCTCATAGTCGCCCCACATGTAACCTGTATCTACAACCA
GCTGCACCAAGCACCTGTCCAACCCGGCCCGCTGGATGCTATGTGGGGCCCCCAGCCTGGGGTGCA
GGCCCTCTGTCAGGGAGATTCCGGGGCCCTGTGCTGTGCTCGAGCCTGACGGACACTGGGTTACGGCTGG
CATCATCAGCTTTGCATCAAGCTGTGCCAGGAGGACGCTCCTGTGCTGCTGACCAACACAGCTGCTCACAGTT
CCTGGCTCAGGCTCGAGTTCAGGGGCGAGCTTCTGGCCAGAGCCCCAGACCCCGGAGATGAGTGATGA
GGACAGCTGTGTAGCCTGTGGATCCTTGAGGACAGCAGGTCCCCAGGAGGAGCACCCTCCCCATGGCCCTGG
GAGGCCAGGCTGATGCACAGGGACAGCTGGCCTGTGGCGGAGCCCTGGTGTGAGGAGGCGGCTGCTAACT
GCTGCCACTGCTTCAATTGGGCGCCAGGCCCCAGAGGAATGAGCGTAGGGCTGGGACCAACAGCCGAGGAG
TGGGGCCTGAAGCAGCTCATCTGATGGAGCTACACCCACCCCTGAGGGGGGCTACGACATGGCCCTCCTGC
TGCTGGCCCCAGCCTGTGACACTGGGAGCCAGCCTGCGGGCCCCCTCTGCCTATCCTGACCAACACCTGCC

FIG. 1CC

TGATGGGAGCGTGGGTTCTGGACGGGCGCCAGGAGCAGGCATCAGCTCCCTCCAGACAGTGCC
CGTGACCCCTCTGGGGCTAGGGCTGCAGCCGGCTGCATGCAGCTCCTGGGGTGATGGCAGCCCTATTCTG
CCGGGATGGTGTACAGTGTGTGGTGAGCTGCCAGCTGTGAGGGCTGTCTGGGGCACCACACTGGTG
CATGAGGTGAGGGGCACATGGTTCTTGGCCGGCTGCACAGCTTCGGAGATGCTTGCCAAAGGCCCGCCAGG
CCGGCGGTCTTCACCGCGCTCCCTGCCTATGAGGACTGGGTGAGCAGTTTGACTGGCAGGTCTACTTCGCCG
AGGAACAGAGCCGAGGCTGAGCCTGGAGCTGCCCTGGCCAAACATAAGCCAACCAACAGCTGCTGA

>SGPR529_SEQID_28

ATGAGAGCTCCGACCTCCACCTCTCCGCGCTCTGGCGCCGGGCTCTGGCGAAGCTGCTGCCGCTGCTGA
TGGCGCAACTCTGGCGCGCAGAGCGGCGCTGCTCCCCAAACGACACGCGCTTGACCCCGAAGCCTATG
GCGCCCGTGGCGCGGCTCGCAGCCTGGCAGGTCTCGCTCTTCAACGGCCTCTCGTTCCACTGCGCGG
GTCTCCTGGTGACAGAGTTGGTGCTGACGGCCGCGCACTGCGGAAACAAGCCACTGTGGGCTCGAGTAG
GGGATGACCACCTGCTGCTTTCAGGGCGAGCAGCTCCGCCGACGACTCGCTGTGTCTCCATCCCAAGTAC
CACCAGGCTCAGGCCCATCTGCCAAGGCGAAGGATGACGACGATCTCATGTTGCTAAAGCTGGCCAGGC
CCGTAGTCCCGGGCCCGCTCCGGCCCTGCAGCTTCCCTACCGCTGTGCTCAGCCCGGAGACCAGTGCC
AGGTTGCTGGCTGGGCACCAACGCGCCCGGAGAGTGAAGTACAACAAGGCGCTGACCTGCTCCAGCATCAC
TATCCTGAGCCCTAAAGAGTGTGAGGTCTTCTACCTGGCGTGTACCAACAACATGATATGTGTGACTGG
ACCGGGCCAGGACCCCTTGCCAGAGTGAAGTCTGGAGGCCCGCTGTGTGACGAGACCCCTCCAAGGCATCCT
CTCGTGGGGTGTACCCCTGTGGCTCTGCCAGCATCCAGCTGTCTACACCCAGATCTGCAATATACATGTCT
GGATCAATAAAGTCATACGCTCCAACTGA

>SGPR428_1_SEQID_29

AAACGTACAATGTGGCCACAGGCCTGCTTTTCCAAACTCGTCATGGTTACCAATTCATGAACGGCTTCAAGTCC
AGAAATGGTGAGTGCCCGTGGAAGTGAGTATCCAGATGTACGGAACACCTCTGTGGAGGCTCAATCTTACAT
TGGTGGTGGTTCTGACAGCCGCACACTGCTTCCGAAGAACCCTATTAGACATGGCCGTGGTAAATGTCACGT
GGTCATGGGAACGAGAACATTCAGCAACATCCACTCGGAGAGAAAGCAAGTGCAGAAGTCAATTACACAATA
TTACAAACCGCCCCAGCTCGACAGTGACCTCTCTGCTTCTACTTGCCACACCAAGTGAATTCAGCAATTTCAA
AATGCCCTGTGCTGAGGAGGAGGAGGAGGACCTGGGACTGGTGTGGATGGCACAGTGGGTAAACGACCAAT
GGGTATGACCAATATGATGACTTAAACATGCACCTGGAAAAGCTGAGAGTGGTGCAGATTAGCCGGAAGAATG
TGCCAAGAGGGTAAACAGCTGTCCAGGAACATGATTTGTGCTTGGAAACGAGGACCAATGGGCAGGGC
CCAGGAGAAGTAGGGGGCCTCTGTTTCCAGAAAAGAAAGACATGGTACCAGCTGGGTATTATCAG
CTGGGGTGTGGGCTGTGGCCAGAAGAACAATGCCTGGAGTGTACACCGAGTTGTCCAATTTATCTGCTTTGGATCG

FIG. 1DD

AGAGGAAGACTGTGCTGGCAGGAAGCCGTATAAGTATGAGCCAGACTCTGTGTACGCTTTGCTTCTCACCC
TGGGCCATCCTGTTACTGTATTTTGTGATGCTTCTATTATCCT

>SGPR425_SEQID_30

ATGGAAAATATGCTGCTTTGGTTGATATTTTACCCCTGGGTGGACCCCTCAATTGATGATCTGAAATGGAATGG
GATTTTATGTGGCACTTGAGAAAGGTACCCCGGATTGTGAGTGAAAGGACTTTCCATCTCACAGCCCGCATTT
GAGGCAGATGCTAAGATGATGTTAAATACAGTGTGTGGCATCGAATGCCAGAAAGAACTCCCAACTCCAGCCT
TTCTGAATTGGAGGATTATCTTCTCTATGAGACTGTCTTTGAGAAATGGACCCGAACTTAACAGGTTGAAAGTT
CAAGATTTGGTTCTTGAGCCGACTCAAAATATCACCAAAAGGAGTATCTGTTAGGAGAAAGACAGGTGTAT
GGCACCGACAGCAGGTTACGATCTTGGACAAAAGTTCTTAACCAATTTCCCTTTAGCACAGCTGTGAAGCTT
TCCACGGGCTGTAGTGGCATTCTCAITTCCTCTAGCATGTTCTAACTGTGCCACTGTGTTTCATGATGGAAG
GACTATGTCAAAGGGAGTAAAGCTAAGGGTAGGGTTGTTGAAGATGAGGAATAAAAGTGAGGCAAGAAACG
TCGAGGTTCTAAGAGGAGCAGGAGAGAGCTAGTGGTGACCAAAGAGAGGTACAGAGAGCATCTGCCG
GAGAGCGGAAGGTGGGAGAAGAAAGAAATCTGCCGGGTGAGAGGATTGCCAAGGAGGCTTCTCT
TTCAGTGGACCCGGTCAAGAATACCCACATTCGAAAGGCTGGGCACGAGGAGCATGGGGACGCTACCTT
GGACTATGACTATGCTCTTCTGGAGCTGAAGCGTGCTCACAAAAGAAATACATGGAAC TTGGAATCAGCCCAAC
GATCAAGAAAATGCCGTGGTGAATGATCCACTTCTCAGGATTTGATAACGATAGGCTGATCAGTTGGTCTATCG
GTTTTGCAGTGTGTCGACGAATCCATGATCTCTTTACCAATACTGCGATGCTGAGTCGGGCTCCACCGGTTT
GGGGTCTATCTCGTCTGAAGATCCAGACAAAAGAAATTGGAAGCGCAAAATCATTTGCGGTCTACTCAGGGC
ACCAGTGGGTGGATGCCACGGGTTCAGAAGGACTACAACGTTGCTGTTCCGATCACTCCCCATAAAATACGCC
CAGATTTGCCCTCTGGATTACGGGAACGATGCCAATTGTGCTTACGGCTAA

>SGPR548_SEQID_31

ATGGGGACCCAGAAAGGAGCGCAGAGTGGGGTTGGGGAGGGGATACCGGTGGTCAGAAGAAATTTATTAA
CAGTGGATGGGATAAGTCTGTCTGGAGGGATCCTGGTGAGGAGGAGGTCCTGCCTCACCTGGATTCTCT
CACTCCCTCCCGAGACTGCAGCCGAACCCCTGGTCCCTCCTCCACAATGTGGCTTCTCCTCCTCTCCTTCTG
CTGGCATCCACAGCAGCCAGGATGGTGACAAGTTGCTGGAAGGTGACGAGTGTGACCCCACTCCAGCCAT
GGCAAGTGGCTCTCTACGAGCGTGGACGCTTTAACTGTGGCGCTTCCCTCATCTCCCCACACTGGGTGTCT
GGGCCCCACTGCCAAAGCCGCTTCATGAGAGTGGCCTGGGAGAGCACAACTGCGCAAGCGCGATGGCCCA
GAGCAACTACGGACACGTCTCGGGTCAATCCACACCCCGCTACGAAGCGCGAGCCACCGCAACGACATCA
TGTGCTGCGCCTAGTCCAGCCCCGACGCTGAACCCCCAGGTGCGCCCCCGGTGCTACCCACGCGTTGCC
CCACCCGGGGAGGCCTGTGTGCTGGCTGGGCGCTGTGTGCCACAACGAGCCTGGGACCGCTGGGAG

FIG. 1EE

CCCCGGTCAAGTGAGTCTCCAGATACGTTGCATTGTGCCAACATCAGCATTATCTCGGACACATCTTGTGA
 CAAGAGCTACCCAGGGCCCTGACAAACACCATGGTGTGTGCGAGCGCGGAGGCGAGGCGCAGAAATCCTG
 TGAGGGTGACTCTGGGGACCCCTGGTCTGTGGGGCATCCTGCGAGGCAATTGTCTCCTGGGGTGACGTCCCT
 TGTGACAAACACCACCAAGCCTGGTGTCTATACCAAAGTCTGCCACTACTTGGAGTGGATCAGGGAAACCATGAA
 GAGGAACTGA

>SGPR396_SEQID_32

ATGGGCCCTGTGGCTGTGCCCTTCAAGCTGTCTCTGTCTGGGATCTCAGTGTGTGGGCAACCTGTATACTC
 CAGCCGCGTTGTAGTGGCCAGGATGCTGCTGCAGGGCGCTGGCCTTGGCAGGTGAGCCTACACTTTTGACCCAC
 AACTTTATCTGTGGAGGTTCCCTCGTCAGTGAGAGGTTGATACTGACAGCAGCACACTGCATACAAACCGACCTG
 GACTACTTTTTCATATACTGTGTGGCTAGGATCGATTACAGTAGTGACTCAAGGAACGTGTGAAGTACTACGT
 GTCCAAATCGTCATCCATCCCAAGTACCAAGATACAAACGCGAGACGTGCGCTTGTGAAACTGTCCCTCTCAAGT
 CACCTTCACCTTCTGCCATCCTGCCTATTGTCTTGGCTTGGCAGTGTCAAAAGCAGTTGGCAATTCACCCCTTTTGTGG
 GTGACCGGATGGGGAAGTTAAGGAAAGTTACAGATAGAGATTACCAATCTGCCCTTCAGGAAGCAGAAAGTACC
 CATTATTGACCGCCAGCCTTGTGAACAGCTCTACAAATCCATCCGATCTTCTTCCAGCACTGGAGCCAGTCAT
 CAAGGAAGACAAGATTGTGCTGTGTGATACTCAAAACATGAAGGATAGTTGCAAGGGTGATTCTGGAGGGCCTC
 TGTCGTGTACATTGATGGTGTATGGATCCAGACAGGAGTAGTAAGCTGGGGATTAGAATGTGGTAAATCTCTTC
 CTGGAGTCTACACCAATGTAATCTACTACCAAAATGGATTATGCCACTATTTCAAGAGCCAAACAATCTAGACTT
 CTCTGACTTCTTGTCCCTATTGTCTCTACTCTCTGCTCTCTGCTGCTCCCTCTCTGCTTGGACCTTAACACT
 ATACACAGAGTAGGCACTGTAGCTGAAGCTGTTGCTTGCTATACAGGGCTGGGAAGAGAAATGCATGGAGATTAG
 TCCCAGGGGCAGATAA

>SGPR426_SEQID_33

ATGATGTACGCACCTGTTGAATTTTCAGAAGCTGAATTTCTCAGGAGCTGAATATCAAGAAGCAGCAATTTTGGG
 ACTCAGTAGCGTAGCTCTTTTACATTAGCAATTGTAGCAATCATAGGAATTGCAATTGGTATTGTACTCATTTT
 GTTGTGAGGATGATAAGTCTTTTCTATTACCTTGCCTCTTTAAAGTCACAAATATCAAAATATAAGAAATATGG
 CATAAGATCTTCAAGAGAGTTTATAGAAAGGAGTCATCAGATTGAAAGAAATGATGCTAGGATATTTTCGACATTC
 TCTGTAGCGGTCGATTTATCAATCTCATGTTATCAAAATTAAGTCCAGATGAACAAGGTGTGGATATCTTATAG
 TGCTCATATTTTCGATACCCATCTACTGATAGTGTGAACAAATCAAGAAATAATTGAAAAGGCTTTATATCAAAGT
 TTGAAGACCAACAATTGTCTTTGACCATAAACAACCATCATTTAGACTCACACGCTGTGGAATAAGGATGACAT
 CTTCAACATGCCATTACCAGCATCCTCTTCTACTCAAGAAATTGTCCAAGGAAGGAAACAGCTATGGAAGGGG
 AATGGCCATGGCAGGCCAGCCTCCAGCTCATAGGCTCAGGCCATCAGTGTGGAGCCAGCCTCATCAGTAACAC

FIG. 1FF

ATGGCTGCTCAGCAGCTCACTGCTTTTGGAAAAATAAGACCCCAACTCAATGGATTGCTACTTTTGGTGCAAC
 TATAACACCCAGCGAGTGAACGAAATGTGAGGAAAAATTATCTTCATGAGAAATACCATAGAGAAACAATGAA
 AATGACATTGCTTTGTTCAAGCTCTCTACTGAGTTGAGTTTCAAAATATAGTCCAGAGAGTTTGCCTCCAGACT
 CATCTATAAGTTGCCACCTAAACAAGTGTTCGTACAGGATTTGGATCCATTGTAGATGATGGACCTATACA
 AAATACACTTCGGCAAGCCAGAGTGAACCAATAGCACTGATGTGTAAACAGAAAGGATGTGTATGATGGCCT
 GATAACTCAGGAATGTTATGTGCTGGATTCTGGAAGGAAAAATAGATGCATGTAAGGGAGATTCTGTGGACC
 TCTGGTTTATGATAATCATGACATCTGGTACATTGTAGGTATAGTAAGTTGGGACAATCATGTGCACCTTCCCAA
 AAACCTGGAGTCTACACCAGAGTAACATAAGTATCGAGATTGGATTGCCTCAAAGACTGGTATGTAG

>SGPR552_SEQID_34

AGAATAGCAGAGGGTCTGGATGCTGAGGAAGGAGAAATGGCCCTGGCAAGCTAGCCCTCCACAGAACAAATGTCTA
 CCGACGGGAGCCACATGGCTTAGTAACAGCTGGCTTATCACTGCTGCTCACTGCTTCATAAGGGTCCATGATC
 CCAAAGAATGGAATGTTATTTAAGTAACCCACAACACAGTCAAAATATCAAGAAATGTTATAATTCAAGAAACTAC
 CATTACCCCTGCACACGATAATGACATTGCTGTTGTGCATCTATCTCACCAGTGTATATACAGCAACATCCAAA
 AAGCATGTCTTCCAGATGTTAATTATATATTCCTATACAATTCAGAAAGCAGTGGTACTGCATGGGATCATTTAA
 CCTTACGAACAACCTTCTAATGTACTCCACAAGGGATTAGTGAAGATTATAGATAATAGGACCTGCAACAATGGG
 GAGGCAGATGGCAGAGTCATCACATCTGGAATGTTGTGTCCGGTTCTGGAGCCACGTTGGATGCCCTGCC
 AGGGTGACTCTGGTGACCACCTGTTGGTACAGATTCTAAGGCATCCTTGCTAAAGTTCCCTGCTGCTGATTGA
 AAGCTGGAGTAAATGAACGTGCTCTTCCAAACAAGCCTAGTGTCTACACTCAAGTGACATACTAT

>SGPR405_SEQID_35

ATGGTCAGCAAGGGGGAGTTGCTGCAGAGCCAGAGCCACACTATTGTGAGGACAGTGAAAGAGGCCCAACA
 CCTCACAGGTCGGGAGCCCTTCTAGAGGAGGTGGCATTGAGGTGGCATGGAGTTTCCGGGATGCAGCGG
 TGAAGGTGCGTGAAGCCCATGAGGAGGCGGCCGGGAGGGGCGGCGAGGCAAGAGGGCTGTGCCGG
 GACCCAAAGCAGCGCAGGGGTGAGCAGAGGGGCTCGCGGGGTGGACGCTGGAGCAGGAGACCAGG
 GGAGATGCTTAGAGGATAAAAATGAGCGGCAGATGAAGAGATACTCAGGCTGGCACCAGGAAAGGCAGGC
 TCCCAATAGACAGCAACACCTGAAACCGGTGATCAGCAGCTTCCCGTAAGATCTCAGGAGCTGGCGAGGG
 GGCTGGAGCAGGCACACTAAGAGGCAAAATGGCAGAGTTAACTGCTCTATGGCCTTCAAGGGACCTGCGGCT
 GGTCATGAAGAGCGCTCAACTCTGTGTCCAGCAGGCGCAAGAGGCAATTGGCTGGGATGTCGCTGCTGCTT
 CTCTTCGTGGTTGACCATTTCTCAGACCTCCCCCGCCCTGCAGGTGAGGAGGAGTTGGAGGCTTGCGC
 GTTAGAGTGCAGGTGGGCGAGCTGAGGCTCTATGAGGACGACCGGACGAAAGGTGGTTGAGATCGTCCGT
 CACCCCAAGTACAACGAGAGCCTGTCTGCCCAAGGCGGTGCGGACATCGCCCTGCTGAAGCTGGAGGCCCGG

GTCCCGTGTCTGAGCTCATCCACCCGGTCTCGCTCCCGTCTGCCCTGACGTGCCCTCGGGGAAGACCT
GCTGGTGACCGGTGGGTGTCAATTGACGTGGAGAACTACTGCCCTGCCCTCAGCTTGTGGAGGGCAG
GGTGAAGTCAAGGACAACGTCTGTAAACAGACCTGTCCGCCCGCTTCTTCCAACACACTGAGCGGT
TTGAGCGGCTCATCAAGGACGACATGCTGTGTCCGGGACGGAAACACGGCTCCTGGCCAGCGACAACG
GGGGCCCCCTCTGTGCAGGCGGAATTGCACCTGGGTCCAGGTGGAGTGGTGAGCTGGGCAAACTCTGCC
GCCCTCGCGGTATCCCGCATGTACACCGGTACAGAGTACGTGCTGTGATCCGCCAGCCATGCCCTC
AGCTCAGACCCCTGTGTGTCGGAAGATTTGTCTCCCCCAATCCAGATGTTGAAGCCCTAACTCCCAGTG
TGATGGGATCAGGAGCGCGCTGCCCGCCCCGACCTGCAAGAGGCCGAGGTCCCCATCATGAGGACCC
GAGCTGCGAGAGGATGTATCAAAAGGCCCACTGCCACGCCAGGTCAACATCATCAAGGTGCCATGCC
GTGTCAGGAGGAAGGGCAGGGTTCTGCCAGCCGCTCTGAGGACGGAGGACCTCACCCCAACCAACC
CAACACGGAGGTCTCCACGTGCAGACCCAGGCTGAGCCAGCCGAGGACATCTGCCAGAGTGGCTTG
GCCAGTTGTGGTGCCACCACCATGCTGTCTGTCTGTCTTCTGGCTGTCTCTCCCTGGGAGCTGTAGC
ACTGGAGTCCAGTCCCTCCCGAGAAATGACCTGGTGGCATTGTGGGGGCCACAACACCCAGGGGAA
GTGTCGTGGCAGTGGTGTGACCGCGCTCACTGCAATTTCCGGAAGGACACCGACCCGTCCACCTACCGG
ATTACACCAAGGATGTGTATCTGTACGGGGCCGGGGCTGCTGAATGTCAGCCAGATCGTCTCCACCCAA
CTACTGTCTTCTTCTGGGGCAGACATCGCCCTGCTGAAGCTGCCACCAAGTCCCTGGAGTTCAGTGACA
GTGACAACTGCTGGAACACAGGCTGGGCATGCTGGCTTGTGGATATGCTGCCGCTCCTTACCGCCCGCA
GCAGGTGAAGTCTCACACTGAGCAATGCAGACTGTGAGCGGACACCTACGATGCTTTTCTGGTCTGGAG
ACAGAAAGTTTCATCCAGGATGACATGATCTGTGCCGGCCGACGGCCGCCACCTGGAAGGTTGACTCAGG
CGGCCCCCTGGTCTGCAAGAAGAGGTACCTGGCTCCAGCGGGAGTAGTGAGCTGGGATTTACAGTGTAT
CGGCCACGATTGGCGTTACACACGCCACAGACAGCTGCAGGGTGCCAAACCATGCAGACGCCACAGAGAC
CAGCTGGCAGGTTCCAAACCATGCAGAGGCCCAGAGACATGGGCCAGGCCAGGAGTGGTCTGCAGGCCCT
TCACCCACGTACCTGTACCCGACGGCCATCCCCAGGCCCTTACCCATGTCACCTGTACCTGATGGCTGTC
CCCAGCACCTCACCCACGTACCTGTACCCGACGGCGCTCCACAGGCCCTTACCCATGTCACCTGCTACCT
GATGGCTGTCCCAGCACCTCACCCACATCACTGTCTACATGATGGCCGTCCCCAGGCCCTTACCCACATCA
CCTGTACCCCAATGGCTGTCCCCAGCACCTTACCACGTCACTGCCACCCGACGGCCATCCCCAGGCCCTTC
ACCCACATCACCTGTACACGATGGCCATCCCCAGGCCCTTCAACCAAGCCACCTGTCTACACGACGGCCATCCC
AGCACCCCTCACCCACGTCACTGTCTACACGATGGCCGTCCCCAGGCCCATCACCCATGTCACTGCTACACGAT
AG

FIG. 1HH

[illegible]

FIG. 111

>SGPR390_SEQID_38

ATGGAGCCCACTGTGGCTACACCTCGTGCCCAGGACAACCAAGGAAGTCCCCGCTCTGGATGCCGCGGT
GCTGTCGAGCGGCCAGCATTTGGCGTGGTGGCCACCAGCCTTGTCTCTCACCTCGGAGTCCTTTTGGGAGG
AATGAACAACCTCAGACACGCTGCCCTTAAGAGCTGCAACACTCCCTGGGAAGTCTACAGCGTCACTCCTGAAG
CAAGCAAGACCACGAACCCACCAAGGAAGAAATTCGGAACACATCCGAACATCAGCAAGAACAACACTCCGGA
CACACCATCTTTAAGAAATGTAACACTCAGCCCTTCTCTCTACACAGGCTTCCACGTGGACCACACGCGCCGAG
CTGCGGGGAATCCGCTGACCAAGCAGTTTGGCGGGAGACCTCGGACTATCACCGCACGCTGACGCCCCACC
CTGGAGGCACTGCTGCACCTTTCTGCTGCGACCCCTCCAGACGCTGAGCCTGGGCCCTGGAGGAGGAGCTATTGC
AGCGAGGGATCCGGGCAAGGCTGCGGAGCAGCGCATCTCCCTGGCTGCCCTATGGCACAATTGTGTCGGCTGA
GCTCACAGGGAGACATAAGGGACCTTGGCAGAAAGAGACTTCAATCAGGCCGCTGTCCAGGGAACCTCCTTTT
CCTGCGGGAACAGCCAGTGTGTGACCAAGGTGAACCCGGAGTGTGACGACCAAGGAGACTGCTCCGATGGGTC
CGACGAGCGCACTGCGAGTGTGGCTTGCAGCCTGCCCTGGAGGATGGCCGGCAGGATCGTGGCGGCATGGA
AGCATCCCCGGGGAGTTCCGTGGCAAGCCAGCCTTCGAGAGAACAAAGGAGCACTTCTGTGGGCGCGCCATC
ATCAACGCCAGGTGGCTGGTGTCTGCTCACTGCTTCAATGAGTTCCAAGACCCGACGAAGTGGTGGCCTA
CGTGGGTGCGACCTACCTCAGCGGCTCGGAGGCCAGCACCGTGGGGCCAGGTGTCAGATCGTCAAGCA
CCCCCTGTACAACGGGACACGGCCGACTTTGACGTGGCTGTGCTGGAGCTGACCAGCCCTCTGCCTTTCCGGC
CGGCACATCCAGCCCGTGTGCCTCCCGCTGCCACACACATCTTCCACCCAGCAAGAGTGCCTGATCTCAG
GCTGGGGCTACCTCAAGGAGGACTTCCGTAAGCATCTTCTCGGCTGCAATGGTCAAGCCAGAGGTGCTGCA
GAAAGCCACTGTGGAGCTGCTGGACCAGGCACCTGTGCCAGCTTGACGGCCATTCACTCACTGACAGGATG
GTGTGCGCTGGCTACCTGGACGGGAAGTGGACTCCTGCCAGGTGACTCAGGAGGACCCCTGGTCTGCGAG
GAGCCCTCTGGCCGGTCTTTCTGGCTGGCATCGTGAGCTGGGAATCGGTGTCGCGAAGCCCGGCGTCCAG
GGGTCTATGCCCCAGTCAACAGGCTACGTGACTGATCCTGGAGGCCACCACCAAGCCAGCATGCCTCTGGC
CCCCACCATGGCTCCTGCCCTGCCGCCAGCACAGCCTGGCCACCACTGCTGAGAGCCCTGTGCTCAGC
ACCCCCACCAATCGATGCAGGCCCTCAGTACCCTGCTCTTGACTGGTCACCGTTCTCTAAGCTACAAGAATG
TGGGGCCAGGCTGCAATGGAGAAGCCACCCGGTCTGTGGCGGTTTCGGAGCTGCCTCCGGGGAGGTGC
CCTGGCAGGTACGCTGAAGGAAGGTCCCCGGCACTTCTGCGGAGCAACTGTGTGGGGACCGCTGGCTGC
TGCTGCGCGCCCACTGCTTCAACCACACGAAGGTGGAGCAGGTTCCGGCCCACTGGGCACCTGCGTCCCTCCT
GGGCTGGCGGGAGCCCGTGAAGATCGGGCTGCGCGGGTAGTGTGCACCCCCCTTACAACCCCTGGCAT
CCTGGACTTCGACCTGGCTGCTGGAGCTGGCCAGCCCCCTGGCCTTCAACAAATACATCCAGCCTGCTGCC
TGCCCCCTGGCCATCCAGAAGTTCCCTGTGGCGCGGAAGTGCAATGATCTCCGGATGGGAAATACGCAGGAAGG
AAATGCCACCAAGCCCGAGCTCTGCAGAAGGCGTCCGTGGGCATCATAGACCAGAAAAACCTGTAGTGTGCTCT

FIG. 1JJ

ACAAATTCTCCCTCACAGACCGCATGATCTGCGCAGGCTTCCTGGAAGGCAAAAGTCGACTCCTGCCAGGGTGAC
 TCTGGGGCCCCCTGGCTGCGAGGAGGCCCTGGCGTGTATCTGCGAGGGATCGTAGCTGGGTATTG
 GCTGCGCTCAGGTTAAGAAAGCCGGCGGTACACGGCATCACAGGCTAAAGGGCTGGATCCTGGAGATCAT
 GTCTCCAGCCCTTCCCATGTCTCCCCCTCGACCACAAGGATGCTGGCCACCACCCAGCCCGGACGACA
 GCTGGCCTCACAGTCCCGGGGCCACACCCAGACACCCCTGGGCTGCCAGAGGTGACGGGCCAA
 CCTGCCAACTCAACCTTATCTGCCGTGAGCACCACTGCTAGGGACAGACGCCATTTCCAGACGCCCGGAGG
 CCACCACACACCCAGCTACCAGACTGTGGCTGGCGCCGCCGCTCACAGGATTGTGGCGGCAGCG
 CAGCGGCCGTGGGAGTGGCCGTGGCAGGTAGCCTGTGGCTGGCGCCGGGAACACCGTTGCGGGGCC
 GTGCTGGTGGCAGAGGTGGCTGTGTCGGCGCACCTGCTTGACGTCTACGGGACCCCAAGCAGTGG
 GCGCCTTCTAGGACGCCGTTCTGAGCGCGGAGGGCAGCTGGAGCGGTGGCGCGCATCTACAAG
 CACCCGTTCTACAATCTTACACGCTCGACTACGACGTGGCGCTGCTGGAGCTGGCGGGCCGTCGTCGCA
 GCCGCTGGTGGTCCCATCTGCCCTGCCGAGCCCGCCGACCCCGACGCGCACGCGCTGCGTCATCA
 CCGCTGGGCTGGTGGCGAAGGAGGCTCCATGGCGCGCAGCTGCAGAAGCGCGCGCTGCCCTCCTC
 AGCGAGCAGACCTGCCGCCCTTCTACCCAGTGCAGATCAGCAGCCGCTGCTGTGTCGGCTTCCCGCAGG
 GTGGCGTGGACAGCTGCTCGGTGACGCTGGGGACCCCTGGCTGACGGAGCCCTCTGGACGGTGGTG
 TAACTGGGTACACTAGCTGGGGCTATGGCTGTGGCGGCCCACTTCCAGGTGCTATACCCGGGTGGCAGC
 TGTGAGAGGCTGGATAGGACAGCACATCCAGGAGTGA

>SGPR521_SEQID_39

ATGGCAAGATCCCTTCTCCTGCCCTGCAGATCCTACTGCTATCCTTAGCCCTTGAAACTGCAGGAGAAGAAGC
 CCAGGGTGACAAGATTATTGATGGCGCCCCATGTGCAAGAGGCTCCACCCATGGCAGGTGGCCCTGCTCAGT
 GGCAATCAGTCCACTGCGGAGCGTCTGTTCAATGAGCGCTGGTGTCTCACTGCCGCCACTGCAAGATGA
 ATGAGTACACCGTGACCTGGCAGTGATACGCTGGCGCAGGAGAGCTCAGAGGATCAAGGCCCTCGAAGTC
 ATCCGCCACCCCGCTACTCCACACAGACCCATGTTAATGACCTCATGCTCGTGAAGCTCAATAGCCAGGCCA
 GGCTGTATCCATGTTGAAGAAAGTCAGGCTGCCCTCCCGCTCGAACCCTTGAACCCCTGTACTGTCTCC
 GGCTGGGCACTACACGAGCCAGATGTGACCTTCCCTCTGACCTCATGTGCGTGGATGTCAAGCTCATCTC
 CCCCCAGGACTGCACGAAGGTTTACAAGGACTTACTGGAAATTCATGTGTGCGCTGGCATCCCCGACTCCA
 AGAAAACGCCCTGCAATGTGTACTCAGGGGACCGTTGTTGTCAGAGGTACCCCTGCAAGGTCTGGTGTCTG
 GGAACTTTCCTTGGGCCAACCCCAATGACCCAGGAGTCTACACTCAAGTGTGCAAGTTCACCAAGTGGATAA
 ATGACACCATGAAAAAGCATCGCTAA

FIG. 1KK

>SGPR530_1_SEQID_40
GTCTCCACAGTGTGTGGGAAGCCTAAGTGTGGGAAGATCTATGGTGGCGGACGCAGCAGCTGGCCAGT
GGCCATGGCAGGCCAGCCTGCTCTACTGGGGCTCGCACCTCTGTGGAGCTGTCTCATCGACTCCTGCTGGCT
GGTATCAACTACCCACTGCTTTCTCAACAAATCCAGGCCCCGAAGAACTATCAGGTTCTGTGGGAACATCCA
ACTGTATCATCAAAACCCAGCACACCCAGAAGATGTCTGTGCACCGGATCATCACCCATCCAGACTTTGAGAAGCT
CCACCCCTTTGGAGTGACATTGCCATGTTGCAGCTGCACCTGCCCTATGAACCTTCACTTCTACATTGTCCTGT
CTGCTCCCATCCCGGACATGCAGCTGCCCAGTAACGTCTCTGTGGATAACCGGCTGGGGAATGCTCACC
GAAGACCATAAGAGGGTGCAACTGTCAACACCTTCTATCTCCAGGAGGCAAGGTGGGCTCATTTGAGAACAC
ACTCTGTATACCTTATATGGCAAGAACTGCAAGGCGAGACCTAAGCTTTGCACGAGGAGATGCTGTGTGG
GGGGTACTTCTCGACAGGAAGTCCATCTGCAAGGCGATTCTGGGGGCTCTAGTCTGCTACCTCCCCAGT
GCCTGGTCTGCTGGGCTGCCAGCTGGGCGCTGGACTGCCGCACTCTGCCCTACCCAGCATCTTCACCA
GGTCACTACTTCACTCAACTGGATTGACGAAATCATGAGGCTCACTCCTCTTTCTGACCCCCGCGCTGGCTCCTC
AC

>SGPR520_SEQID_41
ATGCTGCTGGCTGTGCTGTCTGTCTACCCCTCCCAAGCTCATGTTGCCACGGGGACCCACTGTACACAG
CCTGCCCCCAGCGCCCTGCAAGTCTTCACTCTCTCTTGGGGCAGAGACTGTGTTGGCCGAACCTAGACT
ACGTTTGTGAAGGGCGTGGCGGAGAGGGCTCCGAGCACTGCCAATGTACGCGGGGCCACGGCCGCATCG
TGGGGGCGAGCGCGCGCCCGGGCTGGCCCTGGCTGGTGAAGCTGCACTCGGCGGCGAGCCTCTG
TGGCGGCGCTCTGTAAGCGGCTCTGGTGTCTACGCGCAGCGCACTGCTTTGTAGGCTGCCGCTCGACC
CGCAGCGCCCGAATGAGCTTCTGTGACTGTACGCTGGCAGAGGGTCCCGGGGAGCAAGCGGAGGAG
GTGCCAGTGAACCGCATCTGCCCAACCCCAAGTTTGACCCGCGGACCTTCCACAACGACCTGGCCCTGGTGC
AGCTGTGACGCGCGGTAGCCCGGGGGATCGGCGCGCCCGTGTGCTGCCCAAGAGCCCAAGAGCCC
CCTGCCGGAACCGCTGCGCCATCGCGGCTGGGGCTGGGGCGCCCTTTCGAAGACGGGCTGAGGCTGAAGCAGTG
AGAGAGCCCGTGTCCCTGTCTCAGCACCGACACCTGCCGAAGAGCCCTGGGGCCCCGGCTGCGCCCCAGC
ACCATGCTCTGCGCCGGTACCTGGCGGGGGCTTGAAGTCTGCTGCCAGGTGACTCGGAGGCCCTGACC
TGTTCTGAGCCTGGCCCCCGCTAGAGAGTCTGTTCCGAGTCACTCTCTGGGGGACGGCTGCGGGGAG
CCAGGGAAGCCCGGGTCTACACCGCGTGGCAGTGTTCAGGACTGGCTCCAGGAGCAGATGAGCGCAGCC
TCCTCCAGCCGAGCCAGCTGCAGGGAGCTTCTGGCTGGGACCCCCCAGGAGCTGCAGGCAGACGCC
GCCCCGCTCTGCGCTTCTATGCCCGCTGTGCCCGGGTCCAGGGCGCTGTGCGGCTGCGGCACCCAG
CAGTGCCTGCAGCGCGCGCGATGCGAGCTGCGCTGCGCGCACACGCTGCTGGGCTGCTGGCGGAAAC

FIG. 1LL

GGCAGGAGCTGCTCGGGCCCTGTCGGGACTGCGGGCCCTGGCCCCCGCCCTGGCTCTCCCCGCTCCAGCG
CTCAGGGAGTCTCTGACCCCGCCGGAGCTGCGGCTTCACTCAGGATCGGGCTGCAGGCACTCGGT
TCCCGAAGCGAGCGCGGAGCCGCGGAGAACCGCTGCCCTGGCTGGAGCCCCCTGCAGACAGAAGT
TGGCTGCCCTGCAGGGGCCCATGCCCTGGATCCTGCAGTCCCCCTCGAGCACCTGGCCATGAACCTTTCATGA
GGTCTGGCAGATCTGGGCTCCAAGACACTGACCGGCTTTTCAGAGCCTGGTGGGCGAGGCTTGGGGGG
CCGGCATGTGGCTTCAGCGGCTGTTGGGCTGGAGCCGCGCACACTGGCTCGAGCCTCCCCGGCTGCT
GGTGCAGGCCCTGCAGGCCCTCCGCGTGGCTGCCCTGGCAGAGGGGAGCCCCGAGGACCCCTGGATGATGT
AGGCGAGGGGCCCGGCTGGAGAGGAGGGGACCCACTCAACCTCAGGTACCCCCCGCCAGGCAACC
CTGA

>SGPR455_SEQID_42

ATGAGTCCTGATATTGCACTGCTGTATCTAAACACAAAGTCAAGTTTGGAAATGCTGTTCAGCCAACTCTGCTTC
CTGACAGCGATGATAAAGTTGAACCAGGAATCTTTGCTTATCCAGTGGATGGGCAAGATTTCCAAACATCAG
AATATTCAAATGTCTACAAGAAATGGAACCTCCCATCATGATGACAGAGCGTGAATACTGTCTCAAGAGCAT
GAACCTCCCTCCCTGGAGAGGACCATGCTGTGCTGGCTTCCCTGATTGGGAATGGACGCCCTGCCAGGGG
GACTCTGGAGGACCACTGGTTTGTAGAAGAGGTGGTGGAACTCTGGATCTTCTGGGATAACTTCTCTGGGTAGC
TGTTGTGCTGGAGTTTCAGTTCCCGTAAGAAACAACCATGTGAAGGCATCACTTGGCAATTTCTCCAAAGTGTC
TGAGTTGATGATTTATCACTCAAACCTGTTACAGGTTTGGATCGGGCCACCCCTCTCAAAAGTGGGCTC
AAGGTATATAACAAAGGCCCTGAGTTCTGTCCAAGAAGTGAATGGAAGCCAGAGAGATAAATAATCTGTATAA
ATTTACAAGTTTAGACATGGAAAAGCAAGTTGGATGTGATCATGACTATGTATCTTTACGATCAAGCAGTGGAGTG
CTTTTAGTAAGTCTGTGGAATAATTTGCCCTTACCATTGCTGGCAGAGACCAGTGAGGCCATGGTTCCATT
GTTTCTGATACAGAGACAGTGGCAGTGGCTTTGAGCTTACCGTTACTGCTGTACAGAAGTCAGAAGCAGGGTC
AGGTTGTGGAGCTGGCTATATTGGTAGAAGAAGGGACAAATCACTCTGCCAAGTATCCTGATTTGTATCCCCAG
TAACACAAGGTGTCAATTGGTTTCAATTGCTCCAGAGAAGCACATTATAAGTTGACATTTGAGGACTTTGCTGTC
AAATTTAGTCCAACTGTATTATGATGCTGTTGTGATTACGGTGATTCTGAAGAAAAGCACAAAGTTAGCTAACT
TTGTGGAATGTTGACCATCACTCAATTCAGTTCTAGTAACATGACGGTGATCTGAAGAAAAGCACAAAGTTAGCTAACT
ATCGTTTACAAGGCTTCAAGGCCAGATTTACCAATTTGCCCTCAGAGTCTTTAAACAAATTTGAACCAAGTTACC
TCCCCAAACAATCCTGTATCTACCGTAAAGCTATTCTGCATGATGTCTGTGGCATCCCTCCATTAGTCCCCAG
TGGCTTCCAGAAGATCGCAGGAGGGGAAGCCCTGCCCCACTGTGGCCATGGCAGGTGGGTCTGAGGT
TTCTAGGCGATTACCAATGTGGAGGTGCCATCATCAACCCAGTGTGATTCTGACCCGACGCCCACTGTGTGCAA
TTGAAGAATAATCCACTCTCCTGGACTATTATTGCTGGGACCATGACAGAAACCTGAAGGAATCAACAGAGCAG
GTGAGAAGGGCCCAACACATAATAGTGCATGAAGACTTTAACACACTAAGTTATGACTCTGACATTTGCCCTAATAC

FIG. 1MM

AACTAAGCTCTCTGGAGTAACTCGGTGGTGAGGCCAGTATGTCTCCACACAGCGCAGCCTCTATTTT
 CCTCGGAGATCTGTCTGTGACCGGATGGGAAGCATCAGTGCAGAGCTCTCTCTGAATGTTTCTTATTAGATG
 GTGGCTAGCAAGTCGCCTACAGCAGATTCAGTGCATGTGTTAGAAAGAGAGGCTGTGAACACACACTTACTATT
 CTGCCCATCCAGGAGGATCACAGAGAAGATGATCTGTGCTGGCTTTCAGCATCTGGAGAGAAAGATTTCTGC
 CAGGGAGACTCTGTGGCCACTAGTATGTAGACATGAATAATGGTCCCTTGTCTCTATGGCAATTGTCAGCTG
 GGGAGCTGGCTGTGCCAGCCATGGAAGCCGGGTGATTTGCCAGAGTGATGATCTTCTGGACTGGATCCAAT
 CAAAAATCAATGTTAAATTGTTTCAATGTTATTAAACAATAACCTCTTCTTTAGAGTGGGTTTGGAACAGTG
 AGTTGTTGCTCTGAAGCAGAGCTAGAAAAGCCTAGAGGCTTTTCCACACACCGGTATCTACTGGATTATAGA
 GGAAGACTGGAATGTTCTGGGTGCTCAGAGTTTCAGCAAGCAGTATGGCAAAATTTACCATTTGAGTATCTGTCA
 CTCCTGGGGTCTCTGTGTCAAGACTCAGTTCTAATTTATGAAGAAAGACACAGTAAAGAAAAGACGGCA
 GGTGGATTACATGGAAGAGACTTTACTCAATGACTTTTCATGAGTCTGGACCGCTGGTGAGGGTGACATTTCCAT
 GCCCTTGACGAGGTGCAATTTGGTATAAGCTATAATTGCTTGAAGTCTTAGGTCCAAAGGACAGTAAATAACC
 AGACTTTCCTCCAAAGTTCAACACAGAGAGCAGTGGTCCCTGTGAGGATGTTCTTCTGACCAAGCCAGAGGATC
 ATGCAGATCCCAAGAAATTCACAGAACTACTATGGTTGCCAATGGAGATTAGTAGCCCCCTTTAAATCACATCA
 TTCAGCTTAATATTAACTTCCCGATGAAGCCCAACACTTTTGTCTGTCTATGGTCTGCTGTTTACGAAGG
 ATTTGGACCAGGAAAAAATTAATAGTAGAATGTTGATGAGCAGTCTGAGCTTTCTTGGTTCTTAAGCCAAATTCAGC
 ACCAAGAAAGCACAGCTTCTTGTGGGGAGACTGCAGTATCTATGAAAATGATGTATACCTTCTATCTTTCTTGCCC
 TACAGAACACCTGTTACCATGCACCTGCCTCATGAGGTGTTTGGAGAAATTAATAA

>SGPR507_2_SEQID_43

ATGAAATATGTCTTCTATTTGGGTGCTCGCTGGGACATTTTCTTGTGACTCATCTGTTCAGAAAGAAGACC
 CTGCTCCCTATTTGGGTACCTCAAGTCTCACTTCAACCCCTGTGTGGCGTCTCTCATCAACCCAGCTGGGTGC
 TGGCCCCAGCTCACTGCTATTTACCAATCTGAAGTGTGCTGGGAAATTTCAAGAGCAGAGTCAGAGACGGTA
 CTGAACAGACAAATTAACCCCATTCAGATCGTCCGCTACTGGAACACAGTACATAGCGCCCCACAGGATGACCTCA
 TGCTCATCAAGCTGGCTAAGCCTGCCATGCTCAATCCCAAGTCCAGCCCCCTTACCCTCGCCACCACCAATGTCA
 GGCCAGGCACCTGTCTACTCTCAGGTTTGGACTGGAGCCCAAGAAACAGTGGGCTTTGGCAGCTGGAGCCA
 CCAGGCCATCTGACTCTGCACAGAGGCCAGCCATTCTGATTGGCAGAGACACAAATTCACATGAACAAGGCCG
 ACACCTGACTTGGGCAGAACCTGGAGGCCCCCGTGATGCTGATCGAGAATGCCAAAAACAGAACAAAGGAA
 AAAGCCACAGGAATTCCTTATGTGTGAAATTTGTGAAAGTATTCAGCCGAATTTTGGGAGGTGGCCGTTGCTA
 CTGTCACTGTCAAAGACAAAGTCCAGGGAATCGAGGTGGGCACTTTCATGGGAGGGGACGTCGGCATCTACAC
 CAATGTTACAAATATGTATCCTGTGATTGAGAACACTGCTAAGGACAAGTGA

FIG. 1NN

>SGPR559_SEQID_44
ATGGGGGAAAATGATCCGCCTGCTGTTGAAGCCCCCTTCTCATCCGATCGCTTTTGGCCTTGATGATTTGAAA
ATAAGTCCTGTTGCACACAGATGCAGATGCTGTTGCTGCACAGATCCTGTCACTGCTGCCATTGAAGTTTTTCCCA
ATCATCGTCATTGGGATCATTTGATATAGCACTGGCCATTGGCTGGGCATCCACTTCGACTGCTCAGGG
AAGTACAGATGTCGCTCATCTTTAAGTGTATCGAGCTGATAGCTGATGTGACGGAGTCTCGGATTGCAAGAC
GGGAGGACGAGTACCGCTGTGTCGGGTGGTGAGATGCCGTGCTCCAGGTGTTACAGCTGCTTCGT
GGAAGACCATGTGCTCCGATGACTGGAAGGTCACACGCAATGTTGCCCTGTGCCAACTGGGTTTCCCAAGC
TATGTGAGTTCAGATAACCTCAGAGTGAGCTCGCTGGAGGGCAGTTCGGGAGGAGTTTGTGTCATCGATCA
CCTCTTGCCAGATGACAAAGGTGACTGCATTACACCACTCAGTATATGTGAGGGAGGATGTGCCTCTGGCCACG
TGGTTACCTTGCACTGACAGCCTGTGGTATAGAAAGGGCTACAGCTACGCATCGTGGTGGAACATGTCC
TTGCTCTCGCAGTGCCCTGGCAGGCCAGCCTTCAAGTCCAGGGCTACACCTGTGCGGGGCTCTGTCTCA
CGCCCTGTGGATCATCACTGCTGCACACTGTGTTATGACTTGTAACCTCCCAAGTCATGGACCATCCAGGTGG
GTCTAGTTTCCCTGTTGGACAATCCAGCCCCATCCCACTTGGTGGAGAAGATTGTCTACCACAGCAAGTACAAGC
CAAAGAGGCTGGGCAATGACATCGCCCTTATGAAGCTGCGCGGCCACTCACGTTCAATGAAATGATCCAGCCT
GTGTGCCCTGCCCAACTCTGAAGAGAACTTCCCGATGGAAAGTGTGCTGACGTCAAGATGGGGGCCACAG
AGGATGGAGCAGGTGACGCCCTCCCTGTCTGAACCAACGCGGCCGTCCTTTGATTTCCAAAGATCTGCAAC
CACAGGGACGTGTACGGTGGCATCATCTCCCTCCATGCTCTGCGCGGCTACCTGACGGGTGCGTGGACA
GCTGCCAGGGGACAGCGGGGCGCCCTGGTGTCAAGAGAGGAGGCTGTGGAAGTTAGTGGAGCGGACCA
GCTTTGGCATCGGCTGCGCAGAGGTGAACAAGCCTGGGGTGTACACCCGCTGTACACCTCCTTCTCTGGACTGGAT
CCACGAGCAGATGGAGAGAGACCTAAAAACCTGA

>SGPR567_1_SEQID_45
ATGGAGAGGGACAGCCACGGGAATGCATCTCCAGCAAGAACACCTTCAGCTGGAGCATCTCCAGCCCGAGGCAT
CTCCAGCTGGGACACCTCCAGGCCGGGCATCTCCAGCCAGGCATCTCCAGCCAGGCATCTCCAGCTGGGAC
ACCTCCGGGCCGGCATCTCCAGCCAGGCATCTCCAGCTGTACACCTCCAGCCGGGCATCTCCAGGCCG
GGCATCTCCAGCCAGGCATCTCCAGCCAGGCATCTCCAGCCAGGCATCTCCAGCCGGGCATCTCCGGCT
CTGGCATCATTCCAGGTCTCATCCGGCAGGTATCATCCGCCAGGTCAAGCTCGGTGACAACTCCCAAC
CAGAGTGTACCTGTTAGAGCAACACCAAGTGGGGTGTACCCATCCGATCATCTCTGCCAGGTCAAGCACCAG
CAACCAGGCCACAGGGAGAGCCCAAGTCCAGTTCTGGCAGGGCCACACAGGATCAGGTACAAGGAGCAGA
GGGAGAGCTGTCCCAAGCACGCTGTTGCTGTGACGGGGTGGTGGACTGCAAGCTGAAGAGTGACGAGCTGG
GCTGCGTGAGGTTGACTGGGACAGTCTGTCTTAAATCTACTCTGGTCTCCCATCAGTGGCTTCCCATCT
GTAGCAACAACTGGAATGACTCTACTCAGAGAAGACCTGCCAGCAGCTGGGTTTCGAGAGTGTCTCACC

FIG. 100

ACCGAGTTGCCACAGGATTTGCCAACAGCTTCTCAATCTTGAGATACAACCTCCACCATCCAGGAAAGCCTC
 CACAGGTCTGAATGCCCTTCCAGCGGTATATCTCCCTCCAGTGTCCCACTGCGGACTGAGGGCCATGACCGG
 GCGGATCGTGGAGGGCGCTGGCTCGGATAGCAAGTGGCTTGGCAAGTGAGTCTGCACCTTCGGCACCCAC
 CCACATCTGTGAGGCACGCTCATTGACGCCCAGTGGTGCTCACTGCCGCCCACTGCTTCTTCTGACCCCGG
 GAGAAAGTCTGGAGGGCTGGAAGGTGTACCGGGCACAGCAACCTGCACCAAGTTGCCGTAGGCAAGCCTCC
 ATTGCCGAGATCATCAACAGCAATTACACCGATGAGGAGGACGACTATGACATCGCCCTCATCGGCTGTC
 CAAGCCCTGACCCCTGTCGCTCACATCCACCCTGCTTGCCTCCCATGATGGACAGACCTTTAGCCTCAATG
 AGACCTGCTGGATCACAGGCTTTGGCAAGACCAGGGAGACAGATGACAAGACATCCCTTCCCTCCGGAGGT
 GCAGGTCAATCTCATCGACTTCAAGAAATGCAATGACTACTTGGTCTATGACAGTTACCTTACCCCAAGGATGAT
 GTGTGCTGGGACCTTCGTGGGGCAGAGACTCCTGCCAGGGAGACAGCGGGGGCCTCTTGTCTGTGAGCA
 GAACAACCGCTGGTACCTGGCAGGTGTACCAAGCTGGGGCACAGGCTGTGCCAGAGAAACAAACCTGGTGTG
 TACACCAAGTGACAGAAGTTCTCCCTGGATTTACAGCAAGATGGAGAGCGAGGTGCGATTTCAGAAATCCTAA

>SGPR479_1_SEQID_46

ATGGCTGCCCCCTGCTTCCGTTCATGGGGCCCACTCGGGCCCTCTGCCCTGGCCCTTCTGCTGCTGCTGCTG
 TGGCCCTCCCGGTGCGAGCATTTGGTCCACAGACAGCCAGAGAACCAAGGAATCTCCCTAACTGGCAGCGT
 GGCCTGTGGTGGCCCAAGCATGAGGGGAAATCCTGGCGCGTCCCTGCCCGGAGAGGAAGTGGCCGTG
 GCAGGTACGCTGCACTACGCGAGGCTCCACGCTGCGGGGCTCCATCCTCAATGAGTACTGGTGTGTCA
 GCTGCGCACTGCTTACAGGGACAGAAATACAAATCTATGACATGTACGTAGGCCCTCGTAAACCTCAGGGTG
 GCCGGCAACCAACCCAGTGGTATGAGGTGACAGGGTGATCCTGCACCCCAATATGAGATGTACCAACCCCAT
 CGGAGGTGACGTGGCCCTGGTGCAGCTGAAGACCCGCAATGTGTTTCTGAGTCCGTGCTCCCGTTTGCCTTG
 CAACTCCAGAAAGTGAACCTTACAGTGCCAAATTGCTGGGCTACGGGATGGGACTAGTCTCAAACAAGGTGAG
 ACCTCAGACGAGCTGCAGGAGGTGCAGCTCCCGCTGATCCTGGAGCCCTGGTGCCACCTGCTCTACGGACACA
 TGTCTACATCATGCCCGACATGCTGTGTGCTGGGACATCCTGAATGCTAAGACCGTGTGTAGGGCGACTCC
 GGGGGCCCACTTGTCTGTGAATTC AACCGCAGCTGGTTGCAGATTGGAATTGTGAGCTGGGGCCGAGGCTGCT
 CCAACCTCTGTACCCCTGGAGTGTATGCCAGTGTTCCTATTTCTCAAAATGGATATGTGATAACATAGAAATCAC
 GCCCACTCCTGCTCAGCCAGCCCTGCTCTCTCTCCAGCTCTGGGGCCCACTCTCAGCGTCTAATGGCCATGC
 TGGCTGGCTGCTCAGTGTGTGA

>SGPR489_1_SEQID_47

ATGAGTCTCAAAATGCTTATAAGCAGGAACAAGCTGATTTTACTAGGAATAGTCTTTTTTGAACGAGGTAAT
 CTGCAACTCTTTTCGCTCCCAAGCTCCAGTGTGGGCAGAGTCTGTTAAGGTACAGCCTTGAATATTATTTA

FIG. 1PP

ACATTTTCAGTCGCATTCTTGGAGGAAGCCAAAGTGAGAGGGTTCCCTATCCCTGGCAGGTATCTCTGAAACAAA
GGCAGAAAGCATATTTGTGGAGGAAGCATCGTCTACACACAGTGGGTGATCACGGCGCTCACTGCAATTGCAAAC
AGAAACATTGTGCTACTTTGAATGTTACTGCTGGAGAGTATGACTTAAGCCAGACAGACCCAGGAGAGCAAACT
CTCACTATTGAAACTGTATCATATACATCCACATTTCTCCACCAAGAAACCAATGGACTATGATATTGCCCTTTTGAA
GATGGCTGGAGCCTTCCAAATTTGGCCACTTTGTGGGCCCATATGTCTCCAGAGCTGCGGGAGCAATTTGAGG
CTGGTTTTATTGTACAACTGCAGGCTGGGGCCGCTTAAGTGAAGGTGGCGTCTCTCACAACTTTGCAAGGAA
GTGAATCTGCCTATTTGACCTGGGAAGAGTGTGTGGCAGCTCTGTTAACACTAAAGAGGCCCATCAGTGGGAA
GACCTTTCTTTGCACAGGTTTTCTGATGGAGGAGAGACGCGATGTCAGGGAGATTCAGGAGGTTCACTCATGT
GCCGGAATAAGAAAGGGCCTGGGACTCTGGCTGGTCAATTTGGGAGGCTCAGGTGGGAGGATCGCTTGAGTC
CAGGAGTTCAAGACCAAGCCTAGGCAACAAGTGAGACTCTGTCTACAAATAATTTCTTCAAAAATTAGCCCG
GTGTGGCACCTGGTGAGTGAGCAGGATGTCAATAGTACGGGGCTGAGGGAAAGCTGCACITCCAGAAAGC
CTCCACCTATATTATGAGAGCAAGCAACGGTGTCTGGACCCCTGCTGGTACCAGAGGAAATGCATGTGTTGCT
CAGTTTTCCACCTAGATGTTGAGTCTTGTCAACACAGTTACCTGTCAATGTATTTTAGAAGACAGACCCATT
GGAAATTTTGTGGAGAAAGCCTCCCTCATCCATTTCTATTGGCTCTAATCTCTAAGGCTGAAATTCGTCTCTG
ATGCCACAGATTATGCAGCTGGGTTTAACTTACCCTATAAGCTCTTAACCAAACTACATTCCTGTTGCAGTTA
CTTAACCTGTCCTTTTGAAGAAGGTCTCATAACAGAGTCTAACTATCCTGAAACTACAGTGACAAAGGCTAACTGT
GACTGGATTTTCAAGCCTCCAAACATCACCTAATTAAGCTTTTCATTCAGAGTCTGAAATAGAAAGAAAGTGGAG
ACTGCACCTCCGACTATGTGACAGTGCACAGCGATGTAGAAAGGAAGAAATAGCTCGGCTGTGTGGCTAT
GATGTCCCCACCCCTGTGCTGAGCCCCCTCCAGCATCATGCTCATCAGCTTCCATTCCAGATGAAACGGGACCTG
CAGGGGCTTTTCAGGCTATAGTCTCCTTCTTCTAAAGCAGTATACCCAGATTTAAACATCTCCATATCAGAGGAT
GAGTCAATGTTTCTGGAGACATGA

>SGPR465_1_SEQID_48

CGGTGGCCATGGCAGGCCAGTCTCCTCTACCTAGCGGGGCACATCTGTGGAGCTGCCCTCATCGACAGCAACT
GGGTGGCCTCTGCTCACTGCTTCCAAAGATGCATCTTCCCTCCACGGGCCCGCTGCCACTAACCCATCT
GATTACCGGATCCTGCTTGGGTATGACCAGCAAGCCATCCCACAGAGCACAGCAAGCAGATGACAGTGAATAA
GATCATGGTGCACGCTGACTATAACGAGTTGCACCGCATGGGAGTGACATCACCCCTGCTGCAGCTGCACCATC
ATGTGGAATTCAGCTCCACATCTCCCGCTGCCCTCCGGAACCAACCAAGTGGCTGGCCCTGACAGCTCC
TGCTGGATATCTGGTTGGGAATGTCACCGAGGATGCTTCTGCTGCCCTGAGCCCTTCCAACTTCAGGAGGCAGA
GGTCGGTGTGATGGACAACACTGTCTGCGGATCCTTTTCCAGCCCCAGTACCCCGCCAGCCAGCAAGCAGCAGTG
ACTACACCATCCACGAGGACATGCTGTGCGCTGGGACCTCATACAGGAAAGGCCATTTGCCGAGTGAATCC
AGGGTCCCCTCGTCTGCCCATTAATGGCACCTGGTTCTCTGATGGGGCTGTCTAGTTGGAGCCTCGACTGCTG

FIG. 1QQ

CTACCCGTCGGTCCAGGGTCTTACCAGGCTCCCTACTTACCACCTGGATCAGCCAGAAGAAGAGGGAGA
 GCACCCCTCCAGATCCCGCTTGGCTCCTCCTCAGGAACACCCAGCCCTGGACAGCATGACCTCTCAGGG
 CATCGTCCACAAGCCCGGCTCTGCGCAGCCCTTCTGGCTGCTCACATGTTCCCTCCTGCTGCTGATTCCTCTGG
 GGAGCCTG

>SGPR524_1_SEQID_49
 ATGGACAAAGAAACAGCGATGTTTCAGCCGACCTGCTGACCTGAAAATATCCAATATCTCAGTCCAAGTGGTC
 AGTGCCCAAAAGAGCTGCCAGTGAGACGACACCCGTTGCCAGGGAGACGACTACCATTGCCAGGAAGACGAC
 CACCACAAGAGACCCATTGGCAAGGCCAAACCCAGAAAGCAATCCAAAGAAAAGTTCCCTTTTGGAAATGTACAAA
 ATAAATCATCTCTTCACAGTATTTTATTCATCCTAGCAGTCATAGCCTGGACACTTCTGTGGCTGTATATCAGT
 AAACAGAAAGCAAAGATGCTTTTACTTTGCTGGGATGTTTCGCATCACCAACATTGAGTTCTTCCCGAATACC
 GACAAAAGGAGTCCAGGGAATTTCTTTAGTGTACGGACTGTGCAGCAAGTGATAAACCTGGTTTATACAACAT
 CTGCCCTTCTCCAAATTTATGAGCAGTCTGTTGTGCAGATGTCAGCAGCAACAACAAGCGGCTCCTTTGTCC
 ACTTTGGATTGTTTGTATGCCACGTGCCAAGGCCACATCTTCTGTGAAGACTGTGTGCCGCCATCTTGAA
 GGACTCCATCCAGACAAGCATATAAACCGGACCTCTGTGGGAGCTTGCAAGGACTGGCTGTGGACATGGAC
 TCTGTGGTACTAAATGGTGATTGTTGGTCAATTCCTAAAAAAGAAAGAAAGAAATGGTGTCTCCACAG
 ACAAGGCTGCTCAGTACTTCTATGCAGAGCATCTGTCTCCTCACTACCCGCTGGAGATTTCTGAGCCTCAG
 GGAGGCTGATGTGCTCACTTCAAGCTGGTGCCATAGTGGGTACCTGATTGCTCTCAATCAAGTCCATCCAAA
 TCGAAGCCGACAACCTGTGCTCACTGACTCCCTGACCATTACGACTCCCTTTGCCCATCCGGAGCAGCATCTGT
 ACAGAAATTTGTGAACCCACAAGAACATTAAATGTCATTTGTTTCTACAAATAATCTCATGTTGGTGACATTTAAGTCT
 CCTCATATACGGAGGCTCTCAGGAATCCGGGCAATTTTGAGGTCAATCCAGAAACAAAGTGTGAACACACAGTG
 TTGGTCAAAGACATCAGTGGCTTTGAAGGGAAATTTCAAGCCCATATTACCCGAGCTACTATCCTCCAAAATGCA
 AGTGTACCTGGAAATTCAGACTTCTCTATCAACTCTTGGCATAGCAGTGAATCTATAACTATTCAATAACCAAG
 AAGAGTATGAAAGGCTGTGAGCATGGATGGTGGAAATTAATGAGCACATGTAAGTGTGGCTCCTACATGGATCAT
 CAGACAATTTTCGAGTCCCCAGCCCTCTGGTTACATTCAGCTCCAGTCCAGTTCAGGCTTCAGACAAAGCCA
 CTTTGGCAGAAATATGGCAGTTACAACATCAGTCAACCCCTGCCCTGTTGGATCTTTTAGATGCTCCTCCGGTTTAT
 GTGTCCCTCAGGCCCAGCGTTGTGATGGAGTAAATGACTGCTTTGATGAAGTGTGAAGTGTGCTGAGCC
 CTCAACCTGCCTGCAATACCAAGTCCCTTCAGGCAGCATGGCCCTCTCATCTGTGATGGCTTCAGGGACTGTGAG
 AATGGCCGGGATGAGCAAAACCTGCACCTCAAGTATTCATGCAACAACAGAACTTTTAAGTGTGGCAATGATATT
 TGCTTTAGGAACAAAATGCAAAATGTGATGGACAGTGGATTGTCCAGATGGAAGTGTGAAGAGGCTGCAC
 CTGCAGCAGGAGTTCTCCGCCCTTCAACCGCATCATCGGAGGCACAGACACCCCTGGAGGGGTTGGCCGTGG
 CAGGTACGCCCTCCACTTTGTTGGATCTGCCTACTGTGGTGCCTCAGTCACTCTCCAGGGAGTGGCTTCTTCTGCA

FIG. 1RR

GCCCACTGTTTTTCATGGAAACAGGCTGTGAGATCCACACCATGGACTGCACACCTCGGGATGTATGTTCAGGG
 GAATGCCAAGTTTGTCTCCCGGTGAGAAGAAATTGTGGTCCACGAGTACTATAACAGTCAGACTTTTGATTATGA
 TATTGCTTTGCTACAGCTCAGTATTGCCCTGGCCTGAGACCCCTGAACAGCTCAATTCAGCCAATATGCATTCCTCC
 CACTGGTCAGAGAGTTCGACGTGGGAGAAAGTCTGGTAACCTGGCTGGGGCGAAGACACGACGAGAGATAAT
 AAAGGCTCCCTCGTTCTGCAGCAAGCGGAGGTAGAGCTCATTGATCAACGCTCTGTGTTCCACCTACGGGAT
 CATCACTTCTCGGATGCTGTGCAGGCATAATGTGAGCAAGAGAGATGCCTGCAAGGAGATTTCGGGTGGAC
 CTTTATCTTGTGGAAGAAAGTGTGAAATGGATTTTGAAGCTGGGATTTGCTGCAAGGACATGGATGTGGAC
 GACCAAACTTTTCTGGTGTACACAAGGGGTGTCAAACCTTTGTTCCCTGGATTCAATAATATGTCCTTCTCTTTT
 GTAA

>SGPR422_SEQID_50

ATGACATTGAACAAATTAAGACCTTTTTGCAGGGAAAGACAGTGGGATTTGGCACCCGAAAGCAGAAATGCTG
 AAGCCATGGATGATTGCCGTTCTCATTTGTGTTGCCCTGACAGTGGTGGCAGTGACCATAGGCTCTCCTGGTTAC
 TTCCTAGTATTGACCACAAAAGGAGTACTATCATGGCTCCTTTAAATTTTAGATCCACAAATCAATAACAATTT
 CGGACAAAGCAACACATATCAACTTAAGGACTTACGAGAGACGACCCGAAATTTGGTGTATTCTTTGAAAATGTAC
 CTTTCTTTTGTGTACAGTCCAGAGGAAGATGGTGTGAAAGTAGATGTCATTATGGTGTCCAGTTCCCTCTA
 CTGAACAAAGGGCAGTAAGAGAGAGAAAGAAATCCAAAGCATCTTAATCAGAAGATAAGGAATTTAAGAGCCTTGC
 CAATAAATGCCCTCATCAGTTCAGTTCAAGTTAATGTGGCCATGGTCAAGAAATGGCAATGTGGGGCCAGGTTCCGGAGCA
 GGAGAGGCTCCAGGCTGGAGGAGGAGTCCCTGGTCCACCAATGAGCTCATCAACAGGGGAGTTAACTGTCC
 AAGCAAGTTGTGGTAACGAGTTGTCCATTAAACGTCAACAGAAATAGCATCTGGAGTCAATGCACCCAGGCGG
 CCTGGCCTTGGCAAGCTTCCCTTCAGTATGATAACATCCATCAGTGTGGGGCCACCTTGATTAGTAACACATGGC
 TTGTCACTGCAGCACACTGCTTCCAGAAGTATAAAATCCACATCAATGGACTGTAGTTTGGAAACAAAATCAA
 CCTCCCTTAATGAAAGAAATGTCAGAAGATTTATATCCATGAGAAGTACCCTCTGCAGCAAGAGAGTACGA
 CATTGCTGTTGTCAGGTCCTTCCAGAGTCACCTTTTCGGATGACATACGCCAGATTTGTTGCCAGAAAGCCTC
 TGCACTCCTTCCAAACCAATTTGACTGTCCACATCAGAGATTGGAGCACTTTACTATGGTGGGAATCCCAAAT
 GATCTCCGAGAAGCCAGAGTGAAATCATAAAGTACGATGTCTGCAAGCAACCCACAGGTGTATGGCAATGATATA
 AAACCTGGAATGTTCTGTGCCGATATATGGAAGGAATTTATGATGCCTGCAGGGGTGATCTGGGGACCTTTA
 GTCACAAGGGATCTGAAAGATACGTGGTATCTCATTTGGAATTTGAAGCTGGGAGATAACTGTGTCAAAGGAC
 AAGCCTGGAGTCTACACACAAGTGACTTATACCGAAACTGGATTGCTTCAAAAACAGGCATCTAA

FIG. 158

>SGPR538_SEQID_51
ATGAGCCTGATGCTGGATGACCAACCCCTATGGAGCCCAAGTATGCAGAGGAGGCCAGGACCTGGGATCT
TCAGAGCAGAGCCTGGAGACCAGCAGCATCCCATTTCTCAGGCGGTGTGCTGGCGTTCCATGCGACGTGGCTG
TGCACTGCTGGAGCCCTGGGCTGCTGGCCGTCAGGTGTGGCTCATGGCTCCTAGTGTATCTGTGT
CCTGCTGCCCTCAGCCCCATTTCCGGGACCTTGAGGATGAGGAGATAACTTTGAGCTGCTCAGAGGCCAGCGC
TGAGGAAGCTCTGCTCCCTGCACTCCCAAAACAGTATCTTTCAGATAAACAGCGAAGACTTCTTGTGGAAGC
GCAAGTGAGGGATCAGCCACGCTGGCTCCTGGTCTGCCATGAGGGCTGGAGCCCCCGCTGGGCTGCAGAT
CTGCTGAGCCCTGGGCATCTCAGACTCACTCACCACAAGGAGTAACCTCACTGACATCAAACTCAACAGTTC
CCAGGAGTTTGCTCAGCTCTCTCTAGACTGGAGGCTTCTGAGGAGCGGTGGCAGCCAGGAACAACACTGC
ACTTCTGGTCAAGTTGTTCCCTCAGATGCTCTGAGTGTGAGCGAGGCCCTGGCTTCCCGGATAGTTGTGG
GCAGTCTGTGGCTCCTGGCGCTGGCGTGGCAGGCCAGCGTGGCCCTGGGCTTCCGGCACACGTGTGGGG
CTCTGTGCTAGCGCCACGCTGGTGGTACTGCTGCACATTGTATGCACAGTTTCAGGCTGGCCCGCTGTCCA
GCTGGCGGTTATGCGGGCTGGTACCCACAGTCCCGTCAGGCCCAAGGCTCTGGTGGAGAGGA
TTATCCACACCCCTCTACAGTCCCCAGAACTACATGACTACGACGTGCCCTCCTGAGGCTCCAGACCGCTCTC
AACTTCTCAGACACTGTGGCGCTGTGTGCTGCCGCCAAGGAACAGCATTTTCCGAGGGCTCGCGGTGCT
GGTGTCTGGCTGGGCCACACCCACCTAGCCATACTTACAGCTCGGATATGCTCCAGGACACGGTGGTGCC
CTTGTTACGCACTCAGCTCTGCAACAGCTCTTGGCTGTACAGCGGAGCCCTACCCCCGCTGTTGCGCTG
GCTACCTGGACGGAAGGCTGATGATGCCAGGGAGATAGCGGGGCCCTAGTGTGCCAGATGGGGACA
CATGGCGCCTAGTGGGGTGTGCTCAGCTGGGGGCTGCGTGCGCAGAGCCCAATCACCCAGGTGTCTACGCCA
AGGTAGCTGAGTTTCTGGACTGGATCCATGACACTGCTCAGGACTCCCTCCTCTGA

>SGPR527_1_SEQID_52
ATGGCCCGGCACCTGCTCCTCCCTTGTGATGCTTGTATCAGTCCCATCCAGGAGCCCTCCAGGACTCAGC
TCTCAGTCTCTACCCAGGAAGAACCTGAAGATCTGAGTCTGGGCGCCCTGAGCCCTCGGCCGTCATCGTGGG
GGCTCAACCGCGCAGCCGGCACCTGGCTTGGCAAGTGAAGCTGCACCATGGAGGTGGCCACATCTGCGGG
GGCTCCCTCATCGCCCTCCTGGTCCCTCTCCGCTGCTCAGTGTTCATGACGAATGGGACGCTGGAGCCCCG
CGGCCGAGTGTGCTGCTGGCGTGCACCTCCAGGACGGGCCCTGGAGCGGCGCACACCCGCGCA
GTGGCCGCCATCGTGGTCCCGGCCAACTACAGCCAAAGTGGAGCTGGCGCGGACCTGGCCCTGCTGCGCCTG
GCCTACCCGCCAGCTGGGCCCGCCGCTGTGGCTGTCTGCCCTGCCCGCCCTCACACCGCTTCGTGCAC
GGCACCGCCTGCTGGGCCACCGGCTGGGAGACGTCCAGGAGGACATCCTGCTCCTCCCTGGGTGCTA
CAGGAAGTGGAGCTAAGGCTGCTGGCGGAGGCCACCTGTCAATGTCTACAGCCAGCCCGGTCCCTTCAACC

FIG. 1TT

TCACTCTCCAGATATTGCCAGGGATGCTGTGTGCTGCTACCCAGAGGGCCGAGGACACCTGCCAGGGTGA
CTCTGGGGGGCCCTGCTGTGAGGAAGGGCCGCTGTTCCAGGCAGGAATCACCAGCTTTGGCTTTGGC
TGTCAGCGAGAAACCGCCCTGGAGTTTCACTGCTGTGGCTACCTATGAGGCATGGATACGGGAGCAGGTGAT
GGTTTCAGAGCCTGGCCCTGCCCTTCCACCCAGCCCAAGAACCCAGTCAGATCCCCAGAGCCAGGGAG
GAGAACTGCACCAATTGCCCTGCCCTGAGTGGGGAAGCCCGGCCAGGGCCCTGGCCCTGGAGGGCCAG
GTGATGGTGCCAGGATCCAGACCCCTGCCATGGGGCGCTGGTGTCTGAAAGCTGGGTCTTGGACCTGCCAGCT
GCTTTCTGACCCGAACAGCTCCGACAGCCACCCCGACCTCGACGCTGGCGCTGCTGCTGCCCTCGCG
CCGCGCGGAGCGGTGGCGCCTGGTGACGACGAGAACGCTTCGTGGGACAAACGCTCGGACCTGG
CGCTGCTGAGCTGCGCACGCCCGTGAACCTGAGCGCGCTTCGCGCCCGTGTGCTACCCACCCGGAAC
ACTACTTCTGCCCCGGAGCCGCTGCCGCTGGCCCGCTGGGGCCGCGGGGAACCCGCTTGGCCAGGC
GCGCTGCTGGAGGCGGAGCTGTTAGCGGCTGGTGGTCCACTGCTGTACGGCCGCGAGGGCGGCAGTA
CCGCTGCCCGGAGACCCGCGCACGCGCTCTGCCCTGCCCTACCAAGAAAGGAGGAGGTGGCAGCTGCTGG
ACTCATGGCCCATGGATCAGCCATGTGACTCGGGGAGCTACCTGGAGGACCACTAGCTGGGATTGGGGCC
CTGATGGGGAGGAGACTGAGACACAGACTTGTCCCCACACACAGAGCATGCTGCTGTGCCCTGCGGCTGGA
GGCTGCTCCAGTGGGGTCCCTGTGGCCCTGGCTGGCAGAGTGCTGTGGCTGGTATCGAGTCTGCACCTGG
GATCCTCCTGCCCCAGGCTGGTCTTGCGAGCCACTCACTGTCTCAGGCCAGGCTCTACAAACAGTGCCTT
ACATTGAAGTGTATCTGGGCCGGGAGGGCCAGCTCCCTCCACAGGCCACAGGTATCCCGCTTGGTCA
CAGCATCCGCTGCCCGCAGACCTGGGACTCAGGCCCGCCCTGGCCCTCCTGGAGCTGAGCTCCCGGGTGA
GCCCTCCCATCAGCCCTGCCCATCTGTCTCCACCCGGCGGTATCCCGGGGCGCAGCTGCTGGGTGTTG
GGCTGGAAGAACCCAGGACCGAGTCCCTGTGGCTGCTGTCTCCATCTTGACACACAGCATCTGTGACTG
CCTCTATCAGGGCATCCTGCCCGCTGGAAACCTCTGTGCTCTGTATGAGAGGGCAGGAGACAGGTGTGAG
ATGACCTCAGCACCGCCCTCCTGTGCCAGATGACGGAAGGTCTCTGGATCCTCGTGGCATGGCTGTTCAG
GGAGCCGGGAGCTGTTTGTGCTGCTGCTGCTGAGAGGCTGGATCTCCAGACAGTGGGAGAGGCCAAT
CCTGCCCGCCAGTGGCTCCCGACACTGGCCCACTGGAGGAGCAATCTCTGCCCGCCAGAACTGGCCAAAGCC
TCGGGATCCCGCATGCAGTCTACTTCTGCTCCTGCTGACTCTCCTGATCCAGAGCTGA

>SGPR542_SEQID_53

GCCATGGGGCTCGGGTTAGGGGCTGGGGAGCTCCTCTGCTGACTGTGGCCACCGCCCTGATGCTGCCCGTG
AAGCCCCCGCAGGCTCCTGGGGGCCAGATCATCGGGGCCACGAGGTACCCCCCACTCCAGGCCCTAC
ATGGCATCCGTGCGCTTCGGGGGCCAACATCACTCGGAGGCTTCTGCTGCGAGCCCGCTGGGTGCTCGG
CCGCCCCACTGCTTCAGCCACAGAGACCTCCGCACTGGCCTGTGTGCTGGCGCCACGCTCCTGAGTACTGC
GGAGCCACCCAGCAGGTGTTTGGCATCGATGCTCTCACCACACACCCCCGACTACCCACCCCATGACCCACGCC

FIG. 1UU

AACGACATCTGCCTGCTGCAGCTGAACGGCTCTGCTGTCTCTGGGCCCTGCAGTGGGGCTGCTGAGGCTGCCAG
 GGAGAAAGGGCCAGGCCCCCAGCGGGACAGCGGGACACGGTGCCGGTGCTGGCTGGGCTTCGTGCTGACTTTG
 AGAGCTGCCGCTGACTGATGAGGCCAAGTCCGAGTGTGACCCGACGCTGCAACAGCTCCTGGAA
 GGGCCACCTGACACTTACCATGCTCTGCACCCGAGTGGGACAGCCACAGACGGGCTTCTGCTCGGCCGAC
 TCCGGAGGGCCCCCTGGTGTGCAGGAACCGGCTCACGGCTCGTTTCTCTCGGCCCTCTGGTGGCGGAC
 CCCAAGACCCCGACGTGTACACGCAGGTGTCCGCCCTTGTGGCCTGGATCTGGGACGTGGTTCGGCGGAGCA
 GTCCCCAGCCCCGCCCCCTGCCTGGGACCACCAAGGCCCCCAGGAGAGCCGCTGA

>SGPR551_SEQID_54

ATGCCCCGTGGCCGAGGCCCCCAGGTGGCTGGCGGGCAGGGGACGGAGGTGATGGGAGAAAGCGGAGCC
 AGAGGGGATGTTCAAGGCTGTGAGGACTCCAAGAGAAAGCCCGGGGTACTCCGCCCTGGTGGCCCTGTTT
 GTGCTGTGGCCCTGCTGCTGGCTTCGGCGGGGTGCTACTCTGGTATTTCTAGGTACAAGCGGAGG
 TGATGGTCAGCCAGGTACTCAGGCAGTCTGCGTGTACTCAATCGCCACTTCTCCAGGATCTTACCCGCCGG
 GAATCTAGTGCCCTTCGCGAGTGAACCGCCAAAGCCAGAGATGCTAAGGAGCTCATCACCAGCACCCGCT
 GGAACTTACTAACCTCCAGCTCCGTCTATTCTTTGGGAGGGACCCCTCACCTGCTTCTTCTGTTCAATCT
 CCAATCCCCGAGCACCGCCGGCTGATGCTGAGCCCCGAGTGGTGACGCACTGCTGGTGAGGAGCTGCT
 GTCCACAGTCAACAGCTCGGCTGCCCTCCCTACAGGGCCGAGTACGAAGTGACCCCGAGGGCTAGTGATC
 CTGGAAGCCAGTGTAAAGACATAGCTGCATTGAATTCCACGCTGGTGTACCGCTACAGCTACGTACGTGGGCA
 GGGCCAGGTCCTCCGGCTGAAGGGCCCTGACCACCTGGCTCCAGCTGCCCTGTGGCACCTGCAGGGCCCCAA
 GGACCTCATGCTCAAACTCCGGCTGGAGTGGACGCTGGCAGAGTGCAGCCGACCGACTGGCCATGTATGACGTG
 GCCGGCCCCCTGGAGAGAGGCTCATCACCTCGGTGTACGGCTGCAGCCGCGCAGGAGCCCGTGGTGAGGTT
 CTGGCGTCGGGGCCATCATGGCGTCTGTGGAAGAAGGGCCTGCACAGCTACTACGCCCTTCGTGCTCT
 CCGTGACGCCGTGTTCCAGGCCGTGAAGTGAACCTGACGCTGGACAACAGGCTCGACTCCAGGGCGT
 CCTCAGCACCCCGTACTTCCCGAGCTACTCGCCCCAAACCCACTGCTCCTGGCACCTCAGGTGCCCTCTC
 TGACTACGGCTTGGCCCTCTGGTTGATGCCATGCATGAGGAGGCGAGATGATTTGCCGTGCACCCAG
 GGCCAGTGGACGATCCAGAACAGGAGGCTGTGTGGCTTGGCATCTGCAGCCCTACGCCGAGAGGATCCCCG
 TGGTGGCCACGGCCGGATCACCATCAACTTCACTCCAGATCTCCCTCACCGGGCCCGGTGCGGGTGCA
 CTATGGCTGTACAACCAAGTCGGACCCCTGCCCTGGAGAGTTCCTCTGTTCTGTGAATGGAATGACTCTGTGCTCCCTG
 CCTGTGTGGGTCAAGGACTGCCCCAACGGCTGGATGAGAGAACTGCGTTGCAGAGCCACATTCCAGTG
 CAAAGAGGACAGCATGCATCTCACTGCCCAAGTCTGTGATGGCAGCCTGATTGTCTCAACGGCAGCGATG
 AAGAGCAGTGCCAGGAAGGGTGCCATGTGGGACATTCACTTCCAGTGTGAGGACCGGAGCTGCGTGAAGAA
 GCCCAACCCGCGAGTGTATGGCGGCCGACTGCAGGACGGCTCGGATGAGGAGCACTGTGACTGTGGCCT

FIG. 1W

CCAGGGCCCTCCAGCCGCATTGTTGGTGAGCTGTGTCTCCGAGGGTGAGTGGCCATGGCAGGCCAGCCTC
 CAGGTTCCGGGTCGACACATCTGTGGGGGCCCTCATCGCTGACCGCTGGGTGATAACAGCTGCCCACTGCT
 TCCAGGAGGACAGCATGGCTCCACGGTGTGGACCGTGTTCCTGGCAAGGTGTGCAGAACTCGCGGTG
 GCCTGGAGAGGTGTCCTTCAAGGTGAGCCGCTGCTCCTGCACCCGTACACGAAAGGACAGCCATGACTAC
 GACGTGGCGTGTGCAGCTGACACCCGGTGTGGCTCGGCCCGTGCGCCCGTGCCTGCCCGCCG
 CGCTCCCACTTCTCGAGCCCGCTGCACCTGCTGGATTACGGCTGGGGCGCTTGCAGAGGGCGGCCCC
 ATCAGCAACGCTCTGCAGAAAGTGATGTGAGTTGATCCACAGGACCTGTGCAGGAGGCCCTATCGCTACCA
 GGTGACGCCACGCTGTGTGCGCGCTACCGCAAGGCAAGAGGATGCCGTGCAGGGTGAATCAGGTGGT
 CCGCTGGTGTGCAAGGCACCTCAGTGGCCGCTGTTCTTGGCGGGCTGGTCACTGGGGCTGGGTGGC
 CGGCCTAACTACTTCGGCGTCTACACCCGCATCACAGGTGATCAGCTGGATCCAGCAAGTGGTGACCTGA

>SGPR451_SEQID_55

GACCTGCCGCCATCTTGCTCACCAGCCTCCAAATGCGGCTGGGGCTCCTGAGCGTGGCGCTGTTGTTGTGG
 GGAGCTCTCACTTATACTCAGACCACTACTCGCCCTCTGGAAGGCACAGGCTCGGCCCTCGCCGAAACCGG
 GGCTAGTTCAGCAGGCTGAGGCCGTCGCAAGAGGCTCCGGCGGAGGAGGAGGCGCATGCAA
 AGGATTGTGGAACAGCACCGCTTAAGGATGTGTGCAAGGCTCGGATTATAGGGGCAACGAAAGCACAAGCT
 GGCGCATGGCCGTGGTGGTGAGCCTGCAGATTAAATATGGCCGTGTTCTTGTTCATGTATGGGGGAACCCCT
 AGTGAGAGAGAGGTGGTCCCTCAGAGCTGCCCAGTGCCTAAAGACACTAGCGATCCTTAATGTGGACAGCTG
 TGATTGGAACCTAATAATACATGAGCGCTATCCTCATACCAAGAGATAAAATTAAAGCAATCATATTATCATCCA
 AACTTCATTTTGGAACTTATGTAAATGATATTGCACCTTTTCACTTAAAAAAGCAGTGAGGTATAATGACTATATT
 CAGCCTATTTGCCCTACCTTTTGATGTTTCCAAATCCTGGACGGAACACAAAGTGTTTTATAAGTGGCTGGGAA
 GAACAAAGAGAGAGGTAAACGCTACAAATATTTTACAAGATGCAGAGTGCATTATATTCTCGAGAGATGTGTAA
 TTCTGAGAGGAGTTATGGGGGAATAATTCTAACACTTCATTTGTGCAGGTGATGAAGATGGAGCTTTTGATACT
 TGCAGGGTGACAGTGGGGACCAATTAATGTGCTACTTACCAGAAATATAAAGATTTTTGTAAATGGGAATTACC
 AGTTACGGACATGGCTGTGGTCGAAGAGGTTTCTCGTGTCTATATTGGGCCATCCTTCTACCAAAAGTGGCTG
 ACAGAGCATTTCTTCCATGCAAGCACTCAAGGCATACTTACTATAAATATTTTACGTGGCCAGATCCTCATAGCTT
 TATGTTTTGTCATCTTACTAGCAACAATAA

>SGPR452_1_SEQID_56

AGCCCCCGCAGCCAGGACCCCTGACTGTAGGCTCCAGGCCCTCCCTGGAAGCCCTGGCCACGCTCGCCCCG
 CAGCCCTCAGACTGGCTGTGCTTCGCGGATCTTGGCTGTTTCAGGCTGATGGAGCTGCCCACTCCATGGGCC
 TGGGCAGCAGCTTGAAGTGGCGTGGGCCAAGCCCTCTGGGATGCCCGTCCAGAGATGACCTGGTGGGCAT

FIG. 1WW

TGTGGGGCCACAATGCCCCCGGGGAAGTGCCGTGGCAGGTCAGCCTGAGGGTCTACAGCTACCACTG
GGCCTCCTGGCGCACATCTGTGGGGCTCCCTCATCCACCCAGTGGTGCTGACTGCTGCCCACTGCATT
TTCTGGAAGGACACCGACCCGTCCATCTACCGATCCACGCTGGGACGTGATCTCTACGGGGCCGGGGC
TGCTGAACGTGAGCGGATCATGTCCACCCCAACTATGTCACTGCGGGCTGGTGCGGATGTGGCCCTGCT
CCAGTGCCTGGGTACCTCTCTCCAGAGTCGTGCCGCCCTACCGCTGCAGCAGGCGAGTGTGCA
GGTCTGGAGAACGCCGTCTGTGAGCAGCCCTACCGCAACGCTCAGGGCACACTGGCGACCGGAGCTCATC
CTGATGACATGCTGTGCCGCGAGCGAGGCGGAGACTCCTGCTACGGTACTCCGGCGGCCCTCTGGTCT
GCAGGCTGCGGGGTCCTGGCGCCTGGTGGGGTGGTACGCTGGGCTACGGCTGTACCCCTGCGGGACTTTC
CCGGCGTCTACACCCACGTCCAGATCTACGTGCTCTGGATCCTGCAGCAAGTCGGGAGTTGCCCTGA

>SGPR504_SEQID_57

ATCGGGGGCCACGAGGTGACCCCCCACTCCAGGCCCTACATGGCATCCGTGCGCTTCGGGGGCCAACATCACT
GCGGAGGCTTCCTGCTGCGAGCCCGCTGGGTGGTCTCGGCCGCCAGTGCTTCAGCCACAGG

>SGPR469_SEQID_58

GGAGATTCTGGGGGCCCTGGTCTGTGAATTAATGGCACATGGGTCAGGTGGGATTGTGAGCTGGGGCA
TTGGCTGCGGTGCAAGGATACCCCTGGAGTTTACACAGAAGTTAGTTTCTACAAGAAATGGATT

>SGPR400_SEQID_59

ATGGCAGGAGACAGGTACCGCCCAATGTCAGCAGATACCCCTGGACAGAAACGATGTCTCTTCCTGAAAAAAC
ATTTCTCCTTTCTATAGAGCATCACTCCTTGCTGTTGTAAACACACAGATCCAATAATAGTCGTGGCGAGCTTTT
GAGAGTCAGGTTCTCCCGATTGACAGCAGGGGACGCCGACACCCCAATTCCTCCCTTGGGTCCCTGGAG
CTGCACCTCTGAAGTCTGGTCCCTTCAAGATCTGGCAGGGGTGAAGACCAAGAGAGGAGGGGGACAGAGA
CACGGCACTGCTGCTATGCATTACGCTGCTCCTTCTGCTGGGATTCGGGTGAGCCCCCAGAAATGGTCT
GTGGCGGGCCACAGTCTCATCTGGTATTGCCTCAGGCTTGGGGCTAGTGTGGGCGAGTGGCCCTGGCAGGT
CAGCATCCGCCAGGCTTGATTCACGTCTGCTCAGATACCCCTCATCTCAGAGGAGTGGTGCTGACAGTGGCGA
TCTGCTTCCATTATCCCCCACCTGATTTCCAAGCAACACATCTAGTGCCATCGCTGTGTAGAACTGCCCT
CCCCAGTTTCTGTAGCCCTGTGCTGCTCATCTGCCCTCCCTCATCTGAAGTCACTGAAAGAAATACAAC
CTCCTGCTGGTGACTGGGTATAGGGCTATAGTGAATATCCAAATATCAAGCGTTCTTATACACTGAAGGAGCT
GAAAGTCCCCCTCATTGATCTCCAGACATCGGTGACCACTATCAAAATGAAATCTTGTGCACGGAGTTGAGCT
CATCATCAGTGAAGCTATGATCTGCTCCAAGCTCCAGTGGGCGAGATGGATCAGTGTACTGTAAAGATCCACC
CCTCAGGCACCTTTCACAGGCCCTTGCCTTCCCCAGTGA

FIG. 2A

>SGPr397_SEQID_60
 MKCLGKRRGQAAFLPLCWLFKILQPGHSHLYNNRYAGDKVIRFIPKTEEEAYALKKISYQLKVDLWQPSSISYVSEG
 TVTDVHIPQNGSRALLAFLQEANIQYKVLIEDLQKLEKSSLHTQRNRRSLSGYNYEVYHSLEEIQNWMMHHLNKTSH
 GLHMFSGRSYEGRCFLIKLGRSRRLKRAVWIDCGIHAREWIGPAFCQWFKVKEALLTYKSDPAMRKMLNHLFYFIMP
 VFNVGYYHFSWTNDRFWRKTRSRNSRFRRCRGVDANRNWVKWCGKFGTNDWDPDKVSAGFTLQNMSPEDSHGR
 LMFFCM

>SGPr413_SEQID_61
 MKPLLETLYLLGMLVPGGLGYDRSLAQHRQEIVDKSVSPWSLETYSYNIYHPMGEIYEWREISEKYKEVVTQHFLGV
 TYETHPMYYLKISQPSGNPKKIWMDCGIHAREWIAFCQWFKVKEILQNHKDNSSIRKLLRNLDYVLPVLNIDGYIYT
 WTTDRLWRKSRSPHNGTCFGTDLNRNFNASWCISGASRNCQDQTFCGTGPVSEPETKAVASFIKSKDDILCFLTM
 HSYGQLILTPYGYTKNKSSNHPEMIQVGQKAANALKAKYGTNYRVGSSADILYASSGSSRDWARDIGIPFSYTFELRD
 SGTYGFVLPEAQIQPTCEETMEAVLSVLDDVYAKHWHSDSAGRVTSATMLLGLLVSCMSLL

>SGPr404_SEQID_62
 MVSNDSTWTVTKNGSGDMIFEGNSEKEIPVLNPLVPMVARYIRINPQSWFDNGSICMRMEILGCPLDPNNYYHR
 RNEMTTDDLDLDFKHHNYKEMRQMLKVVNEMCPNITRIYNIKSHQGLKLYAVEISDHPEGEHEVGEPEFHYIAGAHGNE
 VLGRELLLLLVQFVCQEYLARNARIVHLVEETRIHVLPSLNPDPGYEKAYEGGSELGGWSLGRWTHDGINNNFPDLNT
 LLWEAEDRQNVPRKVPNHYIAPEWFLSENATVAETRAVIAWMEKIPFVLGGNLQGGELVAYPYDLVRSPWKTQEH
 TPTPDDHVFRWLAYSASTHRLMTDARRRVCHTEDFQKEEGTVNGASWHTVAGSLNDFSYLHTNCFELSIYVGCDK
 YPHESQLPEEWENNRESLIVFMEQVHRGIKGLVRDSHGKIPNAISVEGINHDIRTANDGDYWRLLNPGEYVVTAKAE
 GFTASTKNCMVGYDMGATRCDFTLSTKNMARIREIMEKFGKQPVSLPARRLKLGRKRQRG

>SGPr536_1_SEQID_63
 MWRCPLGLLLPLAGHLALGAQQGRGRRELAPGLHLRGIRDAGGRYCEQDLCCRGRADDCALPYLGAICYCDLF
 CNRTVSDCCPDFWDFCLGVPFPPIQCGMHGGRIYPVLGTYWDNCNRCTCQENRQWQCDEPCLVDPDMIKAIN
 QGNYGWQAGNHSFAWGMTLDEGIRYRLGTIRPSSVMNMHEIYTVLNPGEVLPATAFEASEKWPNIHEPLDQGNCA
 GSWAFSTAAVASDRVSIHSLGHMTPLVSPQNLLSCDTHQQCGCRGGRLDGAWWFLRRRGVSDHCYPFSGRERDE
 AGPAPPCMMHSRAMGRGKRQATAHCPNSYVNNNDIYQVTPYRLGSNDKEMKELMENGVPQALMEVHEDFFLYK
 GGIYSHTPVSLGRPERYRRHGTHSVKITGWGEETLPDGRTLKYWTAAANSWGPWAGERGHFRIVRGVNECDIESFVL
 GWGRVGMEDMGHH

FIG. 2B

>SGPr414_SEQID_64
MCENCADLVEVLNEISDVEGGDGLQLRKEHTLKIFTYINSWTQRQCLCCFKEYKHLEIFNQVVCALINLVIAQVQVLRD
QLCKHCTTINIDSTWQDESNAQEEPLNIDRECNEGSTERQKSIEKKSSTRICNLTEEESSKSSDPFSLWSTDEKEKLL
LCVAKIFIQIFPLYTAYKHNTHPTIEDISTQESNILGAFCDMNDVEVPLHLRYVCLFCGKNGLSLMDKCFEYGTPTPLP
FLIAHAFITVSNIRIWLHIPAVMQHIIPFRTYVIRYLCKLSDQELRQSAARNMADLMWSTVKEPLDTTLCFDKESLDLAF
KYFMSPTLTMRLAGLSQITNQLHTFNDVCNNESSLVSDTETSIKELADWLISNNVVEHIFGPNLHIEIKQCQVILNFLAAE
GRLSTQHIDCIWAAQQLKHCRIYHDLFPPLRLHLLNLVSALEPSVHTEQTLYLASMLIKALWNNALAAKAQ
LSKQSSFASLLNTNPIGNKKEEEELRRTAPSPWSPAASPQSSDNDTHQSGGSDIEMDEQLINRTKHVQQRLSDTEE
SMQSSDETANSGEDGSSGPGSSSGHSDGSSNEVNSSHASQAGSPGSEVQSEDIADIEALKEEDEDHGHNP
KSSCGTDLNRKLESQAGICLGDSSGTSENGTSSGTGKDLVFNTESLPSVDNRMRLDACSHSEDPEDHISGEMN
ATHIAQGSQESCITRTGDFLGETIGNELFNCRQFIGPQHSHHHHHHHHGHMVDDMLSADDVSCSSSQVSAKSEK
NMADFDGEESGCEEEVLQINSHAELTSHLQQHLPNLASIYHEHLSQGPVVKHGHFNQSNVAVTDINLDNVCKKGNLTLLW
DIVQDEDAVNLSGLINEAEKLLCSLVCWFTDRQIRMRFIEGCLNNGNRSVVISRLPKLFGTFQFGSSYDTHWIT
MWAEEKLNMMKLFDDNLVYIQTVREGROKHALYSHSAEVQVRLQFLTQVSTLGSDFHRLSLEQVDILWHCLVED
SECYDDALHWFLNQVRSDQHMGMEETYKHLFLEKMPQLKPETISMTGLNLFQHLNCLARLATSAYDGCNSSELCG
MDQFWGIALRAQSGDVSRRAIQYINSYINGKTGLEKEQEFISKMESLMIASSLEQESHSSLMVIERGLMLKTHLE
AFRRRFAYHLRQWQIEGTGISSHLKALSDKQSLPLRVVCPAGLPDKMTIEMYPDSQVADLRAEVTHWYENLQKEQI
NQQAQLQEFQGSNRKGEFFPGGLMGPVRMISSGHELTDDYDEKALHELGFKDMQMVFSVSLGAPRRERKGEVQLPA
SCLPPPQKDNIPMLLLQEPHLTTLFDLLEMLASFQPPSGKVAVDDSESLRCEELHHAENLSRRVWELLMLPTCPN
MLMAFQNISDEQSFKAQSDHRSRHEVSHYSMWLLVSWAHCCSLVKSSLADSDHLQDWLKKLTLLIPETAVRHESCSG
LYKLSLGLDGGDSINRSFLLAASTLLKFLPDAQALKPIRIDDDYEEEPILKPGCKEYFWLLCKLVDNHDKDASQTTLLDL
DALARHLADCIRSREILDHQDGNVEDDGLTGLRLATSVVKHKPPKFSREGQEFLRDIFNLLFLLPSLKDRQQPKCKS
HSSRAAYDLLVEMVKGSVENYRLIHNWMAQHMQSHAPYKWDYWPHEDEVRAECRFVGLTNLGATCYLASTIQQLY
MIPERQAVFTAKYSEDMKHKTLLLELQKMFYLMSECKAYNPRPFCKTYTMDKQPLNTGEQKDMTEFFTDLTIKIE
EMSPELKNTVKSFLGGVITNNVSLDCEHVSTAEFFYTRCQVADMKNYESLDEVTKDTELDGNMYTCSQCGKKV
RAEKRACFKKLPRIXSFNTMRYTFNMVTMMKEKVNTHFSFPLRLDMTPYTEDFLMGKSERKEGFEVSDHSDKSEY
EYDLIGVTHTGTADGGHYYSFIRDIVNPHAYKNKWKWYLFNDAEVKPFDASAQLASECFGGEMTTKTYDSVTDKFMDFS
FEKTHSAYMLFYKRMEPEEEENGREYKFDVSSLELWHDNMQFLQDKNIFEHTYFGFMWQLCSCIPSTLPDPKAVS
LMTAKLSTSFVLETFHSKEKPTMLQWIELLTKQFNNSQAACEWFLDRMADDDWWPMQILKCPNQIVRQMFQRLCIH
VIQRLRPVHAHLYLQPGMEDGSDMDTSDVEDIGGRSCVTRFVRTLLIMEHGVKPHSKHLTEYFAFLYEFKAKMGEEES
QFLLSLQAISTMVHFYMGTKGPENPQVEVLSEEEGEEEEEDILSLAEKRYRPAALEKMIALLVVEQSRSEHRLTL
SQTDMAALTGGKGFPLFQHIRDGINIRQTCNLIFSLCRYNNRLAEHIVSMLFTSIAKLTPAANPFFKLLTMLMEFAGG

FIG. 2C

PPGMPFASYILQRIWEIEYNPSQCLDWLAVQTNRKLAHSWWLQNMENWVERFLLAHNYPRVRTSAAYLLVSLIPS
NSFRQMFRSTRSLHIPTRDLPLSPDTTVLHQVYNVLLGLLSRAKLYVDAAVHGTTKLVYPYFSFMTYCLISKTEKLMFS
TYFMDLWNLFQPKLSEPAIATNHNKQALLSFYWNVCADCPENIRLVQNPVWTKNIAFNYLADHDDQDQVWLFNRGMLP
AAYGILRLCCEQSPAFTRQLASHQNIQWAFKNLTPHASQYPGAVEELFNLMLQFIAQRPMREEELEDIKQFKKTTISC
YLRCLDGRSCWTTLSAFRILLESDEDRLLVFNRLGILMTESFNTLHMMYHEATACHVTGDLVELLSIFLSVLKSTRPY
LQRKDVQALIQWQERIEFAHKLLTLNSYSPPELRNACIDVLKELVLLSPHDFLHTLVFLLQHNHCTYHHSNIPMSLGP
YFPCRENIKLIGGKSNIRPPRPELNMCLPTMVETSKGDDVYDRMLLDYFFSYHQFIHLLCRVAINCEKFTETLVKLSV
LVAYEGLPLHALFPKLWTELCQTQSAMSKNICKLLCEDPVFAEYIKCILMDERTFLNNNIVYTFMTHFLLVQSQVFSE
ANCANLISTLITNLISQYNLQSDFSNRVEISKASASLNGDLRALALLSVHTPKQLNPALPTLQELLKSKRTCLQQRNS
LQEQEAKERKTKDDEGATPIKRRRVSSDEEHTVDSCISDMKTETREVLTPSTSDNETRDSIIDPGTEQDLPSPENSS
VKEYRMEVPSSFSEDMNIRSQHAEEQNNGRYDDCKEFKDLHCKSKDSTLAESEFPSTISAVLSDLADLRSCDG
QALPSQDPEVALSLSCGHSRGLFSHMQQHDILDTLCRTIESTIHVTRISGKGNQAAS

>SGPr430_SEQID_65

MSPLKIHGPIRIRSMQTGITKWKEGSFEIVEKENKVSLLVHYNTGGIPRIFQLSHNIKNVLRPSGAKQSRLMLTLQDNS
FLSIDKVPKDAEEMRLFLDAVHQNRPLPAAMKPSQSGSGFAGILGSRTSQKETSRLQSYSDNQASAKRGSLETCKDIP
FRKVLGNPGRGSIKTVAGSGIARTIPSLTSTSTPLRSGLENRTEKRMISTGSELNEDYPKENDSSNNKAMTDPSR
KYLTSREKQLSLKQSEENRTSGLLPLQSSSFYGSRAGSKEHSSGGTNLDRTNVSSQTPSAKRSGLFPQVPVPLSV
KKLRCNQDYTGWNKPRVPLSSHQQQLQGFSNLGNTCYMNAILQSLFSLQSFANDLLKQGPWKKIPLNALIRFAHL
LVKDDICNSETKKDLLKVKNAISATAERFSGYMQNDAHEFLSQCLDQLKEDMEKLNKTWKTETPVSGEENSPDISATR
AYTCPVITNLEFEVQHSIICKACGEIPKREQFNDSLIDLP RRKKPLPPRSIQDSLDFRFAEELEYSCCKCGKCALVRH
KFNRLPRVLIHLKRYSFNVALSLNNKIGQQVIIPRYLTLSHCTENTKPPFTLGWSAHMAMSRPLKASQMVNSCITSPS
TPSKKFTFKSKSSALCLDSDSEDELKRSVALSQRCEMLGNEQQQEDLEKDSKLCPIEPDKSELENSGFDRMSEEL
LAAVLEISKRDASPSLSHEDDDKPTSSPDTGFAEDDIQEMPENPDTMETEKPKITELDPASFTEITKDCDENKENKTP
EGSQGEVDWLQQYDMEREREELQQAQSLQEQEAWQEKEDDDDLKRATELSLQEFNNSFVDALGSDDEDSGNE
DVFDMEYTEAEAEELKRNAETGNLPHSYRLISVWSHIGSTSSSGHYISDVYDIKKQAWFTYNDLEVSKIQEAAVQSDRD
RSGYIFFYMHKEIFDELLETEKNSQSLSTEVGKTTTQAAS

>SGPr496_1_SEQID_66

MTLLAPWYTGPMPMDVNEPSSVTTAPTLSSSLQHISFLATGKKLSLHFGHPRECEVTRIDDKNRRGLEDSEPGAKL
FNNDGVCCCLQKRGPNITSCVSPRTLQISVFLSEKYEKVFESDELFPFGVIGSNIGDAHFQEFRAGISWKPVWDP
DDPIQFPDCCSSSSRIPSVSVLVAVPLVAGHKGAQFIERMLGCFKELKQELTQEGPGGGHPRSAWPPRRHAQWP

FIG. 2D

PEPCEQGEPPVPEAEVEEAETAERKVEAEKVEGKAEAGKAEAGKVDATEKVEVETAGKVDAAGKVETAEG
 PGRRAELKLEPEPEVREAEQEPKQELEDEDENPARSGGGNSDEVPPPTLPSDPPRPPDPSRRSRAPRRPRPQ
 TRLRTPQPRPPRRPPRGPGGGCLDVDFAVGPPGCSHVNSFKVGENWRQELRVYQCFFVWCGTPETRKSKA
 KSCICHVCGTHLNRHSLSCVFFGCFTEKIHIEHAETKQHNLAVDLYYGGYCFMCKDYVYDKDIEQIAKEEQGEALK
 LQASTSTEVSHQQCSVPGLGEKFTWETTKPELELLGHNPRRRRITSSFTIGRLGLNLGNTCFMNCIVQALHTHPILRD
 FFLSDRHRCEMPSELCLVCEMSSLFREL YSGNPSHPYKLLHLVWIHARHLAGYRQQDAHEFLAALDVLHRHCKG
 DDVGKAANNPHNCNCIDQIFTGGLQSDVTCQACHGVSTTIDPCWDISLDLPGSCTSFWPMSPGRESSVNGESHIPGI
 TTLTDCRLRRFTRPEHLGSSAKIKGSCQSYQESTKQLTMNKL PVACFHFKRFEHSAKQRRKITYISFPLELDMTPFM
 ASSKESRMNGQLQLPTNSGNNENKYSLFAVNVHQGTLESGHYTSFIRHHKQWFKCDDAVITKASIKDVLDSGEYLLF
 YHKQVLEHESEKVKEMNTQAY

>SGPr495_SEQID_67

MRVKDPTKALPEKAKRSKRPTVPHDEDDSDDIAGVLTQHVSHAISVNHVKRAIENLWSVCSELEERRFYDGGQLVL
 TSDIWLCLKCGFGCGKNSSEQHSLKHFKSSRTEPHCIINLSTWIIWCYECDEKLSTHCNKKVLAQIVDFLQKHASKT
 QTSAFSRIMKLCCEEKCEIDEIQGGKCRNL SVRGITNLGNTCCFNAMQNLAQTYTLTDLMEIKESSTKLKIFPSSDS
 QLDPLVELSRPGPLTSALFLFLHSMKETEGPLSPKVLFNQLCQKAPRFKDFQQQDSQELLHYLLDAVRTTEETKRIQ
 ASILKAFNNPTTKTADDETRKVKVISTVKDPFIDISLPIIEERVSFKPLLWGRMNKYRSLRETDHSDRYSGNVTIENIHQPR
 AKKHSSSKDKSQLIHDRKICIRKLSSETVITYQKNENLEMNGDSL MFASLMNSESRLNESPTDDSEKEASHESNVDA
 DSEPSSESASKQTGLFRSSSGGVQPDGPLYPLSAGKLLYTKETDSGDKEMAEASELRLSSTVTGDDQDFDRENQP
 LNISNLCFLEGKHLRSYSPQNAFQTLSSQSYITTSKECSIQSCLYQFTSMELLMGNNKLLCENCTKNKQYQEETSFAE
 KKVEGVYTNARKQLLISAVPAVLILHLKRFHQAGLSLRKVNHRHVDPLMLDLAPFCSATCKNASVGDKVLYGLYGIVEH
 SGSMREGHYTAYVKVRTPSRKLSEHNTKKKNVPGLKAADSESAGQWVHVSDTYLQVWPESRALSQAQYLLFYERV

>SGPr407_SEQID_68

MEYPVPYFRSPNRTLPERIWSNPLLVIAKYTVSWPRQQLAKQANKWMPFVIPSCTLPWDPLELKICYQQRNPYP
 PDPSNFTFLRCLNAFAAVFYLPQPSWHKPEGLKPAGYPRVPDIPYSGGYTLKSTTEAGLHQSLPMVQLPLHPTKG
 SALLKESELNDADWANL MWKRYLEEQEDSKMVDLFVGMKSYLKCQACGYHSMFTKVFVFFCDLSLTIPKKGFAGGK
 VSLRDCLSLFTKEEELELENASGTLPTVKSEVLSTSCVPFGTTQAASTVATTQPCASARLVGTFTMTLVSPNLTRDTE
 GIELTVMKALVDILFKASTDIILFNHDSSSGNKWRKLPEPGGLEKKHEELRLRPLKEEYHWLVLPKL TGSPHRWRP
 RKRALASCSCWCLQRVTMRVMGVQDQKAGNRNQMLLGQRPVIGDVTVNSQTTTRDKACRRPPSHSVFTQSSFWACL
 DPDLFFYGHQSYWMKAHLNDLILREGPVTQMAQSFYWGFPAGGNLSALEMLPDGPAPRTFLQKKSCLFPLFSYILLH
 KAGKLFQPDAGHGLVVKVHAPTRGIVFIMEPRQLGGKSLKLQACALGGMNSGMEPQKSAPFAAGKGLAPPLPVC

FIG. 2E

NLRFKLRVYKFEELWSRAGLGKKSDNHSSRQMPWGAAQVACQHPCKLPRIVAEITPPKLSFGFLNTVQSSVLPTSL
SQFFLNDSPQEEAIPQSLPGSPRTNSFPKDKFVPKDKLVLSLLTMYELDRLF

>SGP453_SEQID_69

MLAMDTCKHVGLQLAQDHSSLNPQKWHCVDCNTTESIWACLSCSHVACGRYIEEHALKHFQESSHPVALEVNEMY
VFCYLCDDYVNDNATGDLKLLRRTLSAIKSQNYHCTTRSGRFLRSMGTGDDSYFLHGAQSLLQSEDQLYTALWHR
RRILMGIFRTWFEQSPIGRKKQEEPFQEKIVKREVKKRRQEELEYQVKAELSMPPRKSRLQLGQAQSTIIEIVSVQVP
AQTPASPAKDKVLSTSENEISQKVSDDSSVKRRPIVTPGVTGLRNLGNTCYMNSVLQVLSHLLIFRQCFLKLDLNQWLA
MTASEKTRSKHPPVTDTVYQMNCEQEKDTGFVCSRQSSLSGLSGGASKGRKMELIQPKETSQYISLCHLHTL
FQVMWSGKWALVSPFAMLSVWRLIPAFRGYAQQDAQEFCELLDKIQRELETTGTSLPALIPTSQRKLIKQVLNVVN
NIFHGQLSQVTCLACDNKSNTIEPFWDLSLEFPERYQCSGKDIAQPCLVTEMLAKFTETEALLEGKIYVCDQCNSKRR
RFSSKPVVLTEAQQLMICHLPQVLRHLKFRFRWSGRNNREKIGVHVGFEEILNMEPYCCRETLKSLRPECFIYDLSAV
VMHKGFGSGHYTAYCYNSEGGFWVHCNDSKLSMCTMDEVCKAQAYILFYTQRVTENGHSKLLPPELLGSGHPN
EDADTSSNEILS

>SGP445_SEQID_70

MRVKDPTKALPEKAKRSKRTPVPHDESDDDIAVGLTCQHVSHAISVNHVKRAIENLWSVCSECLEERRFYDGGQLVL
TSDIWLCLKGFGQCGKNSQSHLKHFKSSRTEPHCIINLSTWIIWCYECDEKLSTHCNKVLAQIVDFLQKHASKT
QTSAFSRIMKLCEEKCETDEIQKGKCRNLVSRGITNLGNTCFFNAVQNLAQTYTLTDLMEIKESSTKLKIFPSSDS
QLDPLVVELSRPGPLTSALFLFLHSMKETEGPLSPKVLNQLCQKRVHLHLI

>SGP401_1_SEQID_71

MTVRNIASICNMGTNASALEKDIDGPEQFPINEHYFGLVNFNGNTCYCNSVLQALYFCRPFRENVLAYKAQQKKENLLTC
LADLFHSIATQKKKVGVIKKFISRLRKENDLFDNMQQDAHEFLNYLLNTIADILQEEKKQEKQNGKLNKGNMNEPA
ENNKPELTWVHEIFQGTLTNETRCLNCETVSSKDEDFDLSDVDEQNTSITHCLRDFSNTETLCSEQKYCETCCSKQ
EAQKRMVKKLPMLVALHLKRFKYMEQLRRTYTKLSYRVVFPLELRLFNTSSDAVNLDRLMYDLVAVVHCGSGPNRGH
YITIVKSHGFWLLFDDIVEKIDAAIEEFYGLTSDISKNSSEGYILFYQSRE

>SGP408_SEQID_72

MVPGEENQLVPKEAPLDHTSDKSLLDANFEPGKKNFLHLTDKDGEQPKILLEDSSAGEDSVHDFRFIGPLPREGSVGST
SDVVSQSYSSILNKSETGYVGLVNQAMTCYLNLSLLQTLFMTPEFRNALYKWEFESEEDPVTSPYQLQRLFVLLQT
SKKRAIETTDVTRSFSGWDSSEAWQQHDVQELCRMFDALQKWKQTEQADLINELYQGKLDYVRCLECGYEGWRI

FIG. 2F

DTYLDIPLIRPYGSSQAFASVEEALHAFIQPEILDGPNQYFCERCKKKCDARKGLRFLHFPYLLTLQLKRFDFDYTTMH
RIKNDRMFTPEELDMSTFIDVEDEKSPQTESCTDGAENEGSCHSDQMSNDFNSDDGVDEGICLETNSGTEKISKIS
GLEKNSLIYELFSVMAHSGAAGGHYYACIKSFSDQWYSFDDQHVSRTQEDIKKTHGGSSGRGYSSAFASSTNA
YMLIYRLKDPARNAKFLVGEYPEHIKNLVQKERELEEQEKQRQERINTCKIKLFC LHPTKQVMMENKLEVHKDKTLK
EAVEMAYKMMDEEVIPLDCRLLVKYDEFHDLERSYEGEEDTPMGLLLGGVKSTYMFDLLLETRKPDQVFSYKPG
EVMVKVHVVDLKAESVAAPITVRAYLNQTVTEFKQLISKAIHLPAETMRVLERCYNDLRLSVSSSKTLKAEGFFRSNKV
FVESSETLDYQMAFADSHLWKLDRHANTIRLVLPEQSPVSYSKRTAYQKAGGDSGNVDDDCERVKGPVGSLSKV
EAILLESTEKLSLSLQQQDGDNGDSSKSTETSDFENIESPLNERDSSASVDNRELEQHIQTSDPENFQSEERSDSD
VNNDRSTSSVDSDILSSSHSDTLNADNAQIPLANGLDSSHITSRRRTKANEGKKTWDTAEEDSGTDSYDESGKS
RGEMQYMYFKAEPYAADGSGEGHGWLMVHVDKRITLAAFKQHLEPFVGVLSHFVKVFRVYASNQEFESVRLNETL
SSFSDDNKITIRLGRALKKGEYRVKYQLLVNEQEPCKFLLDVAFKGMTVRQSKKEELPQLREQCGLELSIDRFRLRK
KTWKNPQTVFLDYHIYEEDINISSNWEVFLVDGVEKMKMSQLAVLSRRWKPSMKLDPFQEVVLESSSVDELREK
LSEISGIPLDIEFAKGRGTFPCDISVLDIHQDLDNWPKVSTLNVWPLYICDDGAVIFYRDKTEELMELTDEQRNELMKK
ESSRLQKTGHRVTYSPRKEKALKIYLDGAPNKDLTQD

>SGP480_SEQID_73

MGAKESRIGFLSYEEALRRVTDVELRLKDAFKRTGGLSYMGQHCIFIREVLGDGVPKVAEVIYCSFGGTSKGLHFN
NLIVGLVLLTRGKDEEKAKYIFSLFSSESGNYVIREEMERMLHVVDGKVPDTRKCFSEGEKVNEYKFRNWLFNLKDAF
TFSRWLLSGGVVTLTDDSDTPTFYQTLAGVTHLEESDIIDLEKRYWLLKAQSRRTGRFDLETFGLVSPPIRPSLSEGL
FNAFDENRDNHIDFKEISCGLSACCRGPLAERQKCFKVFVDVDRDGLSRVELRDMVVALLEVWKNRDTDDIPELHM
DLSDIVEGILNAHDTTKMGHLLTLEDYQIWSVKNVLANEFNLFFQVCHIVLGLRPATPEEEGQIIRGWLERESRYGLQA
GHNWFIISMQWWQQWKEYVKYDANPVVIEPSSVLNGGKYSFGTAHPMEQVEDRIGSSLSYVNTTEEFSDNISTAS
EASETAGSGFLYSATPGADVCFARQHNTSDNNNQCLLGANGNILLHLNPQKPGAIDNQPLVTQEPVKATSLTLEGRL
KRTPQLIHGRDYEMPEPVWRALYHWYGANLALPRPVKNKSDIPELELFPYLLFLRQQPATRTQQSNWVNMGNV
PSPNAPLKRVLAYTGCFSRMQTKEIHEYLSQRRLRIKEEDMRLWLYNSENLYLLDDEDHKLEYLKIQDEQHLVIEVRNK
DMSWPEEMSFIANSSKIDRHKVPTEKGATGLSNLNTCFMNSSIQCVSNTQPLTQYFISGRHLYELNRTNPIGMKGHM
AKCYGDLVQELWSGTQKNVAPLKLRTWIAKYAPRFNGFQQQDSQELLAFLLDGLHEDLNRVHEKPYVELKDSDBG
DWEVAAEAWDNHLRRNRSIWDLFHGLRQSVKCKTCGHISVRFDPFNLPLPMDSYMHLITVILKLDGTTTPVRYG
LRLNMDEKYTGKQLSDLCGLNSEQILLAEVHGSIKIFPDQDNQKVRLSVSGFLCAFEIPVPVSPISASSTQTDSS
SPSTNEMFTLTNGDLPRPIFIPNGMPNTVPCGTEKNTNGMVNGHMPSLPDSPTGYIAVHRKMMRTEL YFLSSQ
KNRPSLFGMPLIVPCTVHTRKKDL YDAVWVQVSRSLASPLPQEAASNAHQDCDDSMGYQYPFTLRVVKDGNSCAWC
PWYRFCRGCKIDCGEDRAFINAYIAVDWDPTALHLRYQTSQERVVDEHESVEQSRRAQAEINLDSCLRAFTSEEL

FIG. 2G

GENEMYCSKCKTHCLATKLDLWRLPPIIHLKRFQFVNGRWIKSQKIVKFPRESFDPFAFLVPRDPALCQHKPLTP
 QGDELSEPRILAREVKVDAQSSAGEEDVLLSKSPSSLSANISSPKGSPSSSRKSGTSCPSSKNSSPNSSPRTLGRS
 KGRRLPQIGSKNKLSSKENLDASKENGAGQICELADALSRGHVLGGSQPELVTPQDHEVALANGFLYEHEACGNG
 YSNGQLGNHSEEDSTDQREDTRIKPIYNLYAISCHSGILGGGHYVYAKNPCKWYCYNDSSCKELHPDEIDTDSAY
 ILFYEQQGIDYAFQLPKTDGKKMADTSSMDEDFESDYKKYCVLQ

>SGPr431_SEQID_74

MDKILEGLVSSSHPLPLKRIVVRKVVEAHEWLDEAQCEAMFDLTTRLILEGQDPFQRQVGHQVLEAYARYHRPEFES
 FFNKTFLVGLLHQGYHSLDRKDVAILDYIHNGKLIMSCPSVLDLFSLLQVEVLRMVCEPERPEQLCARLSDLLTDFVQCI
 PKGKLSITFCQQLVRTIGHFQCVSTQERELREYVSVQTKVSNLLQNIWKAEPATLLPSLQEVFASISSTDASFEPVALA
 SLVQHPLQMITVLRSLTTPDPNVKDSMTQALCRMIDWLSWPLAQHVDTWVIALKGLAAVQKFTILIDVTLKIELVFN
 RLWFPLVRPGALAVLSHMLLSFQHSPEAFHLVPHVNVLSHFKNDGLPSSTAFVLVQLTELIHCCMMYHYSGFDPDLYEPI
 LEAKDFPKPSEEKILILNQSAWTSQSNLSASCLSRISGKSETGKTGLINLNTCYMNSVIAQALFMTDFRRQVLSLNL
 NGCNSLMKKLQHLFAFLAHTQREAYAPRIFFEASRPWFTRSQQDCSEYLRFLDLRLHEEEKILKVQASHKPSIELEC
 SETSLQEVASKAAVLTTETPRTSDGEKTLIEKMFGGKLRTHIRCLNCRSTSQKVEAFDLSLAFCPSSSLENMSVQDPAS
 SPSIQDGGGLMQASVPGPSEEPVYVNPPTAAFCIDSLVNEKTIGSPNEFYCSENTSVPNESKILVNKDVQKPGGET
 TPSVTDLLNYFLAPEILTGDNQYYCENCASLQNAEKTMQITEEPEYLILTLRFSYDQKYHVRKILDVNSLPLVLELPVK
 RITSFSSLSSESWSVDFTDLSENLAKKLKPSGTDEASCTKLVYLLSSVWHSGISSESGHYYSYARNITSTDSSYQM
 YHQSEALALASSQSLLGRDPSAVFEQDLENKEMSKEWFLNDSRVFTFSQSVQKITSRFPKDTAYVLLYKKQHST
 NGLSGNNPTSGLWINGDPPLQKELMDAITKDNKLYLQEQELNARARALQAASASCSPRPNGFDDNDPPGSCGPTGG
 GGGGGFNTVGRLVF

>SGPr429_SEQID_75

MAPRLQLEKAAWRWAETVRPEEVSQEHIETAYRIWLEPCIRGVCCRCKGNPNCLVGIGEHWLGEIDENSFHNIDDP
 NCERRKNNSFVGLTNLGATCYNTFLQVWFLNLELRQALYLCPSTCSDYMLGDGQEEKDYEPQTICEHLQYLFALLQ
 NSNRRYIDPSGFVKALGLDTGQQQDAQEFKLFMSLLEDTLISKQKNPDVRNIVQQQFCGEYAYVTVCNQCGRESKLL
 SKFYELNLQGHKQLTDCISEFLKEEKLEGNRYFCENCQSKQNAIRKIRLLSLPCTNLQLMRFVFDRTGTHKKLN
 TYIGFSEILDMPEYVEHKGSYYELSAVLIHRGVSAYSGHYIAHVKDPQSGEWYKFNDIEDIEKMEGKKLQLGIEEDLE
 PSKSQTRPKCKGKGTCHSRNAYMLVYRLQTQEKPNNTTVQVPAFLQELVDRDNSKFFEEWCIEMAEMRKQSVDKGAK
 HEEVKELYQRLPAGAEPEYFVSLEWLQKWLDSTPTKPIDNHACLCSHDKLHPDKISIMKRISYAADIFYSRYGGGPR
 LTVKALCKECVVERCRILRLKNQLNEDYKTVNNLLKAAVKGDFWVGKSSLSRWRQALALEQLDEQDGAEQSNGKM
 NGSTLNKDESKEERKEEEELNFNEDILCPHGELCISENERRLVSKEAWSKLQQYFPAPEFPSYKECCSQCKILEREG

FIG. 2H

EENEALHKMIANEQKTSPLNLFQDKNRPCLSNWPEDTDVLYVQQFFVEEWRKFVRKPTRCSPVSSVGNALLCPHG
GLMFTFASMTKEDSKLIALWPSEWQMIQKLFVDHVIKITRIEVDVNPSETQYISEPKLCPECREGLLCQQQRDLREY
TQATYVHKVDNKKVMKDSAPELNVSSSETEEDKEEAKPDGEKDPDFNQSNNGGTRKQKISHQNYIAYQKQVIRRRSM
RHRKVRGEKALLVSANQTLKELKIQIMHAFSVAPFDQNLSIDGKILSDDCATLGLGVIPESVILLKADEPIADYAAAMDDV
MQVCMPEEGFKGTGLLGH

>SGPr503_SEQID_76

MLSSRAEAMTAADRAIQRLRTGAAVRYKVMKNWGVIGGIAAALAAAGIYVIWGPITERKKRRKGLVPLVNLGNTCF
MNSLLQGLSACPAFIRWLEEFTSQYSRDQKEPPSHQYLSLTLLHLLKALSCQEVTDDEVLDASCLLDVLRMYRWQISS
FEEQDAHELHFHVTSSLEDERDRQPRVTHLFDVHSLEQQSEITPKQITCRTRGSPHPTSNHWKSQLPFFHGRLTSNMV
CKHCEHQSPVREDFDLSLSIPAAWGHPLTLDHCLHFFISSESRDVCDNCKIEAKGTLNGEKVEHQRTTTFVKQ
LKLGLPQCLCIHLQRLSWSSHGTPLKRHEHVQFNEFLMMDIYKYHLLGHKPSQHNPKNKNPGPTLELQDGPAGPT
PVLNQPGAPKTQIFMNGACSPSLLPTLSAPMPFPLPVVPDYSSSTYLFRLMAVWVHHGDMHSGHFVTYRRSPPSARN
PLSTSNQWLWVSDDTVRKASLQEVLSSSAYLLFYERVLSRMQHQSCECKSEE

>SGPr427_SEQID_77

MDLPGDGAAGGGPLAPRRRRRLRFLSRLLALGSRSRPGDSPRPQPGHCDGDGEGGFACAPGPVPAAPGSP
GEERPPGPQQLQLPAGDGARPPGAQGLKNHGNTCFMNAVQCLSDNTDLLAEFLALGRYRAAPGRAEVTEQLAALV
RALWTREYTPQLSAEFKNAVSKYGSQFQGNHQHDALEFLLWLLDRVHEDLEGSSRGPVSEKLPPEATKTSNCLSPS
AQLPLGQSFVQSHFQAQYRSSLTCPHCLKQSNFTDFPFLCVSLPIPLRQTRFLSVTLVFPSSQRFRLRVGLAVPILSTVA
ALRKMVAEEGGVPADDEVILVELYPSGFQSFDEEDLNTIAEGDNVYAFQVPPSPSQGTLSAHLGLSASPRLAAREG
QRFSLSHSESKVILFCNLVSGGQQAARFGPPFLIREDRAVSWAQLQQSILSKVRHLMKSEAPVQNLGSLFSIRVGL
SVACSYLSPKDSRPLCHWAVDRVLHLRRPGPPPHVKLAWEWDSSVKERLFGSLQEERAQDADSVWQQQAHHQHH
SCTLDECQFYTKEEQLAQDDAWKCPHCQVLQQGMVKLSLWTLPDILIHILKRFQVGERRNKLSLTKFPLSGLNMA
PHVAQRSTPEAGLGPWPSWKQPDCLPTSYPLDFLYAVCNHHGNLQGGHYTAYCRNSLDGQWYSYDDSTVEP
LREDEVNTRGAYILFYQKRNSIPPWSASSSMRSTSSSLSDHWLLRLGSHAGSTRGSLLSWSSAPCPSLPQVPDSPIF
TNSLCNQEKGGLPRRLVRGVKGRSISMKAPTTSRKQGPFTKMLRWSFGSKEKPPGASVELVEYLESRRRPRST
SQSIVSLLTGTAGEDEKASPRSNVALPANSEDDGGAIERGPAGVPCPSAQPNHCLAPGNSDGPNTARKLKENAGQD
IKLPRKFDLPLTMPSVEHEKPARPEGQKAMNWKESFQMGSKSSPPSPYMGFSGNSKDSRRGTSELDRLQGTTL
LRSVFRKKENRRNERAEVSPQVPPVSLVSGGLSPAMDGGQAPGSPPALRIPEGLARGLGSLRLERDVWSAPSSLRPR
KASRAPRGSALGMSQRTVPGEQASYGTQFRVKYHTLSLGRKKTLPESSF

FIG. 21

>SGPr092_SEQID_78
 MQLVLRVTIFLPWCFAVPVPPAADHKGWDFVEGYFHQFFLTKKESPLLTQETQTQLLQQFHRNGTDLLDMQMHAL
 HQPHCGVPDGSSTISPGRCKWNKHTLTIRIINYPHDMKPSAVKDSIYNAVSIWSNVTPLIFFQQVQNGDADIKVSWFQ
 WAHEDGWPFDPGGILGHAFLPNSGNPGVVHFDKNEHWSASDTGYNLFLVATHEIGHSLGLQHSGNQSSIMYPTYW
 YHDPRTFQLSADDIQRILHYGEKCSSDIP

>SGPr359_SEQID_79
 MKVLPASGLAVFLIMALKFSTAAPSLVAASPRTWNNYRLAQAYLDKYTYNKEGHQIGEMVARGSNSMIRKIKELQAF
 FGLQVTGKLDQTTMNVIKPRCGVPDVANYRLFPGEKWKNTLTYSKYTPSMSSVEVDKAVEMALQAWSSAVPL
 SFVRINSGEADIMISFENGHDGDSYPFDGPRGTLAHAFAPGEGLGDTDFDNPEKWTMTGTNGFNLTVAHEFGHAL
 GLAHSTDPALMYPTYKYKNPYGFHLPKDDVKGIQALYGPRKVFLGKPTLPHAPHKPSIPDLCDSSSFDVATMLGK
 ELLLFKDRIFWRRQVHLRTGIRPSTITSSFPQLMSNVDAAYEVAERGTAYFFKGPYHWTIRGFMQGPPTIYDFGFP
 RHVQQIDAAYVLRPQKTLFFVGDEYYSYDERKRKMEKDYPKNTEEEFSGVNGQIDAAVELNGYIYFFSGPKTYKYDT
 EKEDVSVVKSSSWIGC

>SGPr104_1_SEQID_80
 MNVALQELGAGSNMVEYKRATLRDEDAPETPVEGGASPDAMEVGFQKGTQRQLLGSRTQLLEVLGASLLAALLGC
 LVALGVQYHRDPSHSTCLTEACIRVAGKILESIDRGVSPCEDFYQFSCGGWRRNPLPDGRSRWNTFNSLWDQNQAI
 LKHLENTTFNSSEAEQKTQRFYLSCLQVERIEELGAQPLRDLIEKIGGWNITGPWDQDNFMEVLKAVAGTYRATPFF
 TVYSISADSKSSNSNVIQDQGLFLPSRDYYLNRANKEVLTAYLDYMEELGMLLGGRPSTSTREQMQQVLELEIQLANI
 TVPQDQRRDEEKYHKMSISELOALAPSMDWLEFLSFLSPLSESDSEPVVYGM DYLQQVSELINRTEPSILNNYLIW
 NLVQKTTSSLDRRFESAQEKLETLTYGTTKSCVPRWQTCISNTDDALGFALGSLFVKATFDRQSKIEAEGMISEIRTA
 EEALGQLVWMDEKTRQAAKEKADAIYDMIGFPDFILEPKELDDVYDGYEISEDSEFFQNMNLNLYNFSKVMADQLRKPP
 SRDQWSMTPTQTVNAYYLPKNEIVFPAGILQAPFYARNHPKALNFGGIGVWGMHELTHAFDDQGREYDKEGNLRPW
 WQNESLAAFRNHTACMEEQYNQYQVNGERLNGRQTLGENIADNGLKAAYNAYKAWLRKHGEEQQLPAVGLTNHQ
 LFFVGFAQVWCVSRTPESSHEGLVTDPHSPARFRVLGTLNSRDFLRHFGCPVGSMPNPGQLCEVW

>SGPr303_SEQID_81
 MPEKRPFERLPADVSPINCSLCLKPDLDDFTFEGKLEAAQVRQATNQIVMNCADIDIITASYPEGDEEIHATGFNYQN
 EDEKVTLSFPSTLQTGTGLKIDFVGELNDKMGFYRSKYTTPSGEVRYAAVTQFEATDARRAFPCWDERAIKATFDIS
 LVVPKDRVALSNMNVDRKPYPDENLVEVKFARTPTSTYLVAFVVG EYDFVETRSDGVCVCVYTPVGKAEQKGF
 ALEVAAKTLFPYNDYFNVVPYPLPKIDLIAIDFAAGAMENWDLVTYRETALLIDPKNSCSSSRQWVALVVGHELAHQWF

FIG. 2J

GNLVTMEWWTHLWLNFGFASWIEYLCVDHCFPEYDIWTQFVSADYTRAQELDALDNHPIEVSUGHPSVDEIFDAIS
YSKGASVIRMLHDYIGDKVKKKTLSI

>SGPr402_1_SEQID_82
MRPAPIALWLRVLALVRPRAVGWAPVRAPVYSSWAVQVSQGNREVERLARKFGFVNLGPIFDGQYFHLRHRG
VVQQLTPHWGHHHLKKNPKVQWFQQQLQRRVKRSVVPTDPWFQSKQWYMNSEAPDLSILQAWSQGLSGQGI
VSVLDDGIEKDHPDLWANYDPLASYDFNDYDPDPQPRYTPSKENRHGTRCAGEVAAMANNFGCGVGVAFNARIGG
VRMLDGTITDVIEAQSLSLQPQHHIYASWGPEDDGRITVDGPGILTREAFRRGVTKRGGGLTLFIWASGNGGLHYD
NCNCDGYTNSIHTLSVGSTTQQGRVPWYSEACASTLTITYSSGVAIDPQIVTTDLHHGCTDQHTGTSASAPLAAGMI
ALALEANPFLTWRDMQHLVVRASKPAHLQAEDWRTNGVGRQVSHHYGYGLLDAGLLVDTARTWLPTQPQRKCAVR
VQSRPTPILPIYIRENVSACAGLHNSIRSLHVVQAQLTLYSRRGDLEISLTSPMGTSTLVAIRPLDVSTEGYNNWVF
MSTHFWDENPQGVWTLGLENKGYFNTGTLRYTLLYGTAEEDMTARPTGPQVTSACVQRDTEGLCQACDGPAYI
LGQLCLAYCPPRFFNHTRLVTAGPGHTAAPALRVCSCHASCYTCRGGSPRDCSTCPPSSSTLDQQQSGCMGPTTPD
SRPRLRAAACPHHRCPASAMVLSLLAVTLGGPVLCGMSMDLPLYAWLSRARATPTKPQVWLPAGT

>SGPr434_SEQID_83
GPGRQGGCAGRRSTALPLRAPLRARRPGRSERMGAATCRGSRIPSGPPVQGERSAPRFGVTSLSLWPADFKNW
RIAGSRQEVALAGEPADQQQTHLRLPYRQTLGYKEDTTNPVCGEPWVSEDLEMTRHWPWEVSLRMENEHVCGG
ALIDPSWVVTAAHCSQGTKEYSVLGTSLQPMNFSRALWVPVRDIIMHPKYWGAFIMGDVALVHLQTPVTFSEYV
QPICLPEPNFLKVGTCWVTGWSQVKQRFSGSTANSMLTPELQEAEVFIMDNKRCDRHYKKSEFPVWPLVLGDMI
CATNYGENLCYGDSSGGPLACEVEGRWILAGVLSWEKACVKAQNPVYTRITKYTKWIKQMSNGAFSGPCASACLLF
LCWPLQPQMGS

>SGPr446_1_SEQID_84
ILTPVCGRTPLRIVGGVDAEEGRWPWQVSVRTKGRHICGGTLVTATWVLTAGHCISRRFHYSVKMGDRSVYNNETSV
VSVQRAFVHPKFSTVTTIRNDLALLQLQHPVNFTSNIQIPICIQENFQVEGTRCWTGWTGKTPERGEKLAISEILQDV
DQYIMCYEECNKIQKALSSTKDVIIKGMVCGYKEQKDCSCQDSSGRLACEYNDTWVQVGVSWGIGCGR

>SGPr447_SEQID_85
MGARGALLALLARAGLGKPEACGHREIHALVAGVESARGRWPWQASLRLRRHRRCGGSLLSRRWVLSAAHCF
QKHYPSEWTVQLGELTSRPTPWNLRAYSSRYKVQDIIVNPDALGVLRNDIALLRASSVTYNAYIQPICIESSTFNFBH

FIG. 2K

RPDCWVTGWGLSPSGTLPPIYNLREAQVTILNNTRCNLYFEQPSRSMIWDMSMFCAGAEAGSDTCKGDSGGPL
VCDKDLWYQVGIVSWGMDCGQPNRPGVYTNISVYFHWIRRVMSHSTPRPNPSQLLLLLALLWAP

>SGPr432_1_SEQID_86

MGSTWGSPPGWRLALCLTGLVSLYALHVKAARDRDYRALCDVGTASCSTVFSSRWGRGFLVEHVLGQDSIL
NQSNSIFGIFYTLQLLLGLQAAQRACGQRPKPQEGNTVPGEWPWQASVRRQGAHICSGSLVADTWLTAAH
CFEKAATELNSWSVLGSLQREGSLPGAEEVGVAALQLPRAYNHYSQGSDLALLQLAHTHTPLCLPQPAHRFPF
GASCWATGWDQDTSAPGTLNLRRLISRPTCNCIYNQLHQRHLSNPARPGMLCGGPQPGVQGPCQDSDGGPVL
CLEPDGHVWQAGIISFASSCAQEDAPVLLTNTAAHSSWLQARVQGAFLAQSPETPEMSDEDESCVACGSLRTAGPQ
AGAPSPWPWEARLMHQQLACGGALVSEEAVALTAHCFIGRQAPPEWSVGLGTRPEEWGLKQLILHGAYTHPEGG
YDMAILLAQPVTLGASLRPLCLPYPDHLPDGERGWVLGRARPGAGISSLQTPVTLLGPRACSRLLHAAPGGDGSP
LPGMVCTSAVGELPSCEGLSGAPLVHEVRGTWFLAGLSFGDACQGPAPPAVFTALPAYEDWVSSLDWQVYFAEEP
EPEAEPGSCLANISQPTSC

>SGPr529_SEQID_87

MRAPHILHLSAASGARALAKLLPLLMAQLWAAEAALLPQNDTRLDPDAYGAPCARGSQPWQVSLFNGLSFHCAGVLV
DQSWVLTAAHCGNKPLWARVGDDHLLLLQGEQLRRTTRSVVHPKYHQSGPILPRTDEHDLMLLKLARPVVPGR
VRALQLPYRCAQPGDQCQVAGWGTTAARRVKYNKGLTCSSITILSPKECEVFYPGVVTNNMICAGLDRGQDPCQSD
SGGPLVCDETLQGILSWGVPYPCGSAQHPAVYTTQICKYMSWINKVIRSN

>SGPr428_1_SEQID_88

NVQCGHRPAFPNSSWLPPFHERLQVQNGECPWQVSIQMSRKHLGGSLHWWVLTAAHCFRRTLDDMAVNVTVV
MGTRTFSNIHSEKQVQKVIHKYKPPQLDSDLALLLATPVQFSNFKMPVCLQEEERTWDWCWMAQWVTTNGYD
QYDDLNMHLEKLRVQISRKECAKRVNQLSRNMICAWNPEGTNGQGPGEVGGPLVCQKKNKSTWYQLGIISWVGVC
GQKNMPGVYTELSNYLLWIERKTVLAGKPYKYEPPDSVYALLSPWAILLYFVMLLLS

>SGPr425_SEQID_89

MENMLLWLIFFTPGWTIDGSEMEWDFMWHLRKVPRIVSERTFHLTSPAFAEADAKMMVNTVCGIECQKELPTPSSEL
EDYLSYETVFENGTRTLTRVKVQDLVLEPTQNTTKGVSVRRKRQVYGTDSRFSILDKRFLTNFPFSTAVKLSGCSGIL
ISPOHVLTAACHVHDGKDYVKGSKLRLVGLLKMNRKSGGKKRRGSKRSRREASGGDQREGTREHLPERAKGGRRR
KKSGRGQRIAEGRPSFQWTRVKNTHIPKGWARGGMGDATLDYDALLEKRAHKKKYMELGISPTIKKMPGGMHFS

FIG. 2L

GFDNDRADQLVYRFCVSDESNDLLYQYCDAESGSTGSGVYLRDKDPDKKNWKRKIIAVYSGHQWVDVHGVQKDYN
VAVRITPLKYAQICLWIHGNDANCAYG

>SGPr548_SEQID_90

MGDPEGSAEWGKGIPVRRNLLTVDGISLCLEGSWWRQKGPASPGFHSLSLRLQPNPGPSSTMWLLLTLSFLLA
STAAQDGDKLLLEGDECAPHSQPWQVALYERGRFNCGASLISPHWVLSAAHCQSRFMRVRLGEHNLKRDRGPEQLR
TTSRVIPHRYEARSHRNDIMLLRLVQPARLNPQVRPAVLPTRCPPHGEACVSGWGLVSHNEPGTAGSPRSQVSLP
DTLHCANISIISDTSCKSYPGRLTNTMVCAGAEGRGAESCEGDSGGPLVCGGILQGIVSWGDPVPCDNTTKPGVYTKV
CHYLEWIRETMKRN

>SGPr396_SEQID_91

MPAGACFTLLLLLGISVCGQPVSRRWGGQDAAAGRWPQVSLHFDHNFICGGSLVSERLILTAACHCIQPTWTTFS
YTVWLGSITVGDSTRKRVKYVYSKIVHPKYQDTTADVALLKSSQVTFTSAILPICLPSVTKQLAIPFCWVTGWGKVKE
SSDRDYHSALQEAEPIDRQACEQLYNPIGIFLPALEPVIKEDKICAGDTQNMKDSCKGDSGGPLSCHIDGWVIQTGV
VSWGLEGCKSLPGVYTNVYYQKWINATISRANNLDFSDFLPVILLSALLRPSCAFGPNTIHRVGTVAEAVACIQGWE
ENAWRFSPRGR

>SGPr426_SEQID_92

MMYAPVEFSEAEFSRAEYQRKQFWDVSVRLALFTLAINAIIAIGIVTHFWEDDKSFYYLASFKVTNIKYKENYGIRSS
REFIERSHQIERMMSRIFRHSVGGGRFIKSHVIKLSPEDEQGVLDILVLIIFRYPPSTDSAEQIKKKIEKALYQSLKTKQLSLTIN
KPSFRLTRCGIRMTSSNMPLPASSSTQRIVQGRETAMEGEWPWQASLQLIGSGHQCGASLISNTWLLTAACHCFWKNK
DPTQWIATFGATITPPAVKRNVRKIILHENYHRETNENDIALVQLSTGVEFSNIVQRVCLPDSSIKLPPKTSVFTGFGSI
VDDGPIQNTLRQARVETISTDVCNRKDVYDGLITPGMLCAGFMEGKIDACKGDSGGPLVYDNHDIWYIVGIVSWGQSC
ALPKKPGVYTRVTKYRDWIAKTGM

>SGPr552_SEQID_93

RIAEGLDAEEGEWPWQASLPQNNVYRRGATWLSNSWLITAACHCFIRVHDPKEWNVLSPQTQSNIKNVIQENYHYYP
AHDNDIAV/VHLSSPVL/YSNIQKACLPDVNIYFLYNSEAVVTAWGSFKPLRTTNSNVLHKGLVKIIDNRTCNNGEADGRVI
TSGMLCAGFLEPRVDACQGDGSGGPLVGTDSKILAKGSLLVKAGVNERALPNKPSVYTVQVTTY

BNSDOCID: <WO__0200B60A2_I_>

FIG. 2N

PLQTLGLEELLQRGIRARLREHGISLAAYGTIVSAELTGRHKGLAERDFKSGRCPGNSFSCGNSQCVTKNPEC
 DDQEDCSGSDGAHCECGLPAPWRMAGRIVGGMEASPGFEPWQASLRENKEHFCAAINARWLVSAAHCFNEFQ
 DPTKWAVYGATYLSGSEASTVRAQVQIVKHPLYNADTADFVAVLELTSPLPFRHIQPVCLPAATHIFPPSKKCLIS
 GWGYLKEDFRKHLPRPAMVKPEVLQKATVELLDQALCASLYGHSLTDRMVCAGYLDGKVDSCQDGGGGLVCEEPS
 GRFFLAGIVSWGIGCAEARPGVYARVTRLRDWILEATTKASMPAPTMAPAPAAPSTAWPTSPESPVVSTPTKSMQ
 ALSTVPLDWVTVPKLQECGARPAMEKPTRVVGFGAASGEVPWQVSLKEGSRHFCEGATVGDRLWLLSAAHCFNHT
 KVEQVRAHLGTASLLGLGSPVKIGLRRVWLHPLYNPGILDFDLAVLELASPLAFNKYIQPVCLPLAIQKFPVGRKCMIS
 GWGNTQEGNATKPELLQKASVGIDQKTCVLYNFSLTDRMICAGFLEGVKVDSCQDGGGGLACEEAPGVFYLAGIVS
 WGIGCAQVKKPGVYTRITRLKGWILEIMSSQPLMSPSTTRMLATTSPRTTAGLTPGATPSRPTPGAASRVGTGQPA
 NSTLSAVSTTARGQTPFPDAPEATHTQLPDCGLAPAALTRIVGGSAAGRGEWPWQVSLWLRREHRCGAVLVAER
 WLLSAAHCFDVYGDPKQWAAFLGTPFLSGAEGQLERVARIYKHPFYNYLTDYDVALLELAGPVRRLVRPICLPEP
 APRPPDGTTRCVITGWGVSREGGSMARQLQKAAVRLTSEQTCRRFYPVQISSRMLCAGFPQGGVDSCSGDAGGPLA
 CREPSGRWWLTGVTSWGYGCGRPHFPGVYTRVAAVRGWIGQHIQE

>SGPr521_SEQID_98

MARSLPLQLILLSLALETAGEEAQDKIDGAPCARGSHPWQVALLSGNQLHCGGVLVNERWWLTAACHCKMNEYTV
 HLGSDTLGDRRAQRIKASKSFRHPGYSTQTHVNDLMLVKLNSQARLSSMVKKVRLPSRCEPPGTCTVSGWGTTTS
 PDVTFPSDLMCVDVKLISPQDCTKVYKDILLENSMLCAGIPDSKKNACNGDSGGPLVCRGTLQGLVSWGTFPCGQPN
 DPGVYTQVCKFTKWINDTMKKHR

>SGPr530_1_SEQID_99

VSTVCGKPKVVGKIYGGRDAAAGQWPWQASLLYWGSHLCGAVLIDSCWLVSSTTHCFLNKSQAPKNYQVLLGNIQLY
 HQTQHTQKMSVHRIITHPDEFKLPFGSDIAMLQLHLP MNFTSYIVPVCLPSRDMQLPSNVSCWITGWGMLTEDHKRV
 QLSPPFYLQEGKVGLIENTLNTLYGQRTAKARPKLCTRRCCVGGYFSTGKSICKGDSGGPLVCYLPSAWVLVGLAS
 WGLDCRHPAYPSIFTRVITYFINWIDEIMRLTPLSDPALAPH

>SGPr520_SEQID_100

MLLAVLILLPLPSSWFAHGHPLYTRLPPSALQVFTLLGAETVLGRNLDYVCEGPCGERRPSTANVTRAHGRIVGSSA
 APPGAWPWLVRQLGGQPLCGGVLVAASWVLTAAHCFVGCSTRSAPNELLWTVTLAEGSRGEQAEVVPVNRILPH
 PKFDPRTFNDLALVQLWTPVSPGGSARPVCLPQEPQEPAGTACAIAGWGALFEDGPEAEAREARVPLLSTDTCR
 RALGPGLRPSTMLCAGYLAGGVDSQDGGGGLTCESEPGRPREVLFGVTSWGDGCGEPKPGVYTRVAVFKDWL
 QEQMSAASSSREPSCRELLAWDPPQELQADAARLCAFYARLCPGSQGACARLAHQQCQLQRRRRCELRLAHTLLGL

FIG. 20

LRNAQELLGPRPGLRRAPALAPALRESPLHPARELRHSGSRAAGTRFPKRRPEPRGEANGCPGLEPLRQKLA
ALQGAHAWILQVPSEHLAMNFHEVLADLSKTLTGLFRAWVRAGLGGRHVAFSGLVGLEPATLARSLPRLLVQALQA
FRVAALAEGEPEGPWMDVGQGPGLERKGGHPLNPQVPPARQP

>SGPr455_SEQID_101

MSPDIALLYLKHVKFGNAVQICLPDSDDKVEPGILCLSSGWGKISKTSYNSVLQEMELPIMDDRACNTVLKSMNLP
PLGRMTLCAGFPDWGMDACQDGGPLVCRGGGWLAGITSWVAGCAGGSPVVRNNHVKASLGIFSKVSELMDF
ITQNLFTGLDRGQPLSKVGSRYITKALSSVQEVNGSQRDKIILKFTSLDMEKQVGDHLYSLRSSSGVLFSKVCCKIL
PSPLLAETSEAMVPFVSDTEDSGSGFELTVTAVQKSEAGSGCSLAILVEEGTNHSKYPDLYPSNTRCHWFICAPEK
HIIKLTFFEDFAVKFSPNCIYDAVVIYGDSEEHKLA LCGMLTITSIFSSNMVTVIYKSDGKNRLQGFKARFTILPSES LN
KFEPLPPQNNPVSTVKAILHDVCGIPFPSPQWLSRRIAGGEEACPHCWQVGLRFLGDYQCGGAINPVWILTA AH
CVQLKNNPLSWTIIAGDHDRLNKESTEQVRRAKHIVHEDFNTLSYDSIALIQLSSPLEYNSVRPVCLPHSAEPLFSS
EICAVTGWGSAELSINVSSLDGGLASRLQQIQVHVLEREVECHTYTSAHPGGITEKMICAGFAASGEKDFCQGD SG
GPLVCRHENGPFVLYGIVSWGAGCVQWPWPGVFARVMIFLDWIOQKINGKLFNSVIKTTISFFRVGLGTVSCCSEAELE
KPRGFFPTPRYLLDYRGRLECSWLVRSASSMAKFTI EYLSLLGSPVCQDSVLIYEERHSKRKTAGGLHGRRLYSMT
FMSPGPLVRVTFHALVRGAFGISYIVLVLPKPKDSKITRLSQSSNREHLVPCEDVLLTKPEGIMQIPRNSHRTT MGCQW
RLVAPLNHIIQLNIINFPMKPTTFVCHGHLRVYEGFGPGKKLIGRMLMSTELSWFLSQFSTKKT TASCGETA VSMKMMY
TSIFLALQNTCYHALPHEVVLRIK

>SGPr507_2_SEQID_102

MKYVYFYLGVLAGTFFADSSVQKEDPAPYL VYLKSHFNPCVGVLIKPSWVLAPAHCYLPNLKVMLGNFKSRVRDGT E
QTINPIQIVRYWNYSHSAPQDDMLIKLAKPAMLNPKVQPLTATTNVRPGTVCLL SGLDWSQENSGLWQLEPPGHLT
LHRGPAIPDWQRHNSHEQGRHPDLRQNLEAPVMSDRECQKTEQGKSHRNSLCVKFVKVFSRIFGEVAVATVICKDKL
QGIEVGHFMGGDVGIYTNVYKYVSWIENTAKDK

>SGPr559_SEQID_103

MGENDPAPAVEAPFSRSLFGLDDLKISPVAPDADAVAAQILSLLPLKFFPIIIGIILALAILGLHIFDCSGKYRCRSSF
KCIELIARCDGVSDCKDGEDEYRCVRVGGQNAV LQVFTAASWKTMCSDDWKGHYANVACAQLGFP SYVSSDNL RVS
SLEGQFREEFVSIDHLLPDDKV TALHHSVYVREGCASGHVVTLQCTACGHRRGYSSRIVGGNMSLLSQWPWQASLQ
FQGYHLCGGSVITPLWIITAAHCYVDLYLPKSWTIQVGLVSLDNPAPSHLVEKIVYHSKYKPKRNLGNDIALMKLAGPLT
FNEMIQPVCLPNSEENFPDGVKVCWTSWGATEDGAGDASPVLNHAAVPLISNKICNHRDVYGGIISPSMLCAGYL TG
GVDSQCQDSSGGPLVCQERRLWKLVGATSF GIGCAEVNKPGVYTRVTSFLDWIHEQMERDLKT

FIG. 2P

[illegible]

>SGPr479_1_SEQID_105
MAAPASVMGPLGPSALGLLLLLVAPPRVAALVHRQPENQGISTGSVACGRPSMEGKILGGVPAPERKWPWQVSV
HYAGLHVCGGSILNEYWVLSAAHCFHRDKNIKIYDMYVGLVNLRVAGNHTQWYEVNRVILHPTYEMYHPIGGDVALVQ
LKTRIVFSESVPVCLATPEVNLTSANCWATGWGLVSKQGETSDELQEVQLPILEPWCHLLYGHMSYIMPDMLCAGD
ILINAKTVCEDSGGPLVCEFNRSWLQIGIVSWGRCGNPLYPGVYASVSFYFSKWICDNIEITPTPAQPAPALSPALGPT
LSVLMAMLAGWSVL

>SGPr489_1_SEQID_106
MSLKMLISRNKLLLGIVFFERKGSATLSLPKAPSCGGQSLVKVQPNYFNIFSRILGGSQVEKGSYPWQVSLKQRQKH
ICGGSIVSPQWVITAACHIANRNVSTLNTAGEYDLSDTDPGEQTLTIETVIHPHFSTKPKMDYDIALLKMAGAFQFGH
FVGPICLPELREQFEAGFICTAGWGRLTEGGVLSQVLQEVNLPILTWEECVAALLTLKRPISGKTFLCTGFPDGGGRDA
CQGDSSGSLMCRNKKGAWDGWSIWEAQVGGLESRSSRPSLGNKVRLCLTNFFKKLAGCGTWCSEQDVIIVSGA
EGKLHPPESLHLYYESKQRCVWTLVPEEMHVLLSFSLHDVESCHHSYLSMYSLEDRIPIGKFCGESLPSSILIGSNLSR
LKVFSDATDYAAGFNLTYKALKPNYIPGCSYLTVLFEGLIQSLNYPENYSDKANCDWIFQASKHHLIKLSFQSLIEES
GDCTSDYVTVHSDVERKKEIARLCGYDVPTPVLSPSSIMLISFHSDENGTCRGFQAIVSFIPKAVPYDNLNISISEDESMFL
ET

>SGPr465_1_SEQID_107
RWPWQASLLYLGHCIGAALDSNWASAAHCFQRCIFPPRAPLSTNPDSYRILLGYDQQSHPTTEHSKQMTVNKIMV
HADYNELHRMGSDITLLQLHHHVEFSSHILPACLPETTWLAPDSSCWISGWMVTEDEVFLPEPFLQAEAEVGVMDN
TVCGSFFQPPQPGQPSSDYTIHEDMLCAGDLITGKAICRVNSRGPLVCPLNGTWFLMGLSSWSLDCDCSPVGPVRVFT
RLPYFTNWISQKKRESTPPDPALAPPQETPPALDSMTSQGIVHKPGLCAALLAAHMFLLLLILLGSL

FIG. 2Q

>SGP524_1_SEQID_108
 MDKENSVDVSAAPADLKISNISQVWSAQKKLPVRRPPLPGRRLPLGRRPPQRPQIGKAKPKKQSKKKVPFWNVQNKIIL
 FTVFLFILAVIAWTLLWLYISKTESKDAFYFAGMFRITNIEFLPEYRQKESREFLSVRTVQQVINLVYTTSAFSKFYEQS
 WVADVSSNNKGGLLVHFVWVFMPRAKGHIFCEDCVAAILKDSIQTSINRTSVGSLQGLAVDMDSVNLNGDCWSFLKK
 KKRKENGAVSTDKGCSQFYAEHLSLHYPLEISAASGRLMCHFKLVAINGYLIRLSIKSIQIEADNCVTDLSLTYYDSLLPIR
 SSILYRICEPTRTLMFVSTNNLMLVTFKSPHIRRLSGIRAYFEVIPEQKCENTVLVKDITGFEKGISSPYPSYPPKCK
 CTWKFTSLSTLGIALKFYNYISITKSMKGCEHGWWNEHEMYCGSYMMDHQTFIRVPSPLVHIQLQCSSLSDKPLLA
 EYGSYNISQPCPVGSFRCSGLCVPOAQRCDGVNDCFDESELCVSPQACNTSSFRQHGPLICDGFRCDCENGRD
 EQNCTQSI PCNNRTFKCGNDICFRKQNAKCDGTVDPCPDGSEEGCTCSRSSSALHRIIGGTDLTLEGGWPWQVSLHF
 VGSAYCGASVISREWLLSAAHCFHGNRLSDPTPWTALHGMVYQGNKAFVSPVRRIVVHEYNSQTFDYDIALQLLSIA
 WPETLKQLIQIPICPTGQVRVSCEKWCWTGWGRRHEADNKGSLVQQAEVELIDQTLCVSTYGIITSRMLCAGIMSG
 KRDAKGDGSGGPLSCRRKSDGKWILTGIVSWGHCGRPNFPGVYTRVSNFVFWIHKYVPSLL

>SGP422_SEQID_109
 MTLNKIKDLFAGKGQWDLAPEAEMLKPMWIAVLVLSLTVAVTIGLLVHFLVFDQKEYYHGSFKILDQPINNFGQSN
 TYQLKDLRETTENLVYSLKMYLSFVCHSPEEDGVKVDVMVFQFPSTEQRAVREKKIQSILNQKIRNLRALPINASSVQV
 NVAMVKNGNVGP GSGAGEAPGLGAGPAWSPMSSSTGELTVQASCGKRVPLNVNRIASGVIAPKAAWPWQASLQY
 DNIHQCGATLISNTWLVTAAHCFQKYKNPHQWTVSFGTKINPPLMKRNVRRFIIHEKYRSAAREYDIAVWQVSSRVTF
 DDIRQICLPEASASFPNLTVHITGFGALYGGESQNDLREARVKIISDDVCKQPQVYGNDIKPGMFCAGYMEGIYDAC
 RGDGSGPLVTRDLKDTWYLGIVSWGDNCGQKDKPGVYTQVTYYRNWIAASKTG

>SGP538_SEQID_110
 MSLMLDDQPPMEAAEQAEEGPGIFRAEPGDQQHPISQAVCWRSRMRGCAVLGALLAGAGVGSWLLVLYLCPA
 ASQPISGTLQDEEITLSCSEASAEALLPALPKTVSFRINSEDFLEAQVRDQPRWLLVCHEGWSPALGLQICWSLGH
 RLTHHKGVNLTDIKLNSSQEFQALSPRLGGFLEAAWQPRNNCTSGQVSLRCSECGARPLASRIVGGQSVAPGRWP
 WQASVALGFRHTCGSVLAPRWVWVTAACHMHSFRLARLSSWRVHAGLVSHSAVRPHQCALVERIIPHPLYSAQNHD
 YDVALLRLQ TALNFSDTVGAVCLPAKEQHFPKGSRCWVSGWGHTHPSHTYSSDMLQD TVPLFSTQLCNSSCVYSG
 ALTPRMLCAGYLDGRADACQGDGSGGPLVCPDGDGTWRLVGVVSWGRACAEPNHPGVYAKVAEFLDWIHDTAQDSSL

FIG. 2R

>SGPr527_1_SEQID_111
 MARHLLPLVLMVISPIGAFQDSALSPTQEEPEDDCGRPEPSARIVGGSNAQPGTWPWQVSLHHGGGHICGGSLIA
 PSWVLSAAHCFMTNGTLEPAAEWSVLLGVHSQDGPLDGAHTRAVAAIWPNANYSQVELGADLALLRLASPASLGPV
 WPVCLPRASHRFVHGTAACWATGWGDVQADPLPLPWVLEVELRLGEATCCLYSQPGPNLTQLPGMLCAGY
 PEGRRDTCQDGGGLVCEEGRWFQAGITSFGFGCGRRNRPGVFTAVATYEAWIREQVMGSEPGPAFTQPQKT
 QSDPQEPREENCTIALPECGKAPRPGAWPWEAQVMVPGSRPCHGALVSESWVLAASCFLDPNSSDSPPRDLDAW
 RVLLPSRPAERVARLVQHENASWDNASDLALLQLRTPVNLASAARPVCLPHEHYFLPGSRCRLARWGRGEPALG
 PGALLEAELLGWWCHCLYGRQGAAPPLPGDPPHALCPAYQEKEEVGSCWTHGPWISHVTRGAYLEDQLAWDWG
 PDGEETETQTCPPHTEHGACGLRLEAAPVGLWPWLAEVHVAGDRVCTGILLAPGWVLAATHCVLRPGSTTVPYIEV
 YLGRAGASSLPQGHQVSRVISIRLPQHLGLRPPLALLELSSRVEPSALPICLHPAGIPPGASCWVLGWKEPQDRVP
 VAAVSILTQRICDCLYQGILPPGTLCVLYAEGQENRCMTSAPLLCQMTGSWILVGMVAVQGSRELF A AIGPEEAWI
 SQTVGEANFLPPSGSPHWPTGGSNLCPPELAKASGSPHAYVFLLLTLLIQS

>SGPr542_SEQID_112
 AMGLGLRGWGRPLLTATALMLPVKPPAGSWGAIQGGHEVTPHSRPMASVRFGGQHCGGFLLRARWVWSAAH
 CFSHRDLRTGLVLGAVHLSAEPTQQVFGIDAL TTHPDYHPMTHANDICLLQLNGSAVLGPVGLRLPGRRARPT
 AGTRCRVAGWGVSDFEELPPGLMEAKVRVLDPDVNCNSSWKGHLLTLMCTRSGDSHRRGFCSDSGGGLVCRNR
 AHGLVSFSGLWCGDPKTPDVYTQVSAFVAWWDVRRSSPQPGPLPGTTRPPGEAA

>SGPr551_SEQID_113
 MPVAEAPQVAGQGGDGGEEAEPEGMFKACEDSKRKARGYRLVPLFVLLALLVLSAGVLLWYFLGYKAEVMVS
 QVYSGSLRVLNRHFSQDLTRRESSAFRSETAKAQMKLKITSTRLGTYYNSSSVYSGEGPLTCFFWFILQIPEHRRL
 MLSPEVQALLVEELLSTVNSSAAVPYRAEYEVDPGLVILEASVKDIAALNSTLGCYRYSYVGGQVLRKGPDLHAS
 SCLWHLQGPDKMLKLRLWTLAECDRLAMVDVAGPLEKRLITSYGCQRQEPVVEVLASGAIMAVVWKKGLHSYY
 DPFVLSVQPWFQACEVNLTDNRDLSQGLSTPYFSPYSPQTHCSWHLTVPSLDYGLALWFDAYALRRQKYDLPC
 TQQQWTIQNRRLCGLRILQPYAERIPVATAGITINFTSQISLTGPGVRVHYGLYNQSDPCPGFEFLCSVNGLCVPACDG
 VKDCPNGLDERNCVCRATFQCKEDSTCISLPKVCDGQPDCLNGSDEEQCEGVPCGTFTFQCEDRSCVKKPNPQC
 DGRPDCRDGSEEHCDCLGQGPSSRIVGAVSSEGEWPWQASLQVRGRHICGGALADRWWVITAACHCFQEDSMAS
 TVLWTVFLGKVVQNSRWPGEVSKVSRLLHPYHEEDSHDYDVALQLDHPVWRSAAVRPVCLPARSHFFEPGLHC
 WITGWGALREGGPISNALQKVDVQLIPQDLCSEAYRYQVTPRMLCAGYRKGGKDACQGDGSGGLVCKALSGRWFLA
 GLVSWGLGCGRPNYFGVYTRITGVISWQQVVT

FIG. 2S

>SGPr451_SEQID_114
DLPPSCSPASKMRLGLLSVALLFVGSSHLYSDHYSPSGRHRLLGPSPEPAASSQQAEEAVRKRLRRRREGGAHAKDCG
TAPLKDVLGSRIGGTEAQAGAWPWVSLQIKYGRVLHVCGGTLVRERWVLTAAHCTKDTSDPLMWTAVIGTNNIH
GRYPHTKKIKAIHPNFILESYNDIALFHLKAVRYNDYIQPICLPFDVFQILDGNTKCFISGWGRTKEEGNATNILQD
AEVHYSREMCNRSERYGGIIPNTSFCAGDEDEGAFDTCRGDSGGPLMCYLPEYKRFFVMGITSYGHGCGRRGFPGVY
IGPSFYQKWLTEHFFHASTQGILTINILRGQILIALCFVILLATT

>SGPr452_1_SEQID_115
SPPQPRTPDCRLQASLEALATLAPQPSDWLCFADLWFEADGAHSMGLGSSLKWAWAKPSGMPVPENDLVGIVG
GHNAPPGKWPWQVSLRVSYHWASWAHICGSLIHPQWVLTAAHCIFWKDTPDSYRIHAGDVYLYGGRGLLNVSR
IVHPNYVTAGLGADVALLQLPGSPLSPESLPPYRLQQASVQVLENAVCEQPYRNASGHTGDRQLILDMLCAGSEG
RDSCYGDSSGGLVCRLRGSRWRLVGWSWGYGCTLRDFPGVYTHVQIYVLWILQQVGELP

>SGPr504_SEQID_116
IIGGHEVTPHSRPMASVRFGGQHHC GGFLLRARWVWSAAQCFSH

>SGPr469_SEQID_117
GDSSGGPLVCELNGTWVQVGIWSWGIGCGRKGYPGVYTEVSFYKKWI

>SGPr400_SEQID_118
MAGEQVTANVSRYPGQKTMSPFECTFLLSYRASLLAVWTHRSNNSRGRAFESQVLPDLTAGDAADPPPLPGGAAL
LKSGPFRIWQGVKTKGEEGDRDTGTAGYAFTLLLLLGISGEPPWVCGRPTVSSGIASGLGASVGQWPWQVSIRQGL
IHVCSDTLISEEWLTVaicfPLSPHPDFQANTSSAIAVVELSPVSVSPVLLICLPSSEVYLLKNTTSCWWTGWYTG
FYIKRSYTLKELKVPLIDLQTCGDHYQNEILLHGVELIIEAMICSKLPVGQMDQCTVRIHPSGTFRPCLPQ

THIS PAGE BLANK (USPTO)